

DAFTAR PUSTAKA

- Ali, A., Bhan, C., Malik, M.B., Ahmad, M.Q. & Sami, S.A. 2018. The Prevention and Management of Contrast-induced Acute Kidney Injury: A Mini-review of the Literature. *Cureus*, 10(9): 1–6.
- Andreucci, M. & Faga, T. 2014. Prevention of contrast-induced nephropathy through a knowledge of its pathogenesis and risk factors. *Sci. World J.*, 2014: 16.
- Andreucci, M., Faga, T., Pisani, A., Sabbatini, M. & Michael, A. 2014. Acute Kidney Injury by Radiographic Contrast Media : Pathogenesis and Prevention. *Biomed Res. Int.*, 2014: 21.
- Andreucci, M., Faga, T., Pisani, A., Sabbatini, M., Russo, D. & Michael, A. 2014. The Choice of the Iodinated Radiographic Contrast Media to Prevent Contrast-Induced Nephropathy. *Advances in Nephrology*, 2014: 1–11.
- Bansal, N., Tighiouart, H., Weiner, D., Griffith, J., Vlagopoulos, P., Salem, D., Levin, A. & Sarnak, M.J. 2007. Anemia as a Risk Factor for Kidney Function Decline in Individuals With Heart Failure. *Am. J. Cardiol.*, 99(8): 1137–1142.
- Becker, J., Babb, J. & Serrano, M. 2013. Glomerular Filtration Rate in Evaluation of the Effect of Iodinated Contrast Media on Renal Function. <http://dx.doi.org/10.2214/AJR.12.8871>, 200(4): 822–826. www.ajronline.org 1 October 2021.
- Bolignano, D., Coppolino, G. & Campo, S. 2008. Urinary neutrophil gelatinase-associated lipocalin (NGAL) is associated with severity of renal disease in proteinuric patients. *Nephrol Dial Transplant*, 23(416): 414–416.
- Bolinska, S., Sobkowicz, B., Zaniewska, J., Chlebińska, I. & Boliński, J. 2011. The significance of anaemia in patients with acute ST – elevation myocardial infarction undergoing primary percutaneous coronary intervention. *Kardiol Pol*, 69(1): 33–39.
- Brunzel, N.A. 2018. *Fundamental of Urine and Body Fluid Analysis*. Fourth edi. Elsevier.
- Cai, L., Rubin, J., Han, W., Venge, P. & Xu, S. 2010. The origin of multiple molecular forms in urine of HNL/NGAL. *CJASN*, 5(12): 2229–2235.
- Caldararu, C., Dogaru, M.T. & Vari, C. 2011. Comparison Between Cockcroft-Gault and MDRD Formula for The Estimation of Glomerular Filtration Rate in Chronic Kidney Disease Patients. *Farmacia*, 59(November).
- Caspi, O., Habib, M., Cohen, Y., Kerner, A., Roguin, A. & Abergel, E. 2017. Acute Kidney Injury After Primary Angioplasty : Is Contrast-Induced. *J Am Heart Assoc*: 1–11.
- Chen, W., Chen, P., Ni, Z., Liu, Y., Guo, W., Jiang, L., Wei, X., Chen, J., Tan, N., He, P. & Guo, Y. 2020. Prognostic of different glomerular filtration rate formulas in patients receiving percutaneous coronary intervention : insights from a multicenter observational cohort. : 1–10.
- Chorin, E., Ben-Assa, E., Konigstein, M., Rofe, M.T., Hochstadt, A., Galli, N., Schnapper, M., Arbel, Y., Rabey, I., Shoshan, J. Ben, Halkin, A., Herz, I., Finkelstein, A., Bazan, S., Keren, G. & Banai, S. 2016. Prevention of post procedural acute kidney injury in the catheterization laboratory in a real-world population. *IJCA*, 226: 1–30. <http://dx.doi.org/10.1016/j.ijcard.2016.10.028>.
- Decavele, A.S.C., Dhondt, L., De Buyzere, M.L. & Delanghe, J.R. 2011. Increased urinary neutrophil gelatinase associated lipocalin in urinary tract infections and leukocyturia. *Clin Chem Lab Med*, 49(6): 999–1003.
- Faucon, A.L., Bobrie, G. & Clément, O. 2019. Nephrotoxicity of iodinated contrast media:

- From pathophysiology to prevention strategies. *Eur. J. Radiol*, 116: 231–241. <https://doi.org/10.1016/j.ejrad.2019.03.008>.
- Filiopoulos, V., Biblaki, D. & Vlassopoulos, D. 2014. Neutrophil gelatinase-associated lipocalin (NGAL): A promising biomarker of contrast-induced nephropathy after computed tomography. *Ren Fail*, 36(6): 979–986.
- Geus, H.R.H. De, Bakker, J., Lesaffre, E.M.E.H. & Noble, J.L.M.L. 2011. Neutrophil Gelatinase-associated Lipocalin at ICU Admission Predicts for Acute Kidney Injury in Adult Patients. *Am J Respir Crit Care Med*.
- Haase, M., Devarajan, P., Haase-Fielitz, A., Bellomo, R., Cruz, D.N., Wagener, G., Krawczeski, C.D., Koyner, J.L., Murray, P., Zappitelli, M., Goldstein, S.L., Makris, K., Ronco, C., Martensson, J., Martling, C.R., Venge, P., Siew, E., Ware, L.B., Ikizler, T.A. & Mertens, P.R. 2011. The outcome of neutrophil gelatinase-associated lipocalin-positive subclinical acute kidney injury: A multicenter pooled analysis of prospective studies. *JACC*, 57(17): 1752–1761.
- Halliwell, B. 1988. Albumin-An important extracellular antioxidant? *Biochemical Pharmacology*, 37(4): 569–571.
- Hanif, M.O., Bali, A. & Ramphul, K. 2021. Acute Renal Tubular Necrosis. *StatPearls*. <https://www.ncbi.nlm.nih.gov/books/NBK507815/> 3 October 2021.
- Hebert, S.A. & Molony, D.A. 2015. The CKD-EPI equation for EGFR predicted adverse outcomes after PCI better than other equations. *Ann. Intern. Med.*, 163(10): JC12.
- Heil, W. & Ehrhardt, V. 2008. *Reference Range for Adult and Children*. 9th ed. Germany: Roche Diagnostics GmbH.
- Inker, A. 2011. Frequently Asked Questions About GFR Estimates Table Of Contents. *National Kidney Foundation*.
- Kashani, K., Cheungpasitporn, W. & Ronco, C. 2017. Biomarkers of acute kidney injury: The pathway from discovery to clinical adoption. *Clin Chem Lab Med*, 55(8): 1074–1089.
- Kellum, J.A., Lameire, N., Aspelin, P., Barsoum, R.S., Burdmann, E.A., Goldstein, S.L., Herzog, C.A., Joannidis, M., Kribben, A., Levey, A.S., MacLeod, A.M., Mehta, R.L., Murray, P.T., Naicker, S., Opal, S.M., Schaefer, F., Schetz, M. & Uchino, S. 2012. Kidney disease: Improving global outcomes (KDIGO) acute kidney injury work group. KDIGO clinical practice guideline for acute kidney injury. *ISN*, 2(1): 1–138.
- Khilchuk, A., Abdulkarim, D., Vlasenko, S., Abugov, S., Scherbak, S., Popov, V., Vorobyovskii, D., Agarkov, M., Sarana, A. & Karmazanashvili, E. 2019. Novel Biomarkers of Contrast-Induced Kidney Injury after Endovascular Procedures: An Interventional Cardiologists Notion. *World J. Cardiovasc. Dis*, 09(12): 942–957.
- Koyner, J.L., Vaidya, V.S., Bennett, M.R., Ma, Q., Worcester, E., Akhter, S.A., Raman, J., Jeevanandam, V., O'Connor, M.F., Devarajan, P., Bonventre, J. V. & Murray, P.T. 2010. Urinary biomarkers in the clinical prognosis and early detection of acute kidney injury. *Clin J Am Soc Nephrol*., 5(12): 2154–2165.
- Kusirisin, P., Chattipakorn, S.C. & Chattipakorn, N. 2020. Contrast-induced nephropathy and oxidative stress: mechanistic insights for better interventional approaches. *J Transl Med*, 18(1): 1–35. <https://doi.org/10.1186/s12967-020-02574-8>.
- Lerman-Garber, I., Calva-Mercado, J.J., Martínez-Sibaja, C., Del Castillo, C.F., Sánchez-Javier, R.M., Lara, E., Villalobos, I. & Gómez-Pérez, F.J. 2000. Leukocyturia in women with diabetes and its clinical implications. *Archives of Medical Research*, 31(2): 210–215.
- Levey, A., Stevens, L. & Schmid, C. 2009. A New Equation to Estimate Glomerular Filtration Rate. *Journal of Nephrology & Therapeutics*, 06(05): 604–612.
- Levey, A.S., Coresh, J., Greene, T., Stevens, L.A., Zhang, Y. (Lucy), Hendriksen, S.,

- Kusek, J.W. & Lente, F. Van. 2006. Using Standardized Serum Creatinine Values in the Modification of Diet in Renal Disease Study Equation for Estimating Glomerular. *Ann. Intern. Med.*
- Levey, A.S. & Inker, L.A. 2016. GFR as the ‘gold Standard’: Estimated, Measured, and True. *Am. J. Kidney Dis.*, 67(1): 9–12.
- Lipsey, M., Furebring, M.I.A., Rubertsson, S. & Larsson, A. 2011. Significant differences when using creatinine, modification of diet in renal disease, or cystatin C for estimating glomerular filtration rate in ICU patients. *Ups. J. Med. Sci.*, 116(September 2010): 39–46.
- Macedo, E. & Mehta, R.L. 2013. Measuring renal function in critically ill patients: tools and strategies for assessing glomerular filtration rate. *Curr Opin Crit Care*., 19(6): 560–566.
- Mangan, C. 2018. Renal physiology: blood flow, glomerular filtration and plasma clearance. *Anaesth. Intensive Care Med.*, 19(5): 254–257.
- Mardhika, A.S., Somasetia, D.H. & Wulandari, D.A. 2019. Korelasi antara Kadar Neutrophil Gelatinase Associated Lipocaline Urin dengan Laju Filtrasi Glomerulus pada Variasi Waktu untuk Mendeteksi Dini Gangguan Ginjal Akut pada Anak Sepsis. , 21(38): 1–7.
- Matsumoto, T. & Kumazawa, J. 1993. Urinary tract infection. *Nippon rinsho. Japanese journal of clinical medicine*, 51 Suppl: 492–502.
- McIlroy, D.R., Wagener, G. & Lee, T. 2010. Biomarkers of Acute Kidney Injury. *Anesthesiology*, 112(4): 998–1004.
- Murata, K., Baumann, N.A., Saenger, A.K., Larson, T.S., Rule, A.D. & Lieske, J.C. 2011. Relative performance of the MDRD and CKD-EPI equations for estimating glomerular filtration rate among patients with varied clinical presentations. *Clinical Journal of the American Society of Nephrology*, 6(8): 1963–1972.
- Myers, G.L., Miller, W.G., Coresh, J., Fleming, J., Greenberg, N., Greene, T., Hostetter, T., Levey, A.S., Panteghini, M., Welch, M. & Eckfeldt, J.H. 2006. Recommendations for Improving Serum Creatinine Measurement: A Report from the Laboratory Working Group of the National Kidney Disease Education Program. *Clinical Chemistry*, 18: 5–18.
- Ning, M., Mao, X., Niu, Y., Tang, B. & Shen, H. 2018. Usefulness and limitations of neutrophil gelatinase-associated lipocalin in the assessment of kidney diseases. *J Lab Precis Med*, 3(3): 1–10.
- Nunes, M.B.G., Filho, A.C., Alvares, V.R.C., Meneguz-Moreno, R., Lamas, E., Loures, V., Chamié, D. & Abizaid, A. 2018. CKD-EPI versus Cockcroft-Gault formula for predicting contrast-induced nephropathy following percutaneous coronary intervention in patients without significant renal impairment. *Rev Port Cardiol.*, 37(1): 25–33. <http://dx.doi.org/10.1016/j.repece.2017.05.010>.
- Nusca, A., Miglionico, M., Proscia, C., Ragni, L., Carassiti, M., Pepe, F.L. & Sciascio, G. Di. 2018. Early prediction of contrast-induced acute kidney injury by a ‘bedside’ assessment of Neutrophil Gelatinase-Associated Lipocalin during elective percutaneous coronary interventions. *PLOS ONE*, 13(5): 1–12.
- O’Callagan, C.. 2006. *At a Glance: Sistem Ginjal*. Edisi kedua. A. Safitri & R. Astikawati, eds. Erlangga.
- Oweis, A.O., Alshelleh, S.A., Saadeh, N., Jarrah, M.I., Ibdah, R. & Alzoubi, K.H. 2020. Long-Term Follow-Up of Contrast-Induced Acute Kidney Injury: A Study from a Developing Country. *Int. J. Vasc. Med*, 2020: 15–20.
- PERKI. 2016. *Panduan praktik klinis (ppk) dan clinical pathway (cp) penyakit jantung dan pembuluh darah*.

- Perrin, T., Descombes, E. & Cook, S. 2012. Contrast-induced nephropathy in invasive cardiology: Incidence, pathophysiology, diagnosis, prevention and prognosis. *Swiss Med Wkly.*, 142: 1–11.
- Pistolessi, V., Regolisti, G., Morabito, S., Gandolfini, I., Corrado, S., Piotti, G. & Fiaccadori, E. 2018. Contrast medium induced acute kidney injury : a narrative review. *J. Nephrol*, 31(6): 797–812. <http://dx.doi.org/10.1007/s40620-018-0498-y>.
- Quintavalle, C., Anselmi, C.V., De Micco, F., Roscigno, G., Visconti, G., Golia, B., Focaccio, A., Ricciardelli, B., Perna, E., Papa, L., Donnarumma, E., Condorelli, G. & Briguori, C. 2015. Neutrophil gelatinase-associated lipocalin and contrast-induced acute kidney injury. *Circ. Cardiovasc. Interv.*, 8(9): 1–11.
- Ronco, F., Tarantini, G. & McCullough, P.A. 2020. Contrast induced acute kidney injury in interventional cardiology: An update and key guidance for clinicians. *Rev. Cardiovasc. Med*, 21(1): 9–23.
- Saito, Y., Watanabe, M., Aonuma, K., Hirayama, A., Tamaki, N., Tsutsui, H., Murohara, T., Ogawa, H., Akasaka, T., Yoshimura, M., Sato, A., Takayama, T., Sakakibara, M., Suzuki, S., Ishigami, K., Onoue, K., Hoshi, T., Hiraya, D., Sakai, S., Maruta, S., Tsumagari, Y., Noguchi, Y., Kakefuda, Y., Watabe, H., Nakanishi, K., Otani, M., Kawabe, M., Hikita, H., Higuchi, Y., Nishida, T., Iida, K., Uematsu, M., Ishihara, T., Kanda, T., Ueda, Y., Nishimoto, Y., Arima, K., Mano, H., Oiwa, K., Matsumoto, M., Yahata, T., Mori, N., Kato, H., Nishino, M., Tachibana, E., Watanabe, Y., Yahagi, T., Yajima, J., Murata, N., Kikushima, K., Nagao, K., Nomoto, K., Akutsu, N., Asakawa, N., Inden, Y., Hayashi, M., Ishii, H., Yoshida, N., Hirai, K., Iwama, H., Kondo, H., Nakagawa, Y., Yamamoto, E., Soejima, H., Yamano, T., Ino, Y., Shiono, Y., Yamaguchi, T., Tamai, K., Inoue, F., Nakajima, T., Sugimoto, H., Horii, M., Yamamoto, Y., Takemoto, Y., Asoda, Y. & Komeda, M. 2015. Proteinuria and reduced estimated glomerular filtration rate are independent risk factors for contrast-induced nephropathy after cardiac catheterization. *Circ j*, 79(7): 1624–1630.
- Sancho-Martínez, S.M., Blanco-Gozaño, V., Quiros, Y., Prieto-García, L., Montero-Gómez, M.J., Docherty, N.G., Martínez-Salgado, C., Morales, A.I., López-Novoa, J.M. & López-Hernández, F.J. 2020. Impaired Tubular Reabsorption Is the Main Mechanism Explaining Increases in Urinary NGAL Excretion Following Acute Kidney Injury in Rats. *Toxicol. Sci*, 175(1): 75–86.
- Seeman, T., Vondrák, K. & Zieg, J. 2017. Urinary Neutrophil Gelatinase-Associated Lipocalin Does Not Distinguish Urinary Neutrophil Gelatinase-Associated Lipocalin Does Not Distinguish Acute Rejection from Other Causes of Acute Kidney Injury in Pediatric Renal Transplant Recipients. , (January).
- Silvain, J., Nguyen, L.S., Spagnoli, V., Kerneis, M., Guedeney, P., Vignolles, N., Cosker, K., Barthelemy, O., Le Feuvre, C., Helft, G., Collet, J.P. & Montalescot, G. 2018. Contrast-induced acute kidney injury and mortality in ST elevation myocardial infarction treated with primary percutaneous coronary intervention. *Heart*, 104(9): 767–772.
- Smertka, M. & Chudek, J. 2012. Using NGAL as an early diagnostic test of acute kidney injury. *Renal Failure*, 34(1): 130–133.
- Souza, D.F., Reis, S.S.A., Botelho, R. V. & Ferreira-Filho, S.R. 2015. Relative and Absolute Changes in Urinary Neutrophil Gelatinase-Associated Lipocalin and Correlation with Small Increases in Serum Creatinine Levels after Coronary Angiography: An Observational Study. *Nephron*, 129(2): 84–90.
- Thomas, G., Sehgal, A.R., Kashyap, S.R., Srinivas, T.R. & Kirwan, J.P. 2011. Metabolic Syndrome and Kidney Disease : A Systematic Review and Meta-analysis. *Clin J Am Soc Nephrol*, 6: 2364–2373.

- Torregrosa, I., Montoliu, C., Urios, A., Andrés-Costa, M.J., Giménez-Garzó, C., Juan, I., Puchades, M.J., Blasco, M.L., Carratalá, A., Sanjuán, R. & Miguel, A. 2015. Urinary KIM-1, NGAL and L-FABP for the diagnosis of AKI in patients with acute coronary syndrome or heart failure undergoing coronary angiography. *Heart Vessels*, 30(6): 703–711.
- Vijay, S., Tiwari, B. & Singh, A. 2013. Contrast induced nephropathy: Pathophysiology and prevention. *Heart India*, 1(2): 39.
- Vlachopoulos, G., Schizas, D., Hasemaki, N. & Georgalis, A. 2019. Pathophysiology of Contrast-Induced Acute Kidney Injury (CIAKI). *Curr. Pharm. Des.*, 25(44): 4642–4647.
- Waas, T., Schulz, A., Lotz, J., Rossmann, H., Pfeiffer, N., Beutel, M.E., Schmidtman, I., Münzel, T., Wild, P.S. & Lackner, K.J. 2021. Distribution of estimated glomerular filtration rate and determinants of its age dependent loss in a German population - based study. *Scientific Reports*, 11: 1–13. <https://doi.org/10.1038/s41598-021-89442-7>.
- Wang, X. 2021. Combined Effects of Dyslipidemia and High Adiposity on the Estimated Glomerular Filtration Rate in a Middle-Aged Chinese Population. *Diabetes Metab Syndr Obes*, 14: 4513–4522.
- Widaningsih, Y., Natsir, R., Arif, M., Bahrin, U., Kasim, H., Mangerangi, F., Kadir, N.A., Anggraeni, R., Haruna, N., Hatta, M. & Usman, A.N. 2017. Analysis of Risk Factors Contrast Induced Nephropathy in Coronary Artery Disease Patients Obtained Percutaneous Coronary Intervention at Wahidin Sudirohusodo Hospital in Makassar, South Sulawesi. *IJSBAR*, 33(1): 145–150. <http://gssrr.org/index.php?journal=JournalOfBasicAndApplied>.
- Woo, K.S., Choi, J.L., Kim, B.R., Kim, J.E., An, W.S. & Han, J.Y. 2012. Urinary neutrophil gelatinase-associated lipocalin levels in comparison with glomerular filtration rate for evaluation of renal function in patients with diabetic chronic kidney disease. *Diabetes Metab J*, 36(4): 307–313.
- Yim, H.E. 2015. Neutrophil Gelatinase-Associated Lipocalin and Kidney Diseases. *Child Kidney Dis*, 19(2): 79–88.