

DAFTAR PUSTAKA

- Adriany, D. T., A. A. B. dan M. I. B. (2020). Perbandingan Metode Isolasi DNA Terhadap Nilai Kemurnian DNA untuk Pengujian White Spot Syndrom Virus (WSSV) pada Lobster Bambu (*Panulirus versicolor*). *Jurnal Prosiding Simposium Nasional Kelautan Dan Perikanan*, (7), 187–194.
- Al-Harhi, H., Syed, A., & El-Enshasy, H. A. (2019). Production, purification and evaluation of biodegrading potential of PHB depolymerase of *Stenotrophomonas* sp. RZS 7. *PLoS ONE*, 15(1).
- Bugnicourt, E., Cinelli, P., Lazzeri, A., & Alvarez, V. (2014). Polyhydroxyalkanoate (PHA): Review of synthesis, characteristics, processing and potential applications in packaging. 8(11), 791–808.
- Christova, N., Kabaivanova, L., Nacheva, L., Petrov, P., & Stoineva, I. (2019). Biodegradation of crude oil hydrocarbons by a newly isolated biosurfactant producing strain. *Biotechnology and Biotechnological Equipment*, 33(1), 863–872.
- Claridge, J. E. (2004). Impact of 16S rRNA gene sequence analysis for identification of bacteria on clinical microbiology and infectious diseases. *Clinical Microbiology Reviews*, 17(4), 840–862.
- Daane, L. L., Harjono, I., Barns, S. M., Launen, L. A., Palleroni, N. J., & Häggblom, M. M. (2002). PAH-degradation by *Paenibacillus* spp. and description of *Paenibacillus naphthalenovorans* sp. nov., a naphthalene-degrading bacterium from the rhizosphere of salt marsh plants. *International Journal of Systematic and Evolutionary Microbiology*, 52(1), 131–139.
- Goudarztalejerdi, A., Tabatabaei, M., Eskandari, M. H., & Mowla, D. (2019). Optimization Conditions for Maximum Oil Bioremediation and Biopolymer Production by *Pseudomonas*. *Iranian Journal of Science and Technology, Transaction A: Science*, 43(4), 1439–1446.
- Green, M. R., & Sambrook, J. (2018). Isolation and Quantification of DNA. *Cold Spring Harbor Protocols*, 403–414.
- Janda, J. M., & Abbott, S. L. (2007). 16S rRNA gene sequencing for bacterial identification in the diagnostic laboratory: Pluses, perils, and pitfalls. *Journal of Clinical Microbiology*, 45(9), 2761–2764.
- Kamnev, A. A., Dyatlova, Y. A., Kenzhegulov, O. A., Vladimirova, A. A., Mamchenkova, P. V., & Tugarova, A. V. (2021). Fourier transform infrared (FTIR) spectroscopic analyses of microbiological samples and biogenic selenium nanoparticles of microbial origin: Sample preparation effects. *Molecules*, 26(4).
- Kansiz, K., Billman-Jacobe, H., & McNaughton, D. (2000). Quantitative determination of the biodegradable polymer poly(β -hydroxybutyrate) in a recombinant *Escherichia coli* strain by use of mid-infrared spectroscopy and multivariate statistics. *Applied and Environmental Microbiology*, 66(8), 3415–3420.
- Khan, A. B., Khattak, M. I., Tarar, O. M., Habib, F., Jamil, K., Yasmin, A., & Parvez, S. (2013). Separation of polyhydroxyalkanoates-producing bacterial strains using PHA synthase gene and their evaluation for PHA deposition. *Brazilian Archives of Biology and*

- Khare, E., Chopra, J., & Arora, N. K. (2014). Screening for MCL-PHA-producing fluorescent *pseudomonas* and comparison of MCL-PHA production under iso-osmotic conditions induced by PEG and NaCl. *Current Microbiology*, 68(4), 457–462.
- Kim, T. K., Tri, M., Shin, H. D., & Lee, Y. H. (2005). Molecular structure of the PHA synthesis gene cluster from new mcl-PHA producer *Pseudomonas putida* KCTC1639. *Journal of Microbiology and Biotechnology*, 15(5), 1120–1124.
- Kshirsagar, S. D., Mattam, A. J., Jose, S., Ramachandrarao, B., & Velankar, H. R. (2020). Heavy hydrocarbons as selective substrates for isolation of asphaltene degraders: A substrate-based bacterial isolation strategy for petroleum hydrocarbon biodegradation. *Environmental Technology and Innovation*, 19, 100832.
- Lathwal, P., Nehra, K., Singh, M., & Rana, J. S. (2018). Characterization of Novel and Efficient Poly-3-hydroxybutyrate (PHB) Producing Bacteria Isolated from Rhizospheric Soils. *Journal of Polymers and the Environment*, 26(8), 3437–3450.
- Mohammed, S., Panda, A. N., & Ray, L. (2019). An investigation for recovery of polyhydroxyalkanoates (PHA) from *Bacillus* sp. BPPI-14 and *Bacillus* sp. BPPI-19 isolated from plastic waste landfill. *International Journal of Biological Macromolecules*, 134, 1085–1096.
- Mohandas, S. P., Balan, L., Jayanath, G., Anoop, B. S., Philip, R., Cubelio, S. S., & Bright Singh, I. S. (2018). Biosynthesis and characterization of polyhydroxyalkanoate from marine *Bacillus cereus* MCCB 281 utilizing glycerol as carbon source. *International Journal of Biological Macromolecules*, 119, 380–392.
- Możejko-Ciesielska, J., & Kiewisz, R. (2016). Bacterial polyhydroxyalkanoates: Still fabulous? *Microbiological Research*, 192, 271–282.
- Muhammadi, Shabina, Afzal, M., & Hameed, S. (2015). Bacterial polyhydroxyalkanoates-eco-friendly next generation plastic: Production, biocompatibility, biodegradation, physical properties and applications. *Green Chemistry Letters and Reviews*, 8(3–4), 56–77.
- Nandiyanto, A. B. D., Oktiani, R., & Ragadhita, R. (2019). How to read and interpret ftir spectroscopy of organic material. *Indonesian Journal of Science and Technology*, 4(1), 97–118.
- Narita, V., Arum, A. L., Isnaeni M, S., & Fawzya, N. Y. (2014). Analisis Bioinformatika Berbasis WEB untuk Eksplorasi Enzim Kitosanase Berdasarkan Kemiripan Sekuens. *Jurnal Al-Azhar Indonesia Seri Sains Dan Teknologi*, 1(4), 197.
- Philla, S. (2011). Handbook of Bioplastic and Biocomposites Engineering Applications. Wiley, USA.
- Sakai, M., Ezaki, S., Suzuki, N., & Kurane, R. (2005). Isolation and characterization of a novel polychlorinated biphenyl-degrading bacterium, *Paenibacillus* sp. KBC101. *Applied Microbiology and Biotechnology*, 68(1), 111–116.
- Shen, M., Song, B., Zeng, G., Zhang, Y., Huang, W., Wen, X., & Tang, W. (2020). Are biodegradable plastics a promising solution to solve the global plastic pollution? *Environmental Pollution*, 263, 114469.
- Shi, H. P., & Lee, C. M. (2007). Phosphate removal under denitrifying conditions by *Brachymonas* sp. strain P12 and *Paracoccus denitrificans* PP15. *Canadian Journal of Microbiology*, 53(6), 727–737.

- Shibulal, B., Al-Bahry, S. N., Al-Wahaibi, Y. M., Elshafie, A. E., Al-Bemani, A. S., & Joshi, S. J. (2017). The potential of indigenous *Paenibacillus ehimensis* BS1 for recovering heavy crude oil by biotransformation to light fractions. *PLoS ONE*, 12(2), 1–18.
- Subudhi, S., Batta, N., Pathak, M., Bisht, V., Devi, A., Lal, B., & Al khulifah, B. (2014). Biofloculant production and biosorption of zinc and lead by a novel bacterial species, *Achromobacter* sp. TERI-IASST N, isolated from oil refinery waste. *Chemosphere*, 113, 116–124.
- Tamura, K., Nei, M., & Kumar, S. (2004). Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proceedings of the National Academy of Sciences of the United States of America*, 101(30), 11030–11035.
- Tu, W., Zhang, D., Wang, H., & Lin, Z. (2019). Polyhydroxyalkanoates (PHA) production from fermented thermal-hydrolyzed sludge by PHA-storing denitrifiers integrating PHA accumulation with nitrate removal. *Bioresource Technology*, 292, 121895.
- Verlinden, R. A. J., Hill, D. J., Kenward, M. A., Williams, C. D., & Radecka, I. (2007). Bacterial synthesis of biodegradable polyhydroxyalkanoates. *Journal of Applied Microbiology*, 102(6), 1437–1449.
- Vijay, R., & Tarika, K. (2018). Banana peel as an inexpensive carbon source for microbial polyhydroxyalkanoate (PHA) production. *International Research Journal of Environmen*.7(1), 28–36.
- Wei, Y. H., Chen, W. C., Huang, C. K., Wu, H. S., Sun, Y. M., Lo, C. W., & Janarthanan, O. M. (2011). Screening and evaluation of polyhydroxybutyrate-producing strains from indigenous isolate *Cupriavidus taiwanensis*. *International Journal of Molecular Sciences*, 12(1), 252–265.
- Yasin, G., Bhanger, M. I., Ansari, T. M., Muhammad, S., Naqvi, S. R., Ashraf, M., Ahmad, K., & Talpur, F. N. (2013). Quality and chemistry of crude oils. *Journal of Petroleum Technology and Alternative Fuel* 4(3), 53–63.