



DAFTAR PUSTAKA

- Baneyx,F., dan M.Mujacic. 2004. Recombinant protein folding and misfolding in *Escherichia coli*. Nat. Biotechnol. 22(11): 1399-1408.
- Briand,L., G.Marcion, A.Kriznik, J.M.Heydel, Y.Artur, C.Garrido, R.Seigneuric, dan F.Neiers. 2022. A self-inducible heterologous protein expression system in *Escherichia coli*. Sci. Rep. 6(33037): 1-11.
- Costanzo,J.A., C.J.O'Brien, K.Tiller, E.Tamargo, A.S.Robinson, C.J.Roberts, dan E.J.Fernandez. 2014. Conformational stability as a design target to control protein aggregation. Protein Eng. Des. Sel. 27(5): 157-167.
- Cummins,P.M., K.D.Rochfort, dan B.F.O'Connor. 2017. Ion-exchange chromatography: Basic principles and application. Methods Mol. Biol.1485: 209-223.
- Darby, N., dan T.E.Creighton. 1995. Disulfide Bonds in Protein Folding and Stability. In: B.A.Shirley (Ed.). Protein Stability and Folding. Humana Press, New Jersey, pp: 219-252.
- Duong-Ly,K.C., dan S.B.Gabelli. 2014. Using Ion Exchange Chromatography to Purify a Recombinant Expressed Protein. In: J.Lorsch (Ed.) Methods in Enzymology. Academic Press, Netherlands, pp: 95-103.
- Feller,G., O.L.Bussy, dan C.Gerday. 1998. Expression of psychrophilic genes in mesophilic hosts: assessment of the folding state of a recombinant α -amylase. Appl. Environ. Microbiol. 64(3): 1163-1165.
- Getachew,A., F.Woldesenbet. 2016. Production of biodegradable plastic by polyhydroxybutyrate (PHB) accumulating bacteria using low cost agricultural waste material. BMC Res. Notes 9(509): 1-9.
- Gradissimo,D.G., L.P.Xavier, dan A.V.Santos. 2020. Cyanobacterial polyhydroxyalkanoates: A sustainable alternative in circular economy. Molecules 5(8): 1-23.
- Hausjell,J., J.Weissensteiner, C.Molitor, H.Halbwirth, dan O.Spadiut. 2018. *E.coli* HMS174(DE3) is a sustainable alternative to BL21(DE3). Microb. Cell Factories 17(169): 1-9.
- Hong,J., W.Park, H.Seo, I.Kim, dan K.Kim. 2020. Crystal structure of an acetyl-CoA acetyltransferase from phb-producing bacterium *Bacillus cereus* ATCC 14579. Biochem. and Biophys. Res. Commun. 533(3): 442-448.
- Jha,S.K., dan S.Marqusee. 2013. Kinetic evidence for a two-stage mechanism of protein denaturation by guanidium chloride. Proc. Natl. Acad. Sci. USA.



John,M.E., dan G.Keller. 1996. Metabolic pathway engineering in cotton: Biosynthesis of polyhydroxybutyrate in fiber cells. Proceedings of the “National Academy Sciences of the United States of America”, Washington,D.C., November 1996.

Kocharin,K., Y.Chen, V.Siewers, dan J.Nielsen. 2012. Engineering of acetyl-CoA metabolism for the improved production of polyhydroxybutyrate in *Saccharomyces cerevisiae*. AMB Express 2(52): 1-11.

Konigsberg,W. 1972. Reduction of disulfide bonds in proteins with dithiothreitol. Methods in Enzymol. 25: 185-188.

Kramer,R.M., V.R.Shende, N.Motl, C.N.Pace, dan J.M.Scholtz. 2012. Toward a molecular understanding of protein solubility: increased negative surface charge correlates with increased solubility. Biophys. J. 102(8): 1907-1915.

Kroemer,T. 2022. How does IPTG induction work?. <https://www.goldbio.com/>. Diakses tanggal 20 Mei 2022.

Kumar, D. 2018. Chromatographic Protein Refolding/Renaturation. Elsevier, Belanda.

Liu,P., L.V.Avramova, dan C.Park. 2009. Revisiting absorbance at 230 nm as protein unfolding probe. Anal. Biochem. 389: 165-170.

Luo,S., M.Mcneill, T.G.Myers, R.J.Hohman, dan R.L.Levine. 2008. Lon protease promotes survival of *Escherichia coli* during anaerobic glucose starvation. Arch. Microbiol. 189(2): 181-185.

Miles,A.J., S.G.Ramalli, dan B.A.Wallace. 2022. DichroWeb, a website for calculating protein secondary structure from circular dichroism spectroscopic data. Protein Sci. 31(1): 37-46.

Monera,O.D., C.M.Kay, dan R.S.Hodges. 1994. Protein denaturation with guanidine hydrochloride or urea provides a different estimate of stability depending on the contributions of electrostatic interactions. Protein Sci. 3: 1984-1991.

Mozejko-Ciesielska,J., K.Szacherska, dan P.Marchinak. 2019. *Pseudomonas* species as producers of eco-friendly polyhydroxyalkanoates. J. Polym. Environ. 27: 1151-1166.

Pace,C.N., S.Trevino, E.Prabhakaran, dan J.M.Scholtz. 2004. Protein structure, stability and solubility in water and other solvents. Philos. Trans. R. Soc. B. Biol. Sci. 359(1448):1225-1235.

Pignataro,M.F., M.G.Herrera, dan V.I.Dodero. 2020. Evaluation of peptide/protein self-assembly and aggregation by spectroscopic methods. Molecules 25(4854): 1-35.

Poirier,Y. 2003. Genetic modification of primary metabolism/Biopolymers. In: B.Thomas (Ed.) Encyclopedia of Applied Plant Sciences, Elsevier, pp: 477-484.



Rani,G.U., dan S.Sharma. 2022. Biopolymers, Bioplastics and Biodegradability: An Introduction. In: M.S.J.Hashmi (Ed.) Encyclopedia of Materials: Plastics and Polymers, Elsevier, pp: 474-486.

Raza,Z.A., S.Abid, dan I.M.Banat. 2018. Polyhydroxyalkanoates: Characteristics, production, recent development and applications. Int. Biodeterior. Biodegr. 126: 45-56.

Reusch, R. 2014. Biogenesis of ion channels. J. Biochem. Biophys. 1(1): 1-24.

Rizkia,P.R., S.Silaban, K.Hasan, D.S.Kamara, T.Subroto, S.Soemitro, dan I.P.Maksum. 2015. Effect of *isopropyl-β-D-thiogalactopyranoside* concentration on *prethrombin-2* recombinant gene expression in *Escherichia coli* ER2566. Procedia Chem. 17:118-124.

Rosano,G.L., dan E.A.Ceccarelli. 2014. Recombinant protein expression in *Escherichia coli*: Advances and challenges. Front. Microbiol. 5(172): 1-17.

Segura-Campos,M.R., L.Espinosa-Garcia, L.A.Chel-Guerrero, dan D.A.Betancur-Ancona. 2012. Effect of enzymatic hydrolysis on solubility, hydrophobicity, and in vivo digestibility in cowpea (*Vigna unguiculata*). Int. J. Food Prop. 15 (4): 770-780.

Singh,A.K., L.Sharma, N.Mallick, dan J.Mala. 2017. Progress and challenges in producing polyhydroxyalkanoate biopolymers from cyanobacteria. J. Appl. Phycol. 29(3): 1213-1232.

Song,J.M., An.Y.J, Kang.M.H, Lee.Y, dan Cha S. 2012. Cultivation at 6-10°C is an effective strategy to overcome the insolubility of recombinant proteins in *Escherichia coli*. Protein Expr. Purif. 82(2): 297-301.

Teteki,S.S. 2021. Kloning dan Ekspresi Gen *Acetyl-CoA Acetyltransferase (phaA)* pada *Escherichia coli* BL21 (DE3). Fakultas Pertanian. Universitas Gadjah Mada. Skripsi.

Trevino,S.R., J.M.Scholtz, dan C.N.Pace. 2008. Measuring and increasing protein solubility. J. Pharm. Sci. 97(10): 4155-4166.

Waegeman,H., dan W.Soetaert. 2011. Increasing recombinant protein production in *Escherichia coli* through metabolic and genetic engineering. J. Ind. Microbiol. Biotechnol. 38(12): 1891-1910.

X.Han., W.Ning, X.Ma, X.Wang, dan K.Zhou. 2020. Improving protein solubility and activity by introducing small peptide tags designed with machine learning models. Metab. Eng. Commun. 11: 1-9.

Yamaguchi,H., dan M.Miyazaki. 2014. Refolding techniques for recovering biologically active recombinant proteins from inclusion bodies. Biomolecules 4(1): 235-251.



Ylinen,A., H.Maaheimo, A.Angelescu-Hakala, M.Penttila, L.Salusjarvi, dan M.Toivari.
2021. Production of d-lactic acid containing polyhydroxyalkanoate (PHA) polymers
in yeast *Saccharomyces cerevisiae*. J. Ind. Microbiol. Biotechnol. 48(7): 1-12.

Zayas,J.F. 1997. Functionality of Proteins in Food. Springer, Germany.

Zhao,M., X.Tao, F.Wang, Y.Ren, dan D.Wei. 2018. Establishment of a low-dosage-IPTG
inducible expression system construction method in *Escherichia coli*. J. Basic
Microbiol. 58(9): 806-810.