

## LIST OF FIGURE

Fig. 1 The regional map at East Java and Bali. The research area include Ijen caldera and Mt. Ijen. ....	14
Fig. 2 The topological map of Ijen caldera, faults and post-caldera volcanoes. Both Kukusan and Genteng is located in inside of Ijen caldera and the distance between them is approximately 4 km. (Caudron et al., 2015)..	17
Fig. 3 The research schedule from March 2018 to December 2020. ....	23
Fig. 4 An example of coalescence bubble. ....	28
Fig. 5 The flow chart of this study from problem statement to conclusion....	29
Fig. 6 Outcrops of volcanic deposits. The length of yellow bar in the pictures is 1m. The description is in the text. (a) Kukusan deposit at location I05. The white ash underlies two scoria layers. (b) Kukusan deposit at location I17. (c) Kukusan deposit at location I99. (d) Genteng deposit at location I43.....	33
Fig. 7 Stratigraphy of Kukusan volcanic deposits. The thickness of scoria increases as the distance from Kukusan vent decreases. The alphabet and number next to layer is the sample name.....	34
Fig. 8 Dispersal map of Kukusan and Genteng scoria layer. Black square: location of outcrop, number above black square: location number, number below black square: thickness of scoria layer (cm), black triangle: location of volcano vent, solid line: isopach, dashed line: estimated isopach.....	35
Fig. 9 (a) Grain size distribution of pyroclastic deposits. Md and $\sigma$ is the median diameter and deviation, respectively (Walker, 1971). (b) The ratio of scoria type A, B and lithic materials of 8-16mm clasts. The number is the total pieces that is observed in one sample set. (c) Bulk density distribution of scoriae.....	38
Fig. 10 (a) The surface texture of scoria from Kukusan (location I89A). (b) The surface texture of scoria from Genteng (location I43B). ....	40
Fig. 11 Petrographic image of scoria. The black bar is 1mm. Phenocryst of plagioclase (Pl), olivine (Ol), orthopyroxene (Opx) and Fe-Ti oxide can be observed in these pictures. (a) Kukusan scoria under open nicole (location I89A). Bubbles are rich in abundance. (b) Kukusan scoria under crossed nicoles. Not only phenocryst but also micro-phenocryst of Pl can be seen. (c) Genteng scoria under open nicole (location I43A). (d)	

- Genteng scoria under crossed nicoles. The shape of Opx is rounded. ...43
- Fig. 12 (a) Thin section of Kukusan modeD scoria. The red square is the approximate part of observation of (c). (b) Thin section of Genteng mode scoria. The red square is the approximate part of observation of (d). (c) BSD image of bubbles in Kukusan scoria. (d) BSD image of bubbles in Genteng scoria. ....44
- Fig. 13 (a) Thick section of Kukusan modeD scoria. A square is the approximate location of observation of (c). (b) Thick section of Genteng mode scoria. The red square is the approximate part of observation of (d). (c) BSD image of microlites in Kukusan scoria. (d) BSD image of microlites in Genteng scoria. ....46
- Fig. 14 The classification of eruption types by dispersal and fragmentation (Walker, 1973). Kukusan is classified as sub-Plinian eruption, while Genteng is classified as Strombolian eruption. ....49
- Fig. 15 Comparison of bubble number density ( $N_{\text{bub}}$ ) for explosive eruption against magma  $\text{SiO}_2$  content. Solid line and dashed lines are contours of decompression rate with a water content of 4 and 8 wt.%, respectively (Toramaru, 2006). References: (1) This study, (2) Mangan and Cashman 1996, (3) Lautze and Houghton 2005, 2007, 2008 (4) Toramaru 2006 (5) Sable et al. 2006 (6) Criswell 1987, Klug and Cashman 1994 .....51
- Fig. 16 Comparison of microlite number density ( $N_{\text{mic}}$ ) for explosive eruption against microlite crystallinity (bubble-free). References: (1) This study, (2) Sable et al., 2006 (3) (Toramaru et al., 2008) .....52
- Fig. 17 The rate of water exsolution rate of Kukusan and Genteng is calculated by MND of pyroxene,  $\text{SiO}_2$  content and initial water content. ....55
- Fig. 18 (left) A model of vertical cross-section of conduit. High microlite crystallinity increases the viscosity of melt and suspend the bubble ascent to the upper side of conduit. (right) A sketch of conduit of Kukusan and Genteng. Higher ascent rate and higher microlite crystallinity in Kukusan make bubbles immobile relative to the melt. On the other hand, lower ascent rate and lower microlite crystallinity in Genteng make bubbles ascend easier. ....60

## LIST OF TABLE

Table 1 The comparison of focus, location, method and result of previous studies and this study. GSD, grain size distribution; XRF, X-ray fluorescence; SEM, scanning electron microscope.....	14
Table 2 The equipment list in this study. ....	21
Table 3 Bulk chemical compositions of major elements of scoria from Kukusan and Genteng volcano. ....	36
Table 4 Vesicularity and phenocryst assemblages of scoriae from Kukusan and Genteng. The classification of density category: vesicular (minimum density), modeV (the more vesicular mode), modeD (the denser mode), dense (maximum density). ....	42
Table 5 Number density and size of bubble in scoriae from Kukusan and Genteng. ....	45
Table 6 Number density, crystallinity and size of microlite in scoriae from Kukusan and Genteng. ....	47
Table 7 The viscosity of melt phase of Kukusan and Genteng estimated by the equation of Hui and Zhang (2007) using the data of chemical composition and typical temperature T and water content Cw. ....	57
Table 8 The relative viscosity of crystal-bearing magma of Kukusan and Genteng estimated by the method of Mader et al. (2013) using the data of microlite crystallinity and aspect ratio. ....	57