

DAFTAR PUSTAKA

- Abd-ellah, H., 2012. Ameliorative effect of captopril against 5-fluorouracil-induced cardiotoxicity in rats: a study with the light and electron microscopes. *J. Appl. Sci. Res.* 8: 863–872.
- Abd, B.H., & Azim, E., 2015. Biochemical studies of captopril against 5-fluorouracil induced heart toxicity in rats. *Int. J. Adv. Res.* 3: 247–261.
- Akaa, P.D., Ahachi, N.C., Vhritherhire, A.R., Agada, E., & Jenrola, A.A., 2017. Five year retrospective study on keloid management. *J. Adv. Med. Med. Res.* 23: 1–8.
- Ali, N.E.M., 2012. Protective effect of captopril against 5-fluorouracil-induced hepato and nephrotoxicity in male albino rats. *J. Am. Sci.* 8: 680–685.
- Andrews, J.P., Marttala, J., Macarak, E., Rosenbloom, J., & Uitto, J., 2016. Keloids: the paradigm of skin fibrosis - pathomechanisms and treatment. *Matrix Biol.* 51: 37–46.
- Arno, A.I., Gauglitz, G.G., Barret, J.P., & Jeschke, M.G., 2014. Up-to-date approach to manage keloids and hypertrophic scars : a useful guide. *Burns* 1–12.
- Bernstein, K.E., Ong, F.S., Blackwell, W.B., Shah, K.H., Giani, J.F., & Gonzalez-villalobos, R.A., 2013. A modern understanding of the traditional and nontraditional biological functions of angiotensin-converting enzyme. *Pharmacol Rev* 65: 1–46.
- Biernacka, A., Dobaczewski, M., & Frangogiannis, N.G., 2011. TGF- β signaling in fibrosis. *Growth factors* 29: 196–202.
- Bijlard, E., Steltenpool, S., & Niessen, F., 2015. Intralesional 5-fluorouracil in keloid treatment : a systematic review. *Acta Derm Venereol* 95: 778–782.
- Chen, C., Yang, S., Zhang, M., Zhang, Z., Zhang, B., Han, D., et al., 2013. In vitro sirius red collagen assay measures the pattern shift from soluble to deposited collagen, in: Welch, W., & Al, E. (Eds.), *Oxygen Transport to Tissue XXXIV*. Springer, New York, pp. 47–53.
- Chen, J., Zhao, S., Liu, Y., Cen, Y., & Nicolas, C., 2016. Effect of captopril on collagen metabolisms in keloid fibroblast cells. *ANZ J. Surg.* 86: 1046–1051.
- Chen, J.J., Jin, P.S., Zhao, S., Cen, Y., Liu, Y., Xu, X.W., et al., 2011. Effect of heat shock protein 47 on collagen synthesis of keloid in vivo. *ANZ J. Surg.* 81: 425–430.
- Choi, M., Ding, Y., & Kim, S., 2012. TGF- β signaling via TAK1 pathway: role in kidney fibrosis. *Semin Nephrol* 32: 244–252.
- Cristina, K., Al-ahj, L.P., Oliveira, R.C., & Magalhães, C., 2014. Cell density and solvent are critical parameters affecting formazan evaluation in MTT assay. *Braz. Arch. Biol. Technol.* 57: 381–385.
- Fang, Q., Wang, X., Zhao, W., Chen, C., Zhang, M., & Shi, B., 2018a. The source of ACE during scar formation is from both bone marrow and skin tissue. *Fed. Am. Soc. Exp. Biol.* 32: 1–10.
- Fang, Q., Wang, X., Zhao, W., Ding, S., & Shi, B., 2018b. Angiotensin-converting enzyme inhibitor reduces scar formation by inhibiting both canonical and noncanonical TGF- β 1 pathways. *Sci. Rep.* 8: 1–10.

- Felice, B. De, Garbi, C., Santoriello, M., Santillo, A., & Wilson, R.R., 2009. Differential apoptosis markers in human keloids and hypertrophic scars fibroblasts. *Mol. Cell. Biochem.* 327: 191–201. doi:10.1007/s11010-009-0057-x
- Forbat, E., Ali, F., & Al-Niami, F., 2017. Treatment of keloid scars using light-, laser- and energy-based devices : a contemporary review of the literature. *Lasers Med Sci* 32: 2145–2154.
- Gauglitz, G., Bureik, D., Dombrowski, Y., Pavicic, T., Ruzicka, T., & Schaubert, J., 2012. Botulinum toxin A for the treatment of keloids. *Ski. Pharmacol Physiol* 25: 313–318.
- Gauglitz, G., & Korting, H., 2011. Hypertrophic scarring and keloids : pathomechanism and emerging treatment strategies. *Mol. Med.* 17: 113–125.
- Geraghty, R.J., Davis, J.M., Downward, J., Freshney, R.I., & Knezevic, I., 2014. Guidelines for the use of cell lines in biomedical research. *Br. J. Cancer* 111: 1021–1046. doi:10.1038/bjc.2014.166
- Gupta, S., & Kalra, A., 2002. Efficacy and safety of intralesional 5-fluorouracil in the treatment of keloids. *Pharmacol. Treat.* 204: 130–132.
- Hietanen, K.E., Järvinen, T.A., Huhtala, H., Tolonen, T.T., Kuokkanen, H.O., & Kaartinen, I.S., 2019. Treatment of keloid scars with intralesional triamcinolone and 5-fluorouracil injections - a randomized controlled trial. *J. Plast. Reconstr. Aesthetic Surg.* 72: 4–11.
- Huang, C., Liu, L., You, Z., Du, Y., & Ogawa, R., 2019. Managing keloid scars : from radiation therapy to actual and potential drug deliveries. *Int. Wound J.* 1–8.
- Huang, L., Cai, Y.J., Lung, I., Leung, B.C.S., & Burd, A., 2013. A study of the combination of triamcinolone and 5-fluorouracil in modulating keloid fibroblasts in vitro. *Br. J. Plast. Surg.* 66: 251–259.
- Huang, L., Wong, Y.P., Cai, Y.J., Lung, I., Leung, C.S., & Burd, A., 2010. Low-dose 5-fluorouracil induces cell cycle G2 arrest and apoptosis in keloid fibroblasts. *Br. J. Dermatol.* 163: 1181–1185.
- Khalid, F.A., Mehrose, M.Y., Saleem, M., Yousaf, M.A., Mujahid, A.M., Rehman, S.U., et al., 2018. Comparison of efficacy and safety of intralesional triamcinolone and combination of triamcinolone with 5-fluorouracil in the treatment of keloids and hypertrophic scars: Randomised control trial. *Burns* 1–7.
- Khamanga, S.M., & Walker, R.B., 2012. In vitro dissolution kinetics of captopril from microspheres manufactured by solvent evaporation. *Dissolution Technol.* 42–51.
- Kontochristopoulos, G., Stefanaki, C., Panagiotopoulos, A., Stefanaki, K., Argyrakos, T., Petridis, A., et al., 2005. Intralesional 5-fluorouracil in the treatment of keloids: An open clinical and histopathologic study. *Dermatologic Surg.* 52: 474–479.
- Kumar, P., Nagarajan, A., & Uchil, P.D., 2018. Analysis of cell viability by the MTT assay. *Cold Spring Harb Protoc.* 469–472.
- Kurniaty, I.R., Siswati, A.S., & Wirohadidjojo, Y.W., 2020. Efektivitas Kombinasi 5-Fluorourasil dan Timolol pada Proliferasi dan Deposisi Kolagen Biakan

Fibroblas Keloid. Universitas Gadjah Mada.

- Ladin, D.A., Hou, Z., Patel, D., Mcphail, M., Olson, J.C., Saed, G.M., et al., 1998. p53 and apoptosis alterations in keloids and keloid fibroblasts. *Wound Repair Regen.* 6: 28–37.
- Lee, H.J., & Jang, Y.J., 2018. Recent understandings of biology, prophylaxis and treatment strategies for hypertrophic scars and keloids. *Int. J. Mol. Sci.* 19: 1–19.
- Mahmood, A., & Ali, S., 2017. Microbial and viral contamination of animal and stem cell cultures : common contaminants , detection and elimination. *J. Stem Cell Res. Ther.* 2: 149–155. doi:10.15406/jsrt.2017.02.00078
- Mari, W., Alsabri, S.G., Tabal, N., Younes, S., Sherif, A., & Simman, R., 2016. Novel insights on understanding of keloid scar: article review. *J. Am. Coll. Clin. Wound Spec.* 7: 1–7.
- Mauviel, A., Verrecchia, F., Wendling, J., & Aim, E., 2003. 5-Fluorouracil Blocks Transforming Growth Factor beta– Induced alpha2 Type I Collagen Gene (COL1A2) Expression in Human Fibroblasts via c-Jun NH 2 -Terminal Kinase / Activator Protein-1 Activation. *Mol. Pharmacol.* 64: 707–713.
- Meerlo, J. Van, & Cloos, J., 2011. Cell sensitivity assays: The MTT assay. *Methods Mol. Biol.* 731: 237–245.
- Mohammadi, A.A., Parand, A., Kardeh, S., & Janati, M., 2018. Efficacy of topical enalapril in treatment of hypertrophic scars. *Wound J. Plast. Surg.* 7: 326–331.
- Morihara, K., Takai, S., Takenaka, H., Sakaguchi, M., Okamoto, Y., Morihara, T., et al., 2006. Cutaneous tissue angiotensin converting enzyme may participate in pathologic scar formation in human skin. *J. Am. Acad. Dermatol.* 54: 251–257.
- Motoki, T.H.C., Isoldi, F.C., Filho, A.G., & Ferreira, L.M., 2018. Keloid negatively affects body image. *Burns* 1–5.
- Murphy, A.M., Wong, A.L., & Bezuhly, M., 2015. Modulation of angiotensin II signaling in the prevention of fibrosis. *Fibrogenes. Tissue Repair* 8: 1–7.
- Nanda, S., & Reddy, B., 2004. Intralesional 5-fluorouracil as a treatment modality of keloids. *Dermatol Surg* 30: 54–57.
- Niazi, F., Hooshyar, S., Hedayatyanfard, K., Ziai, S., Doroodgar, F., Niazi, S., et al., 2018. Detection of angiotensin II and AT1 receptor concentrations in keloid and hypertrophic scar. *J. Clin. Aesthet. Dermatol.* 11: 36–39.
- Pakyari, M., & Farrokhi, A., 2013. Critical role of transforming growth factor beta in different phases of wound healing. *Adv. Wound Care* 2: 215–224.
- Phan, T., Lim, I.J., Bay, B., Qi, R., Huynh, H.T., Lee, S., et al., 2002. Differences in collagen production between normal and keloid-derived fibroblasts in serum-media co-culture with keloid-derived keratinocytes. *J. Dermatol. Sci.* 29: 26–34.
- Pramar, Y., Gupta, V. Das, & Bethea, C., 1992. Stability of captopril in some aqueous systems. *J. Clin. Pharm. Ther.* 17: 185–189.
- Qin, H., Liu, R., Nie, W., Li, M., Yang, L., Zhou, C., et al., 2021. Comparison of Primary Keloid Fibroblast Cultivation Methods and the Characteristics of Fibroblasts Cultured from Keloids , Keloid-surrounding Tissues , and Normal Skin Tissues. *Int. J. Morphol* 39: 302–310.

- Rabey, N., Goldie, S., & Price, R., 2017. 5-Fluorouracil for keloid scars (Protocol). *Cochrane Database Syst. Rev.* 1–13.
- Rittié, L., 2017. Method for Picrosirius red-polarization detection of collagen fibers in tissue section. *Methods Mol. Biol.* 1627: 395–407.
- Ryan, J., 2008. Understanding and Managing Cell Culture Contamination What Are the Major. *Corning Inc.* 1–24.
- Sadeghinia, A., & Sadeghinia, S., 2012. Comparison of the efficacy of intralesional triamcinolone acetonide and 5-fluorouracil tattooing for the treatment of keloids. *Dermatologic Surg.* 38: 104–109.
- Salati, S.A., 2019. Keloids – an extensive review in the light of recent. *J. Pakistan Assoc. Dermatol* 29: 225–249.
- Shah, M., Shah, A., Ninama, K., Mahajan, R., & Bilimoria, F., 2014. Efficacy of intralesional 5-fluorouracil, triamcinolone acetonide, and CO2 laser for treatment of keloids and hypertrophic scars - A comparative study. *J. Integr. Heal. Sci.* 2: 3–6.
- Shah, V. V, Aldahan, A.S., Mlacker, S., & Alsaidan, M., 2016. 5-fluorouracil in the treatment of keloids and hypertrophic scars : a comprehensive review of the literature. *Dermatol. Ther. (Heidelb).* 6: 169–183.
- Shaheen, A., 2018. Comprehensive review of keloid formation. *Clin Res Dermatol* 4: 1–18.
- Shivaswamy, K., Shyamprasad, A., Sumathy, T., & Suparna, M., 2015. Clinical efficacy of low dose intralesional 5-fluorouracil in the treatment of keloids. *J. Evol. Med. Dent. Sci.* 4: 16229–16231.
- Teofoli, P., Barduagni, S., Ribuffo, M., Campanella, A., & Pita, O. De, 1999. Expression of Bcl-2, p53, c-jun and c-fos protooncogenes in keloids and hypertrophic scars. *J. Dermatol. Sci.* 22: 31–37.
- Tripathi, S., Soni, K., Agrawal, P., Gour, V., Mondal, R., & Soni, V., 2020. Hypertrophic scars and keloids : a review and current treatment modalities. *Biomed. Dermatology* 4: 1–11.
- Tucci-viegas, V.M., Hochman, B., & Ferreira, L.M., 2010. Keloid explant culture : a model for keloid fibroblasts isolation and cultivation based on the biological differences of its specific regions. *Int. Wound J.* 7: 339–348.
- Vijayakumar, A., Sanghvi, H., & Murdeshwar, P., 2015. Effect of Enalapril on Hypertrophic Scars and Keloids. *J. Adv. Plast. Surg. Res.* 1: 10–13.
- Xu, J., Yang, E., Yu, N.Z., & Long, X., 2017. Radiation therapy in keloids treatment: history, strategy, effectiveness, and complication. *Chin Med J* 130: 1715–1721.