

DAFTAR PUSTAKA

- Aep, W. 2006. Budidaya tanaman kedelai (*Glycine max* (L.) Merrill). Tesis. Universitas Padjadjaran. Bandung.
- Adie, M. dan Krisnawati, A. 2007. Biologi tanaman kedelai. *Balai Penelitian Kacang-kacangan dan Umbi-umbian* (BALITKABI). Malang.
- Adisarwanto, T. 2005. *Budidaya kedelai dengan pemupukan yang efektif dan pengoptimalan peran bintil akar*. Penebar Swadaya, Jakarta.
- Adrienne, B.N., Andrea, L and Boyce, C.K. 2011. The evolution and functional significance in the angiosperms. *Funct, Plant Biol* 38: 535-552.
- Agisimanto, D., Martasari, C dan Supriyanto, A. 2007. Perbedaan primer RAPD dan ISSR dalam identifikasi hubungan kekerabatan genetik jeruk siam (*Citrus suhuniensis* L. Tan). *Indonesia.J. Hort* 17 (2): 101-110.
- Agrawal, P.C. 2014. Genetic diversity analysis among soybean (*Glicine max* (L.) Merr.) genotype through ISSR molecular marker. *Ijapsa* 01: 1-7.
- Alamri, S. 2013. Comparative analysis of soybean accessions using ISSR and RAPD markers. Thesis. Laurentian University.
- Anggraeini, E. 2008. Random Amplified Polymorphic DNA (RAPD) suatu metode analisis DNA dalam menjelaskan berbagai fenomena biologi. *Biospecies* 1 (2):73-76.
- Anonimus. 2008. *Mutu Kedelai Nasional Lebih Baik dari Kedelai Impor* [Siaran Pers]. Jakarta: Badan Litbang Pertanian. <http://pustaka.litbang.deptan.go.id/ppi/lengkap/sp1202081.pdf>
- Ansari, A., Rahman, M dan Husain, M.A. 2011. Comparison among clustering in multivariate analysis of rice using morphological traits, physiological traits and simpe sequence repeat markers. *America-Eurasian. Agr and Env Sci.* 11 (6): 876-882.
- Araujo, F., Pacheco, M.V., Vieira, C., Felix, F.C and Chagas, K.P.T. 2016. ISSR molecular markers for the study of the genetic diversity of *Mimosa caesapiniaefolia* Benth. *IDESIA (Chile) Junio* 34: 47-52.
- Azrai, D.W. 2009. Teknik isolasi DNA genom tanaman pepaya dan jeruk dengan menggunakan modifikasi buffer CTAB. *Buletin Teknik Pertanian* 14 (1): 12-16.
- Baihaki, A. 2000. *Teknik rancang dan analisis penelitian pemuliaan*. Universitas Padjajaran. Diktat mata kuliah: Bandung. 91 hlm.
- Brown, G., Thompson, J., Nelson, R., and Warburton, M. 2000. Evaluation of genetic diversity of soybean introductions and North American ancestor using RAPD and SSR markers. *Crop Science* 40: pp: 815-823

- Bustaman, M dan S. Moeljopawiro. 1998. Pemanfaatan teknologi sidik jari DNA di bidang pertanian. *Zuriat* 9 (2): 7-90.
- Carter, T.E., Nelson, R., Sneller, C.H., and Cui, Z. 2004. *Soybeans: Improvement, Production and Uses*, (3rd-Ed). Madison, Wisconsin, USA. pp 25-30
- Chaerani, H.N dan Utami, D.W. 2011. Keragaman genetik 50 aksesori plasma nutfah kedelai berdasarkan penanda mikrosatelit. *Jurnal AgroBiogen*. 7 (2): 96-105.
- Charters, Y.M., Robertson, A., Wilkinson, M.J and Ramsay, G. 1996. PCR analysis of oilseed rape cultivars (*Brassica napus* L. ssp. *oleifera*) using 5'-anchored simple sequence repeat (SSR) primers *Theor. Journal of Applied Genetics* 92: 442-447.
- Chavez, H., Borromeo, T., Borines, N.O and Gentallan, R.Jr. 2016. Phenotypic diversity of soybean (*Glycine max* (L.) Merr.) accessions in the Philippines for utilization. *Legume research* 40 : 9-15.
- Chowduri, A., Sriniver, P., Tongpamnak, P., Saksoong, P and Chattwchirawong, P. 2002. Genetic relationship among exotic soybean introductions in Thailand: Consequence for varietal registration. *Science Asia* 28 : pp. 227-239.
- Diwan, N and Cregan, P.B. 1997. Automates sizing of fluorescent-labeled simple sequence repeat (SSR) markers to assay genetic variation in soybean. *Theor. Appl. Genet* 95: 723-733.
- Doyle, J.J., and Doyle, J.L. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* 19:11-15.
- Ellison, A.M., Bukley, H.L., Miller, T.E and Gotelli, N.J. 2004. Morphological variation in *Sarracenia purpurea* (Sarraceniaceae) geographic, environmental, and taxonomic correlates. *Am. J. Bot.* 91 (11): 1930-1935.
- Fenta, B.A., Schluter, U., Garcia, B.M., DuPlessis, M., Foyer, C.H and Kumert, K.J. 2011. Identification and application of phenotypic and molecular markers for abiotic stress tolerance in Soybean. In *Soybean-Genetics and novel techniques for yield enhancement*, Prof. Dora Krezhova (Ed.) pp 181-195.
- Fukushima, D. 2013. *Recent progress in research and technology on soybeans*. Kikkoman Corporation Noda, Chiba. 237-278.
- Gepts, P. 2005. Legumes as a model plant family. Genomics for food and feed report of the cross-legume advances through genomics conference. *Plant Physiol* 13: 1228-1235.
- Giancola, S.S., Poltri, M., Lacaze, P., and Hopp, H.E. 2002. Feasibility of integration of molecular markers and morphological descriptors in a real study of a plant variety protection system for soybean. *Euphytica* 127: 95-113.
- Gijzen, M., Miller, S.S., Kuflu, K., Buzzell, R.I., and Miki, B.L.A. 2003. Soybean seed lustre phenotype and surface protein cosegregate and map to linkage group. *E Genome* 46: 659-664.

- Godwin, I.D., Aiken, E.A.B and Smith, L.W. 1997. Application of inter-simple sequence repeat (ISSR) markers to plant genetics. *Electrophoresis* 18: 1524–1528.
- Gower, J.C. 1971. A General coefficient of similarity and some of its properties. *Biometrics* 27: 857-874.
- Gyoung, A., Crawford, G.W., Liu, L., Sasaki, Y and Chen, X. 2011. Archaeological soybean (*Glycine max*) in East Asia: does size matter?. *PLOS ONE* 6: 11-17.
- Hammer, K and Morimoto, Y. 2011. *Classification of infraspecific variation in crop plants*. In: Guarimo, L. (Eds). *Collecting plant genetic diversity: Technical guidelines*. Biodiversity International.
- Hamilton, M.B. 2009. *Population genetics*. Wiley-Blackwell. UK. P: 357.
- Handayani, F. 2015. *Analisis keragaman genetik lai (*Durio kutejensis* (Hassk.) Becc.) berdasarkan penanda morfologis dan Inter Simple Sequence Repeat*. . Tesis (tidak dipublikasikan). Fakultas Pertanian Universitas Gadjah Mada: Yogyakarta.
- Handayani. 2014. *Persilangan untuk merakit varietas unggul baru kentang*. Bandung (ID): Balai Penelitian Tanaman Sayur.
- Hanelt, P. 1986. *Formal and non-formal classification of the infraspecific variability of cultivated plants- advantages and limitations*. In: Styles, B.T. (Ed.). *Infraspecific Classification of Wild and Cultivated Plants*. The Systematic association Special Volume. No. 29. Oxford University Press, New York. Pp. 139-156.
- Henderson, A. 2006. Traditional morphometrics in plant systematics and It's role in palm systematic. *Bot. J. Linn. Soc.* 151: 103-111.
- Hidayat, O.D.1985. *Morfologi tanaman kedelai*. (Eds.). Puslitbangtan: Bogor. Hal 73-86.
- Hiromoto, D.M and Velo, V.A.1986. The genetic base of brazilian soybean (*Glycine max* (L.) Merrill) Cultivars. *Brazilian Journal of Genetics* 9 (2): 295-306.
- Hutchinson, J.B. 1990. *Essays in crop plant evolution*. Cambridge University Press. New York. pp 341-367.
- Hymowitz, T. 1995. *Evaluation of wild perennial glycine species and crosses for resistance to hakopsora*". In Sinclair, J.B.; Hartman, G.L. *Proceedings of the Soybean Rust Workshop*. Urbana, IL: National Soybean Research Laboratory. pp. 33–37.
- IBPGR. 1984. *Descriptor for Soybean*. International Board for Plant Resources. Italy. pp. 19-38.
- Izhaki, I., Tsahar E., Paluy, I., and Friedman, J. 2002. Within population variation and interrelationships between morphology, nutritional content, and secondary compound of *Rhamnus alaternus* fruits. *New Phytologist* 156: 217-223.



- Jamilah, C., Waluyo, B., and Kurniawan, A. 2011. *Parameter genetik aksesori tanaman kerabat liar ubi jalar koleksi Unpad untuk peningkatan genetik dan sumber perbaikan karakter ubi jalar*. Makalah disampaikan pada Seminar Nasional Pemuliaan Berbasis Potensi dan Kearifan Lokal Menghadapi Tantangan Globalisasi, diselenggarakan atas Kerjasama Perpipi Komda Banyumas dan Lembaga Penelitian dan Pengabdian kepada Masyarakat Universitas Jenderal Soedirman Purwokerto, Jawa Tengah.
- Jeong, N., Suh, S.J., Kim, M.H., Lee, S., Moon, J.K., Kim H.S., and Jeong, S.C. 2012. Ln is a key regulator of leaflet shape and number of seeds per pod in soybean. *The Plant Cell* 24 (12): 4807-4818.
- Judd, W.S., Campbell, C.S., Kellogg, E.A., Stevens, P.F., and Donoghue, M.J. 2007. *Taxonomy in Plant Systematics—A Phylogenetic Approach*. Sinauer Associates: Sunderland. pp 221-230.
- Kaplan, D.R. 2001. The science of plant morphology; definition, history dan role in modern biology. *American Journal of Botany* 88: 1711-1741.
- Lanham, P.G., and Brennan, R.M. 1999. Genetic characterization of gooseberry (*Ribes grossularia* subgenus *grossularia*) germplasm using rapid, issr and aflp markers. *The Journal of Horticultural Science and Biotechnology* 74: 361-366.
- Lorenz, T. 2012. Polymerase Chain Reaction: basic protocol troubleshooting and optimization strategies. *J Vis Exp* (63): 39-98.
- Mandl, F.A. and Buss, G.R. 1981. Comparison of narrow and broad leaflet isolines of soybean. *Crop Sci.* 2:25-27.
- Mariana, B.D., Sugiyatno, A., and Supriyanto, A. 2011. Genetic diversity of local accessions of *Dimocarpus longan* revealed by ISSR markers. *Bul Plasma Nutrafah* 16 (1): 1-7.
- Mokkamul, P., Chaveerach, A., Sudmoon, R., and Tanee, T. 2007. Species identification and sex determination of the genus *Nepenthes* (Nepenthaceae). *Pak. J. Biol. Sci* 10: 561-567.
- Mondini, L., Noorani, A., and Pagnotta, M.A. 2009. Assessing plant genetic diversity by molecular tools. *Diversity*, 1: 19-35; doi: 10.3390/d101001.
- Monpara, J., Chudasama, K., and Thaker, V. 2017. ISSR studies on small and large seed varieties of *Glycine max*. *Journal of pharmacognosy and phytochemistry* 6 (6): 1652-1656.
- Nasution, M.A., Nur, B.G., and Razak, Z. 2011. Keragaman genetik beberapa aksesori markisa berdasarkan penanda ISSR. *J. Agrivigor* 10 (2) :155-167.
- Nelson, R.L., and Wang, P. 1989. Variation and evaluation of seed shape in soybean. *Crop Sci.* 29:147-150.
- Ng, W.L and Tan, S.G. 2015. Inter-Simple Sequence Repeat (ISSR) Markers: Are We Doing It Right?. *ASM Sci. J* 9(1): 30–39.

- Peric, V., Nicolic, A., Babic, V., Sudaric, A., Srebric, M., Dordevic, V and Mladenovic Drinic, S. 2014. Genetic relatedness of soybean genotypes based on agromorphological traits and rapd-markers. *Genetika* 46: 839-854.
- Pfeifer, T.W., Peyyala, R., Ren, Q and Ghabrial, S.A. 2003. Increased soybean pubescence density. *Crop breeding, genetics & cytology* 43 (6): 2071-2076.
- Phatmawati, M. 2009. Optimalisasi ekstraksi DNA dan PCR-RAPD pada *Grevillea spp.* (Proteaceae). *Jurnal Biologi* 13 (1): 12-16.
- Pitopang, R., and Gradstein, S.R. 2004. Herbarium celebense (ceb) and its role in supporting research on plant diversity of Sulawesi. *Journal Biodiversitas* 5: 36-41.
- Poehlman, J.M., and Sleper, D.A. 1995. *Breeding field crops*. Iowa State University Press, Ames, IA.
- Pujiasmanto, B., Moenandir, J., Syamsulbahri., and Kuswanto. 2007. Kajian argoekologi dan morfologi sambiloto (*Andrographis paniculata* ness.) pada berbagai habitat. *Journal Biodiversitas* 8 (4): 326-329.
- Porter, C. 2000. *Inheritance of the Gene(s) Controlling leaflet shape in soybeana crop and soil environmental science*. Blacksburg. Thesis. Virginia Polytecnic institute.
- Rahayu, S.E., dan Handayani. 2010. Keragaman Genetik Pandan Asal Jawa Barat Berdasarkan Penanda *Inter Simple Sequence Repeat*. *Makara, Sains* 14:158-162.
- Rahmah, A. 2013. *Hubungan kekerabatan aksesori Purwoceng (Pimpinella pruatjan Molkenb.) di Pulau Jawa berdasarkan karakter morfologis dan molekular*. (Tesis). Program Pascasarjana Fakultas Biologi Universitas Gadjah Mada: Yogyakarta.
- Rasyad, A dan Idwar. 2010. Interaksi genetic x lingkungan dan stabilitas komponen hasil berbagai genotip kedelai di Provinsi Riau. *J. Agron. Indonesia* 38: 25-29.
- Reddy, M.P., Sarla, N., and Siddiq, E.A. 2002. Intersimple sequence repeat (ISSR) polymorphism and its application in plant breeding. *Euphytica* 128: 9-17.
- Reddy, B.P.K., Priya S.M., and Reddy M.D. 2013. Studies on the stability of hilum colour in soybean (*Glycine max* (L.) Merrill) cultivars. *Current Biotica* 6(4): 509-511.
- Roldan-ruiz, J., Euwijk, F.A., Gilliland, T.J., Dubreuil, P., Dillmann, C., Lallemand., De Loose., and Baril, C.P. 2001. A comparative study of molecular and morphological methods of describing relationships between perennial ryegrass (*Lolium perene* L.) varieties. *Theor. Appl. Genet* 103:1138-1150.
- Rukmana, R., dan Yuniarsih, Y. 1996. *Kedelai budidaya dan pascapanen*. Kanisius. Yogyakarta. pp 5-40.



UNIVERSITAS
GADJAH MADA

Variabilitas Genetik dan Klasifikasi Intraspesies Non Formal Kedelai (*Glycine max* (L.) Merrill) di Sulawesi Berdasarkan Karakter Morfologis dan Molekular

FIONE Y YALINDUA, Dr. Purnomo, M.S; Dr. Budi Setiadi Daryono, M.Agr.Sc

Universitas Gadjah Mada, 2018 | Diunduh dari <http://etd.repository.ugm.ac.id/>

- Sibositi, D. 2010. *Variasi intraspesifik Stevia (Stevia rebaudiana Bertoni) koleksi balai besar penelitian dan pengembangan tanaman obat dan obat tradisional (B2P2TO-OT) berdasarkan karakter morfologis dan molekular*. (Tesis). Program Pascasarjana Fakultas Biologi Universitas Gadjah Mada. Yogyakarta.
- Sambrook, J., and Russell, R.W. 2001. *Molecular cloning: A laboratory manual, 3rd ed*. Cold spring harbor laboratory press. cold spring harbor. N.Y. pp 123.
- Simpson, M.G. 2010. "Chapter 1 Plant Systematics: an Overview". *Plant Systematics (2nd ed.)*. Academic Press. ISBN 978-0-12-374380-0.
- Siregar, U.J dan Olivia, R.D. 2013. Keragaman genetik populasi sengon (*Paraserianthes falcatari* (L) Nielsen) pada hutan rakyat di Jawa berdasarkan penanda RAPD. Institut Pertanian Bogor.
- Shon, M.K., and Blevins, D.G. 1990. Foliar boron applications increase the final number of branches and pods on branches of field-grown soybeans. *Plant Physiol* 92: 602-607.
- Shurtleff, W., and Aoyagi, A. 2012. *History of soybeans and soyfoods*. Lafayette, California. pp. 335-337.
- Sneath, P.H., and Sokal, R.R. 1973. *Numerical Taxonomy*. W.H. Freeman and company. San Fransisco. pp 114-132.
- Sofiari, E., and Kirana, R. 2009. Analysis of segregation pattern and distribution of several pepper characteristic. *J. Hort* 19: 255-253.
- Sokal, R.R., and Sneath, P.H.A. 1963. *Principle of Numerical Taxonomy*. W.H. Freeman Co, San Fransisco.
- Specht, J.E., Williams, J.H and Pearson, D.R. 1985. Near-isogenic analyses of soybean pubescence genes. *Crop Sci* 25: 92-96.
- Suhartina. 2003. *Perkembangan dan deskripsi varietas unggul kedelai 1918-2002*. Balai Penelitian Tanaman Kacang-kacangan dan Umbi-umbian, Malang.
- Suhartina., Hapsari, R.T., dan Purwantoro, A. 2016. Keragaman plasma nutfah kedelai berdasarkan keragaan karakter morfo-agronomis. *Bul. Plasma Nutfah* (2): 109-118.
- Sujata, B., Basavaraha, G.T., and Salimath, P.M. 2011. Gene action and heritability of leaf and reproductive characteristics in soybean. *Breed Sci*. 50: 45-51
- Suprpto. 1997. *Bertanam kedelai*. Penebar Swadaya. Jakarta.
- Suryatini, K.Y. 2011. *Analisis keragaman genetik jarak pagar (Jathropa curcas L) dengan metode ISSR*. Tesis. Universitas Udayana. Denpasar.



UNIVERSITAS
GADJAH MADA

Variabilitas Genetik dan Klasifikasi Intraspesies Non Formal Kedelai (*Glycine max* (L.) Merrill) di Sulawesi Berdasarkan Karakter Morfologis dan Molekular

FIONE Y YALINDUA, Dr. Purnomo, M.S; Dr. Budi Setiadi Daryono, M.Agr.Sc

Universitas Gadjah Mada, 2018 | Diunduh dari <http://etd.repository.ugm.ac.id/>

- Susandarini, R. 2014. *Biosistematika pomelo (Citrus maxima (Burm). Merr.) di Indonesia berdasarkan kajian morfologis, fitokimia dan molekular*. Disertasi (tidak dipublikasikan). Fakultas Biologi Universitas Gadjah Mada: Yogyakarta.
- Susantidiana., Wijaya, A., Lakitan, B and Surahman, M. 2009 . Identifikasi beberapa aksesori jarak pagar (*Jathropa curcas* L.) melalui analisa RAPD dan morfologi. *Jurnal agronomi Indonesia* 37 (2): 167-173.
- Tyagi, S.D and Sethi, J. 2011. Genetic diversity pattern in soybean (*Glycine max* L. Merrill). *Res. J. Agric. Sci.* 2: 288-290.
- Udensi, O., Ikpeme, E.V., Markson, A.A., Edu, E.A.B., Umana and Urua, E.J. 2010. Selection of soybean genotypes using morphological markers. *IJCR* 7: 05-08
- Valliyodan, B. and Nguyen, H.T. 2012. *Biological mechanisms that influence soy protein concentration and composition*. In: Wilson, R. F.ed., *Designing Soybeans for the 21st Century Markets*. AOCS, IL, USA.
- Vanijajiva, O. 2012. The application of ISSR markers in genetic variance detection among durian (*Durio zibethinus* Murr.) cultivar in the Nonthaburi Province, Thailand. *Procedia Engineering* 32 :155-159.
- Verdcourt, B. 1966. A proposal concerning *Glycine* L. *Taxon* 15 : 34-36.
- Vernetti, D.F and de Jesus Vernetti, J.F. 2017. *Qualitative traits in breeding*. In : Lopes da Silva., Borem, A., Sediya, T., Ludke, W. (eds) *Soybean Breeding*. Springer, Cham.
- Wang, J., Ha, W., Ngan, F., But, H., and Shaw, P. 2001. Application of sequence characterized amplified region (SCAR) analysis to authenticate *Panax* species and their adulterants. *Planta Medica* 67: 781-783.
- Wang, K.J. and Takahata, Y., 2007. A preliminary comparative evaluation of genetic diversity between Chinese and Japanese wild soybean (*Glycine soja*) germplasm pools using SSR markers. *Genetic Resources and Crop Evolution* 54: 157–165.
- Wang, K.J., Li, X.H., and Yan, M.F. 2014. Microsatellite markers reveal genetic diversity of wild soybean in different habitats and implications for conservation strategies (*Glycine soja*) in China. *Conservation Genetics* 15: 605-618.
- Wilcox, J. R 2004. *World distribution and trade of soybean*. In: Boerma, H.R. & J.E. Specht., ed., *Soybeans: Improvement, Production and Uses*, *American Society of Agronomy*, Madison, WI, USA. pp 1-14.
- Wilson, R.F. 2008. Soybean: market driven research needs in genetics and genomics of soybean. *Springer* 2: 17-23.



UNIVERSITAS
GADJAH MADA

Variabilitas Genetik dan Klasifikasi Intraspesies Non Formal Kedelai (*Glycine max* (L.) Merrill) di Sulawesi Berdasarkan Karakter Morfologis dan Molekular

FIONE Y YALINDUA, Dr. Purnomo, M.S; Dr. Budi Setiadi Daryono, M.Agr.Sc

Universitas Gadjah Mada, 2018 | Diunduh dari <http://etd.repository.ugm.ac.id/>

- Widiastuti, A.D. 2010. Analisis keragaman genetik manggis (*Garcinia mangostana* L.) hasil iradiasi sinar gamma berdasarkan morfologi, anatomi dan penanda ISSR. Tesis. Institut Pertanian Bogor. Bogor.
- Yan , X., Zhao, H.,Liu, X., Li, Q., Qang, Y., Yuan, C., and Dong, Y., 2013. Phenotypic traits and diversity of different leaf accessions of the wild soybean (*Glycine soja* Sieb. Et Zucc.) in China.*Can J. Plant Sci* 3:168-178.
- Yantama, E. 2012. Keragaman dan heritabilitas karakter agronomi kedelai (*Glycine max* [L.] Merrill) generasi F2 hasil persilangan Wilis dan Malang. Universitas Lampung. Bandar lampung.
- Yeh, F.C., Yang, R.C., Boyle, T.B.J., Ye, Z.H., and Mao, C.J. 1999. *The User-friendly shareware for population genetic analysis*. Molecular Biology and Biotechnology Center. University of Alberta, Edmonton.
- Ye, Y., Wang, Z., and Zhuo, S. 2012. Soy germ isoflavones improve menopausal symptoms but have no effect on blood lipids in early postmenopausal Chinese women: a randomized placebo-controlled trial. *Menopause* 19 (8): 791-800.
- You, M.G., Liu, Y.B., Zhao, T.J., and Gai, J.Y. 1995. Effect of leaf shape on seed yields and its component in soybeans. *Soybean Genet Newlet* 22: 66-70.
- Zhang, J., Specht, J.E., Graef, G.L., and Johnson , B.E. 1992. Pubescence density effects on soybean seed yield and other agronomic traits. *Crop Sci.* 32: 641-648.
- Ziekiewicz, E., Rafalski, A., and Labuda, D. 1994. Genomic fingerprinting by simple sequence repeat (ssr)-anchored polymerase chain reaction amplification. *Genomics* 20:176-183.
- Zulchy, T dan Sutoro. 2016. Keragaman genetik plasma nutfah kedelai (*Glycine max* L.) berdasar karakter morfologis dan hasil. Balai Besar Bitoteknologi dan Sumber Daya Genetik Pertanian.