

BIBLIOGRAPHY

- Aletaha, D., Funovits, J., & Smolen, J. S. (2011). Physical disability in rheumatoid arthritis is associated with cartilage damage rather than bone destruction. *Ann of the Rheumatic Diseases*, 70(5), 733–739.
- Amador-Patarroyo, M. J., Rodriguez-Rodriguez, A., & Montoya-Ortiz, G. (2012). How does age at onset influence the outcome of autoimmune diseases? *Autoimmune Diseases*, 1(1).
- Ancelin, M. L., Scali, J., Norton, J., Ritchie, K., Dupuy, A. M., Chaudieu, I., & Ryan, J. (2017). Heterogeneity in HPA axis dysregulation and serotonergic vulnerability to depression. *Psychoneuroendocrinology*, 77, 90–94.
- Audhya, T., Adams, J. B., & Johansen, L. (2012). Correlation of serotonin levels in CSF, platelets, plasma, and urine. *Biochimica et Biophysica Acta - General Subjects*, 1820(10), 1496–1501.
- Basker, M., Moses, P. D., Russell, S., & Russell, P. S. S. (2007). The psychometric properties of beck depression Inventory for adolescent depression in a primary-care paediatric setting in India. *Child and Adolescent Psychiatry and Mental Health*, 1, 1–7.
- Bortolato, B., Carvalho, A. F., Soczynska, J. K., Perini, G. I., & McIntyre, R. S. (2015). The Involvement of TNF- α in Cognitive Dysfunction Associated with Major Depressive Disorder: An Opportunity for Domain Specific Treatments. *Current neuropharmacology*, 13(5), 558-76.
- Cowen, P. J., & Browning, M. (2015). What has serotonin to do with depression? *World Psychiatry*, 14(2), 158–160.
- Cross, M., Smith, E., Hoy, D., Carmona, L., Wolfe, F., Vos, T., ... March, L. (2014). The global burden of rheumatoid arthritis: Estimates from the Global Burden of Disease 2010 study. *Annals of the Rheumatic Diseases*, 73(7), 1316–1322.
- Gibofsky, A. (2014). Epidemiology, pathophysiology, and diagnosis of rheumatoid arthritis: A Synopsis. *The American Journal of Managed Care*, 20(7 Supplement), S128–S135.
- Hasler, G. (2010). Pathophysiology of depression: Do we have any solid evidence of interest to clinicians? *World Psychiatry*, 9(3), 155–161.
- Jackson-Koku, G. (2016). Beck Depression Inventory. *Occupational Medicine*, 66(2), 174–175.

- Jeong, H., Baek, S. Y., Kim, S. W., Eun, Y. H., Kim, I. Y., Kim, H., ... Cha, H. S. (2017). Comorbidities of rheumatoid arthritis: Results from the Korean National Health and Nutrition Examination Survey. *PLoS ONE*, 12(4), 1–15.
- Kopp, S., & Alstergren, P. (2002). Blood serotonin and joint pain in seropositive versus seronegative rheumatoid arthritis. *Mediators of Inflammation*, 11(4), 211–217.
- Lubberts, E., & van den Berg, W.B. (2013). Cytokines in the Pathogenesis of Rheumatoid Arthritis and Collagen-Induced Arthritis. *Madame Curie Bioscience Database of Landes Bioscience*. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK6288/>
- Matcham, F., Rayner, L., Steer, S., & Hotopf, M. (2013). The prevalence of depression in rheumatoid arthritis: a systematic review and meta-analysis. *Rheumatology*, 52(12), 2136–2148.
- Mateen, S., Moin, S., Shahzad, S., & Khan, A. Q. (2017). Level of inflammatory cytokines in rheumatoid arthritis patients: Correlation with 25-hydroxy vitamin D and reactive oxygen species. *PloS one*, 12(6), e0178879. doi:10.1371/journal.pone.0178879
- McWilliams, D. F., & Walsh, D. A. (2017). Pain mechanisms in rheumatoid arthritis. *Clinical and Experimental Rheumatology*, 35(5), S94–S101.
- Mitani, H., Shirayama, Y., Yamada, T., & Kawahara, R. (2006). Plasma levels of homovanillic acid, 5-hydroxyindoleacetic acid and cortisol, and serotonin turnover in depressed patients. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 30(3), 531–534.
- Oliver, J. E., & Silman, A. J. (2009). Why are women predisposed to autoimmune rheumatic diseases? *Arthritis Research and Therapy*, 11(5), 1–9.
- Sadock, B., Sadock, V., & Ruiz, P. (2015). *Kaplan and Sadock's Synopsis of Psychiatry: Behavioral Sciences/Clinical Psychiatry*. (C. Pataki & N. Sussman, Eds.) (Eleventh). Wolters Kluwer.
- Saldanha, B. D., Kumar, M. N., Capt, S., Ryali, V., Srivastava, K., Capt, S., & Pawar, A. A. (2009). Serum Serotonin Abnormality in Depression.pdf. *MJAFI*, 65, 108–112.
- Sekiduka-Kumano, T., Kawayama, T., Ito, K., Shoji, Y., Matsunaga, K., Okamoto, M., ... Hoshino, T. (2013). Positive association between the plasma levels of 5-hydroxyindoleacetic acid and the severity of depression

in patients with chronic obstructive pulmonary disease. *BMC Psychiatry*, 13.

Slavich, G. M., & Irwin, M. R. (2014). Social signal transduction theory of depression. *Psychiatric Bulletin*, 140(3), 774–815.

Sullivan, G. M., Oquendo, M. A., Huang, Y., & Mann, J. J. (2006). Elevated cerebrospinal fluid 5-hydroxyindoleacetic acid levels in women with comorbid depression and panic disorder. *The International Journal of Neuropsychopharmacology*, 9(05), 547.

Supriyanto, I. (2017). *Correlation between depression score and immune, inflammatory, and neurotrophic states of patients with rheumatoid arthritis*. Universitas Gadjah Mada.

Szeitz, A., & Bandiera, S. M. (2018). Analysis and measurement of serotonin. *Biomedical Chromatography*, 32(1).

Wirasto, R. T. (2017). *Cognitive Visual Perception Therapy for Reducing Depression in Patients with Rheumatoid Arthritis*.

Wulsin, L. R., Musselman, D., Otte, C., Bruce, E., Ali, S., & Whooley, M. a. (2009). Depression and Whole Blood Serotonin in Patients With Coronary Heart Disease From the Heart and Soul Study, 71(3), 260–265.

Zhang, L., Wang, J., Zhang, Q., Fu, T., Yin, R., Wang, Z., ... Gu, Z. (2017). Factors associated with hand joint destruction in Chinese patients with rheumatoid arthritis. *BMC Musculoskeletal Disorders*, 18(1), 1–9.