

DAFTAR PUSTAKA

- Abergel R.P., Pizzurro D., Meeker C.A., *et al.* 1985. Biochemical composition of the connective tissue in keloids and analysis of collagen metabolism in keloid fibroblast cultures. *J Invest Dermatol.* 84:384-8
- Adams BB., Gloster HM. Surgical pearl : excision with suprakeloidal flap and radiation therapy for keloid. *J Am Acad Dermatol* 2002; 47:307-9
- Ali S., Saeed S. 2011. Comparison of The Efficacy of Intralesional Triamcinolone Acetonide and 5-Fluorouracil Tattooing for The Treatment of Keloids. *Dermatol Surg.* 38 : 104-109.
- Al-Attar A., Mess S., Thomassen JM., Kauffman CL., Davison SP. 2006. Keloid pathogenesis and treatment. *Plast Reconstr Surg.* 117: 286-300.
- Alster TS., Williams CM. 1995. Treatment of Keloid sternotomy scars with 585 nm flashlamp-pumped pulsed-dye laser. *Lancet.* 345(8959): 1198-1200.
- Babin RW., Ceilly RI. Combined modalities in the management of hypertrophic scars and keloids. *J Otolaryngol* 1979; 8 : 457-60.
- Bailey JNR., Waite AE., Clayton WJ., Rustin MH. 2007. Application of topical mitomycin C to the base of shave removed keloid scars to prevent their recurrence. *Br J Dermatol.* 156: 628-686.

Bayat A., McGrouther DA., Ferguson M. 2003. Skin scarring. *Br Med J.* 326: 88-92.

Bergman B, Bieleley HC. Keloid. *J A Acad Dermatol.* 2005 ; 33 (10 : 117-23).

Berman B., DUncan MR. 1989. Short term keloid treatment *in vivo* with human interferon alfa-2b results in a selective and persistent normalization of keloidal fibroblast collagen, glycosaminoglycan, and collagenase production *in vitro.* *J Am Acad Dermatol.* 21(4): 694-702.

Berman B., Villa A.M., Ramirez C.C. 2005. Novel opportunities in the treatment and prevention of scarring. *J Cutan Med Surg.* 32-36.

Boyadijev C., Popchristova E., Mazgalova J. 1995. Histomorphologic changes in keloids treated with Kenacort. *J Trauma.* 38(2): 299-302.

Bradner WT. 2001. Mitomycin C: A clinical update. *Cancer Treatment Reviews.* 27: 35-50.

Brent B. 1978. The Role Pressure Therapy in Management of Earlobe Keloids. *Ann Plast Surg.* 1(6): 579-581.

Butler P.D., Longkaker M.T., Yang G.P. 2008. Current Progress in Keloid Research and Treatment. *J Am Coll Surg.* 206:731-741.

Calderon M., Lawrence W.T., Banes A. 1996. Increased Proliferation in Keloid Fibroblast Wounded *in vitro.* *J Surg Res.* 61:343-347.

Carrol LA., Hanasono MM., Mikulec AA., Kita M., Koch RJ. 2002. Triamcinolon stimulates bTGF production and inhibits TGF- β , productin by human dermal fibroblast dermal surgery. *Dermatol Surg*; 28:704-9

Chen MA dan Davidson TM. 2005. Scar Management: Prevention and Treatment Strategies. *Curr Opin Otolaryngol Head Neck Surg*. 13(4): 242-247.

Christophers E. 1974. Growth stimulation of cultured postembryonic epidermal cells by vitamin A acid. *Journal of Investigative Dermatology*. 63: 450-455.

Chung Suyoun, Mitsuko Nakashima, Hitoshi Zembutsu, dan Yusuke Nakamura. 2011. Possible involvement of NEDD4 in Keloid Formation; Its Critical Role in Fibroblast Proliferation and Collagen Production. *Proc Jpn Acad Ser B Phys Biol Sci*. 2011, 87(8): 563-73.

Clark RAF. 1985. Cutaneus tissue repair : Basic biologic consideration 1. *J Am.Acad.Dermatol*; 13 : 701-25

Clugston PA., Vistnes MD., Perry LC., Maxwell GP., Fisher J. 1995. Evaluation of silicone gel sheeting on early wound healing of linear incisions. *Ann Plast Surg*. 34(1): 12-15.

Cohen JK., Diegelmann RF. 1977. The biology of keloid and hypertropic scar and the influence of corticosteroid. *Clinical Plastic Surgery*: 297-310

Curtis AS. dan Seehar GM. 1978. The Control of cell division by tension or diffusion. *Nature*, 274: 52-53.

Eveline Bijlard, Sanne Steltenpool, Frank B. Niessen. 2015. Intralesional 5-Fluorouracil in Keloid Treatment: A Systematic Review. *Acta Derm Venereol*; 95: 778–782

Fitzpatrick. 1999. Treatment of inflamed hypertrophic scars using intralesional 5-FU. *Dermatol Surg*. 25(3): 224-232.

Fitzpatrick TB, Freedberg IM. Eisen AZ., Wolff K., Austen KF., Goldsmith LA., Katz SI., eds. 2003. *Dermatology in general medicine.*, edisi ke 6 New york : Mc Graw Hill:236-45

Fluorouracil. December 2003. Didapat dari : *file://A:\fluorouracil(51—21-8)index from Fluoride Action Network Pesticide Project.htm.*

Freshney RI. 2000. *Primary culture.* Dalam : Culture of animal cells, A manual of basic technique, edisi ke-4. United States : Wiley-Liss Inc.

Friedman D.W., Boyd C.D., Mackenzie J.W., Norton P., Olson R.M., Deak S.B. 1993. Regulation of collagen gene expression in keloids and hypertrophic scars. *J Surg Res*. 55:214-222.

From L, Assaad D. 2003. *Neoplasms, pseudoneoplasms and hiperplasias of the dermis,* Dalam : Fitzpatrick TB, Freedberg IM, Eisen AZ, Wolff K., Austen KF, Goldsmith LA, Katz SI, eds. *Dermatology in general medicine,* edisi ke -6 New York : Mc Graw Hill: 1163-5.

Fujiwara, Y. Muragaki and A. Ooshima. 2005. Keloid-derived fibroblasts show increased secretion of factors involved in collagen turnover and depend on matrix metalloproteinase for migration. *British Journal of Dermatology* 153, pp: 295–300.

Gilchrest BA. 1979. Relationship between actinic damage and cronologic aging in keratinocyte cultures of human skin. *J Invest Dermatol*;72(5):219-23.

Gold MH. 1994. A controlled clinical trial of topical silicone gel sheeting in the treatment of hypertrophic scars and keloids. *J Am Acad Dermatol*. 30(3): 506-507.

Hans I-Chen Harn, Yang-Kao Wang, Chao-Kai Hsu, Yen-Ting Ho, Yi-Wei Huang, Wen-Tai Chiu, Hsi-Hui Lin, Chao-Min Cheng, Ming-Jer Tang. 2015. Mechanical coupling of cytoskeletal elasticity and force generation is crucial for understanding the migrating nature of keloid fibroblasts. *Experimental Dermatology*: 24, 579–584

Harting M., Hicks J.M., Levy M. 2008. Dermal Hypertrophies. In : Wolffk, Goldsmith L.A., Katz S.I., Gilchrest B.A., Paller A.S., Leffel D.J., editor. *Fitzpatrick's Dermatology in General Medicine, 7th ed*. Mc Graw Hill Companies, Inc. 556-558.

Huang C., Murphy G.F., Akaishi S., Ogawa R. 2013. Keloids and hypertrophic scars: Update and future directions. *Plast Reconstr Surg Glob Open*. 1(4): e25.

Huang L., Cai Y.J., Lung I., Leung B.C.S., Burd A. 2013. A study of the combination of triamcinolone and 5-fluorouracil in modulating keloid fibroblast in vitro. *J Plast Reconstr Aesth Surg*. 66: 251-259.

Huang L., Wong Y.P., Cai Y.J., Lung I., Leung C.S., Burd A. 2010. Low-dose 5-fluorouracil induces cell cycle G2 arrest and apoptosis in keloid fibroblasts. *Br J Dermatol*. 163:1181-1185.

Janssen de Limpens AM. 1980. The local treatment of hypertrophic scars and keloids with topical retinoid acid. *Br J Dermatol*. 103: 319-323.

Junqueira LC, Carneiro J. 1982. *Basic histology, edisi ke-3*. California: Lange Medical Publication:89-94.

Junqueira LC, Carneiro J. 2005. *Basic Histology 11th ed*. USA: The McGraw-Hill Companies Inc.

Ketchum LD., Cohen IK., Master FW. 1974. Hypertrophic scars and keloids : a collective review *Plastic Reconstruction Surgery*; 53: 140-54

Khalifa E.S., Adil A.N., Mahmood R.A. 2014. Debulking of Keloid Combined with Intralesional Injection of 5-Fluorouracil and Triamcinolone Versus Intralesional Injection of 5-Fluorouracil and Triamcinolone. *Journal of Cosmetics, Dermatological Sciences and Applications*. 4:85-91.

Kischer CW., Wagner HN., Pindur J., Holubec., Jones M., Ulreich JB., Scuderi P. 1989. Increased fibronectin production by cell lines from hypertrophic scar and keloid. *Connect Tissue Res*. 23:279-288.

Koh HK., Bhawan L., Peled Z., 5-Fluorouracil inhibits keloid fibroblast proliferasi and keloid fibroblast populated collagen lattice contraction. Dipublikasikan 15 januari 2002. Didapat dari : <http://www.journalofburns.com>

Kovalic JJ., Peres CA. 1989. Radiation Therapy Following Keloidectomy: A 20 Year Experience. *Int J Radiat Oncol Biol Phys.* 17(1): 77-80.

Larrabee WF., East CA., Jaffe HS., Stephenson C., Peterson KE. 1990. Intralesional interferon gamma treatment for keloids and hypertrophic scars. *Arch Otolaryngol Head Neck Surg.* 116(10): 1159-1162.

Ledon J.A. *et al.* 2013. Intralesional treatment for keloids and hypertrophic scars: a review. *Dermatol Surg.* 39(12): 1745-1757.

Levinson H, Liu W, Peled Z., 5Fluorouracil inhibits keloid fibroblast proliferation and keloid fibroblast populated collagen lattice contraction. Dipublikasikan 15 januari 2002. Didapat dari : <http://www.journalofburns.com>.

Luo S., Benathan M., Raffoul W., Panizzon RG., Egloff DV. 2001. Abnormal Balance Between Proliferation and Apoptotic Cell Death in Fibroblast Derived from Keloid Lesions. *Plast Reconstr Surg.* 107:87-96.

Machie RM. 1992. Soft Tissue Tumor. Dalam : Champion R.H., Burthorn J.L., Ebling F.J.G. eds. Textbook of dermatology, edisi ke-5. London : Blackwell Scientific Publ: 2073-100.

Manuskiatti W. 2002. Comparison among intralesional corticosteroid, 5-fluorouracil, and 585 nm flashlamp-pumped pulsed-dye laser treatments. *Arch Dermatol.* 138: 1149-1155.

McCoy B.J., Cohen I.K. 1982. Collagenase in keloid biopsies and fibroblasts. *Connect Tissue Res.* 9:181-5.

Mustoe TA., Cooter RD., Gold MH., Hobbs R., Ramelet AA., Shakespeare PG., Stella M., Teot L., Wood FM., Ziegler UE. 2002. International Clinical Recommendations on Scar Managements. *Plast Reconstr Surg.* 110:560-571.

Niessen FB., Spauwen PH., Schalkwijk J., Kon M. 1999. On the nature of hypertrophic scars and keloids: A Review. *Plast Reconstr Surg.* 104 (5): 1435-1458.

Norris JE. 1991. Hypertrophic Scar and Keloid. *Plast Reconstr Surg.* 88(2): 372-373.

Norris JE. 1995. Superficial X-ray Therapy in Keloid Management: A Retrospective Study of 24 Cases and Literature Review. *Plast Reconstr Surg.* 95(6): 1051-1055.

Ogawa R., Mitsuhashi K., Hyakusoku H., dan Miyashita T. 2003. Post operative Electron Beamirradiation Therapy for Keloids and Hypertrophic Scars: Retrospective Study of 147 Cases Followed for More than 18 months. *Plast Reconstr Surg.* 111: 547-553.

Peltonen J., Hsiao L.L, Jaakkola S., Sollberg S., Aumailley M., Timpi R. 1991.

Activation of Collagen Gene Expression in Keloid : Co-Localization of Type I and VI Collagen and Transforming Growth Factor β 1 mRNA. *J Invest Dermatol.* 97: 240-248.

Poochaeron VN and Berman B. 2003. New therapies for the Management of Keloid. *J Craniofac Surg.* 14:654-657.

Pollack SU. 2004. *Management of keloids.* Dalam : Wheeland RG. Cutaneous surgery. Philadelphia : W.B Saunders Co: 688-98.

Puliafito A, Hufnagel, Neveu, Streichan, Sigal, Fygenon, *et al.* 2012. Collective and Single Cell Behavior in Epithelial Contact Inhibition. *Proceedings of the National Academy of Sciences*, 109, 739-744.

Quinn KJ., Evans JH., Courtney JM., Gaylor JD., Reid WH. 1985. Non pressure treatment of hypertrophic scars. *Burns Incl Therm Inj.* 12(2): 102-108.

Ragoowansi R., Cornes PG., Moss AL., Glees JP. 2003. Treatment of Keloids by Surgical Excision and Immediate Postoperative Single Fraction Radiotherapy. *Plast Reconstr Surg.* 111(6): 1853-1859.

Raynolds JEF. 1989. *Corticosteroids* dalam Martindale the extra pharmacopoeia, edisi ke-29. London : The Pharmaceutical Press: 901-2

Ren YM, Zhou XH, Wei ZJ, Lin W, Fan BY, Feng SQ. 2016. Efficacy and safety of triamcinolone acetonide alone and in combination with 5-fluorouracil for

treating hypertrophic scars and keloids: a systematic review and meta-analysis. *Int Wound J*; doi: 10.1111/iwj.12629

Robles D.T., Berg D. 2007. Abnormal wound healing : keloids. *Clin Dermatol.* 25: 26-32.

Scott IU., Greenfield DS., Schiffman J., Nicoleta MT., Rueda JC., Tsai JC., Palmberg PF. 1998. Outcome of primary trabeculectomy with the use of adjunctive mitomycin. *Arch Ophthalmol.* 116: 286-291.

Shaffer JJ, Taylor SC., Boled FC. 2002. Keloid scar : A review with a critical look at therapeutic option. *J Am Acad Dermatol*; 46: S63-97).

Simman R., Alani H., Williams F. 2003. Effect of mitomycin C on keloid fibroblast: An in vitro study. *Ann Plast Surg.* 50: 71-76.

Smith JW, Bellinger CG. 1973. *Keloid and hypertrophic scar* dalam Grabb WC, Smith JW eds. *Plastic surgery a concise guide to clinical practise*, edisi ke-2. Boston : Little Brown & Co; 200 : 740-50

Sproat JE., Dalcin, Weitauer N., Roberts RS. 1992. Hypertrophic sternal scars: silicone gel sheet vs kenalog injection treatment. *Plast Reconstr Surg.* 90(6): 988-992.

Steven PD, Joseph HD, Mark WC., Smitha S., Antai W., dan Amy C. 2009. Efficacy of Intralesional 5-Fluorouracil and Triamcinolone in the Treatment of Keloids. *Aesthetic Surg J* 2009;29:40–46.

Stewart CE., Kim JY., Linda I. 2006. Application of Mitomycin-C for Head and Neck Keloids. *Otolaryngol Head Neck Surg.* 135: 946-950.

Sussman M. 1966. Effect of Increased tissue traction upon tensile strength of Cutaneous incisions in rats. *Proc Soc Exp Biol Med.* 123:38-41.

Talmi YP., Orenstein A., Wolf M., Kronenberg J. 2005. Use of mitomycin C for treatment of keloid: A preliminary report. *Otolaryngol Head Neck Surg.* 132(4): 598-601.

Tredget EE., Nedelec B., Scott PG., Grahary A. 1997. Hypertrophic scars, keloids, and contracture. the cellular and molecular basis for therapy. *Surg Clin North Am.* 77(3): 701-730.

Uitto J, Eisen AZ. Collagen. Dalam : Fitzpatrick TB., Freedberg IM., Eisen AZ., Wolff K., Austen KF., Goledsmith LA., Katz SI. Eds. *Dermatology in general medicine*, edisi ke 6 New york : Mc Graw Hill; 2003 : 165-78)

Ulrich D., Ulrich F., Unglaub F., Platkowski A., Palhua N. 2010. Matrix metalloproteinases and tissue inhibitors of metalloproteinases in patient with different types of scars and keloids. *J Plast Reconst Aesthet Surg.* 63:1015-1021.

Vidhi V. Shah, Sahal S., Adam SA., Keyvan N., Stephanie M., Mohammed A. 2016. 5-Fluorouracil in the Treatment of Keloids and Hypertrophic Scars: A Comprehensive Review of the Literature. *Dermatol Ther (Heidelb)* (2016) 6:169–183

Wirohadidjojo Y.W. dan Dachlan I. 2007. The Effect of Narrow and Broad Band Ultraviolet B on to Keloid Fibroblast-VEGF Expressions. *Berkala Ilmu Kedokteran*. 39(2).

Wirohadidjojo Y.W., Radiono S., Budianto A., Soebono H. 2011. Cellular Viability, Collagen Deposition, and Transforming Growth Factor Beta1 Production among Ultraviolet B - Irradiated keloid Fibroblasts. *Aesthet Plast Surg*. 35(6): 1050-1055.

Wong TW., Chiu HC., Chen JS., Lin LJ., Chang CC. 1995. Symptomatic keloids in two children. Dramatic improvement with silicone cream occlusive dressing. *Arch Dermatol*. 131(7): 775-777.

Yoshino Y, Kubomura K, Ueda H, Tsuge T, Ogawa R. 2018. Extension of flaps associated with burn scar reconstruction: A key difference between island and skin-pedicled flaps. *Burns*. 2018;44(3):683-691