



DAFTAR PUSTAKA

- Amirah, N. binti A. L. (2013). Life Cycle And Ovipositional Preference Of Bactrocera Carambolae Feeding On Starfruits. *Department Zoology*, 1–26. <https://ir.unimas.my/id/eprint/3092/>.
- Astriyani, Kadek Lina., Karlina, N., Supartha, I. W., & Sudiarta, I. P. (2016). *Kemelimpahan Populasi Dan Persentase Serangan Lalat Buah Yang Menyerang Tanaman Buah-Buahan Di Bali*. 5(1), 19–27. Retrieved from <http://ojs.unud.ac.id/index.php/JASB> 19.
- Badan Pusat Statistika Kabupaten Buleleng. (2017). Statistika Hortikultura Kabupaten Buleleng (BPS Kabupaten Buleleng, ed.). <https://www.bulelengkab.go.id/assets/instansikab/82/bankdata/statistik-hortikulturas-tahun-2017-di-kabupaten-buleleng-25.pdf>
- Bernays, Elizabeth A., Chapman, R. F. (1994). Host-Plant Selection by Phytophagous Insects (Bernard D. Roitberg and Murray B. Isman, ed.). <https://www.springer.com/gp/book/9780412031311>
- Broufas, G. D., Pappas, M. L., & Koveos, D. S. (2009). Effect of Relative Humidity on Longevity, Ovarian Maturation, and Egg Production in the Olive Fruit Fly (Diptera: Tephritidae). *Annals of the Entomological Society of America*, 102(1), 70–75. <https://doi.org/10.1603/008.102.0107>
- Chang, C. L., Liang, G. H., & Geib, S. (2017). Proteomic interactions between the parasitoid Diachasmimorpha longicaudata and the oriental fruit fly, Bactrocera dorsalis during host parasitism. *Journal of Asia-Pacific Entomology*. <https://doi.org/10.1016/j.aspen.2017.12.002>
- Chapman, R. F. (1969). *The Insect Structure and Function*. Retrieved from https://www.researchgate.net/profile/Khalid_Khan22/post/What_are_you_r_favorite_entomology_textbooks/attachment/59d6355dc49f478072ea357a/AS%3A273663524311040%401442257876021/download/Chapman+5+th+edition+The+Insects+structure+and+function.pdf
- Commonwealth Agricultural Bureaux International [CABI]. (2020). *Bactrocera dorsalis (oriental fruit fly)*. Tersedia pada <https://www.cabi.org/isc/datasheet/17685> (Diakses pada 24 April 2020)
- Commonwealth Agricultural Bureaux International [CABI]. (2020). *Bactrocera umbrosa*. Tersedia pada <https://www.cabi.org/isc/datasheet/8747>. Diakses pada 24 April 2020)



Dhillon, M. K., Singh, R., Naresh, J. S., & Sharma, H. C. (2005). The melon fruit fly, *Bactrocera cucurbitae*: A review of its biology and management. *Journal of Insect Science*, 5(1). <https://doi.org/10.1093/jis/5.1.40>

Dominiak, B. C., Mavi, H. S., & Nicol, H. I. (2006). Effect of town microclimate on the Queensland fruit fly *Bactrocera tryoni*. *Australian Journal of Experimental Agriculture*, 46(9), 1239–1249. <https://doi.org/10.1071/EA04217>

Doorenweerd, C., Leblanc, L., Jose, M. S., Pham, H. T., & Rubinoff, D. (2018). Descriptions of four new species of *Bactrocera* and new country records highlight the high biodiversity of fruit flies in Vietnam (Diptera, Tephritidae, Dacinae). *ZooKeys*, 2018(797), 87–115. <https://doi.org/10.3897/zookeys.797.29138>

Drew, Richard A.I . and Meredith C. Romig. (2013). *Tropical Fruit Flies Of South-East Asia* (Nigel, Farrar, Alexandra Lainsbury, and Shankari Wilford, ed.). United Kingdom: CABI.

Duyck, P. F., David, P., & Quilici, S. (2004). A review of relationships between interspecific competition and invasions in fruit flies (Diptera: Tephritidae). *Ecological Entomology*, 29(5), 511–520. <https://doi.org/10.1111/j.0307-6946.2004.00638.x>

García Medel, D., Sivinski, J., Díaz-Fleischer, F., Ramirez-Romero, R., & Aluja, M. (2007). Foraging behavior by six fruit fly parasitoids (Hymenoptera: Braconidae) released as single- or multiple-species cohorts in field cages: Influence of fruit location and host density. *Biological Control*, 43(1), 12–22. <https://doi.org/10.1016/j.biocontrol.2007.06.008>

Garcia, F. R. M., & Ricalde, M. P. (2013). Augmentative biological control using parasitoids for fruit fly management in Brazil. *Insects*, 4(1), 55–70. <https://doi.org/10.3390/insects4010055>

Ginting, R. (2009). Keanekaragaman Lalat Buah Diptera (Tephritidae) Di Jakarta, Depok, Dan Bogor Sebagai Bahan Kajian Penyusunan Analisis Risiko Hama (Institut Pertanian Bogor). Retrieved from <https://repository.ipb.ac.id/handle/123456789/5263>

Hafizah, N. E. A. . Z. Y. S. (2016). *Keanekaragaman Spesies Tumbuhan Di Arealnilai Konservasi Tinggi (Nkt) Perkebunan Kelapa Sawit Provinsi Riau (The Diversity Of Plant Species In High Conservation Value Area of Oil Palm Plantation in Riau Province)*. 21(1), 91–98. Retrieved from <https://journal.ipb.ac.id/index.php/konservasi/article/view/13956>



Hasyim, A., Muryati, M., & Kogel, W. J. De. (2016). Population Fluctuation Of Adult Males Of The Fruit Fly, *Bactrocera* Tau Walker (Diptera: Tephritidae) In Passion Fruit Orchards In Relation To Abiotic Factors And Sanitation. *Indonesian Journal of Agricultural Science*, 9(1), 29. <https://doi.org/10.21082/ijas.v9n1.2008.p29-33>

Herlinda, S., Zuroaidah, Y. Pujiastuti, S. Samad, dan T. A. (2008). Spesies Lalat Buah yang Menyerang Sayuran Solanaceae dan Cucurbitaceae di Sumatera Selatan. *J. Hort*, 18(2), 212–220. <http://ejurnal.uin-suska.ac.id/index.php/agroteknologi/article/download/58/48>

Huan, L. I. U., Dong-ju, Z., Yi-juan, X. U., Lei, W., Dai-feng, C., Yi-xiang, Q. I., ... Yong-, L. U. (2019). *Invasion, expansion, and control of Bactrocera dorsalis (Hendel) in China*. (April). [https://doi.org/10.1016/S2095-3119\(18\)62015-5](https://doi.org/10.1016/S2095-3119(18)62015-5)

Huston, M. (1979). A General Hypothesis of Species Diversity. *The American Naturalist*, 113(1), 81–101. <https://doi.org/10.1086/283366>

Huang, Y. B. K., & Chi, H. (2014). Fitness of *Bactrocera dorsalis* (Hendel) on seven host plants and an artificial diet. *Turkiye Entomoloji Dergisi*, 38(4), 401–414. <https://doi.org/10.16970/ted.08838>

Indriyanti, D. R. R. F. F. (2014). Keanekaragaman parasitoid dan predator *Bactrocera* (Diptera : Tephritidae) Di Kabupaten Demak. *Agrineca*, 14. Retrieved from [https://lib.unnes.ac.id/30757/1/Keanekaragaman_parasitoid_dan_predator_Bactrocera_\(Diptera__Tephritidae\)_Di_Kabupaten_Demak.pdf](https://lib.unnes.ac.id/30757/1/Keanekaragaman_parasitoid_dan_predator_Bactrocera_(Diptera__Tephritidae)_Di_Kabupaten_Demak.pdf)

Kraft, K. H., Brown, C. H., Nabhan, G. P., Luedeling, E., De Jesús Luna Ruiz, J., D'Eeckenbrugge, G. C., Gepts, P. (2014). Multiple lines of evidence for the origin of domesticated chili pepper, *Capsicum annuum*, in Mexico. *Proceedings of the National Academy of Sciences of the United States of America*, 111(17), 6165–6170. <https://doi.org/10.1073/pnas.1308933111>

Kuswadi, A. N., I. A. Nasution, M. Indrawatmi dan Darmawi. (1999). Pembibitan Massal Lalat Buah *Bactrocera carambolae* (Drew and Hancock) Dengan Makanan Buatan. Dalam Kumpulan Laporan Akhir Pelaksanaan RUT VI 1997/2000. Badan Tenaga Nuklir Nasional. Jakarta.

Kuswadi, N. dan A. N. (2004). Perendaman Telur, Penggunaan Suhu Rendah dan Aerasi Untuk Perbaikan Pembibitan Massal Lalat Buah *Bactrocera carambolae* (Drew & Hancock) Dalam Teknik Serangga Mandul. *Risalah Seminar Ilmiah Penelitian Dan Pengembangan Aplikasi Isotop Dan Radiasi*. <http://repo-nkm.batan.go.id/7164/>



Li, C., Hirano, H., Kasajima, I., Yamagishi, N., & Yoshikawa, N. (2019). Virus-induced gene silencing in chili pepper by apple latent spherical virus vector. *Journal of Virological Methods*, 273, 113711. <https://doi.org/10.1016/j.jviromet.2019.113711>

Maguran, A. E. (1988). *Ecological Diversity and Its Measurement* (First). Retrieved from <https://www.springer.com/gp/book/9789401573603>

Manurung, B., Prastowo, P., & Tarigan, E. E. (2012). *Bactrocera Dorsalis Complex Pada Pertanaman Jeruk Di Dataran Tinggi Kabupaten Karo Provinsi Sumatera Utara*. 12(2), 103–110. <http://jhpttropika.fp.unila.ac.id/index.php/jhpttropika/article/view/99>

Meats, A., Holmes, H. M., & Kelly, G. L. (2004). Laboratory adaptation of Bactrocera tryoni (Diptera: Tephritidae) decreases mating age and increases protein consumption and number of eggs produced per milligram of protein. *Bulletin of Entomological Research*, 94(6), 517–524. <https://doi.org/10.1079/ber2004332>

Motswagole, R., Gotcha, N., & Nyamukondiwa, C. (2019). Thermal Biology and Seasonal Population Abundance of Bactrocera dorsalis Hendel (Diptera: Tephritidae): Implications on Pest Management. *International Journal of Insect Science*, 11, 117954331986341. <https://doi.org/10.1177/1179543319863417>

Nishida, R. K. H. T. H. F. (1988). *CIS-3,4-Dimethoxycinnamyl Alcohol From The Rectal Glands Of Male Oriental Fruit Fly, Dacus Dorsalis*. 3, 4. Retrieved from https://www.academia.edu/19120940/Cis-3_4-dimethoxycinnamyl_alcohol_from_the_rectal_glands_of_male_oriental_fruit_fly_Dacus_dorsalis

Oka, I.N.(1995). *Pengendalian Hama Terpadu dan Implementasinya di Indonesia*. Yogyakarta: Gadjah Mada University Press. 255hal.

Olievera, N. Wayan Susila; Wayan Supartha. (2016). Keragaman Jenis Lalat Buah dan Tingkat Parasitasi Parasitoid yang Berasosiasi dengan Tanaman Buah-Buahan di Distrik Lautem , Timor Leste. 5(February), 93–102. https://www.researchgate.net/publication/292966665_Jenis_Lalat_Buah_dan_Tingkat_Parasitasi_Parasitoid_yang_Berasosiasi_dengan_Tanaman_Buah-Buahan_di_Distrik_Lautem_Timor_Leste

Plant Health Australia. (2018). *The Australian Handbook For The Identification of Fruit Flies*. <https://doi.org/10.1016/j.jasms.2007.01.008>

Santiatma, I Made Yesta., I Ketut Sumiartha., I Wayan Susila., Made Suparta Utama., Joko Maryono., & Greg Luther. (2016). Identifikasi Lalat Buah (Diptera: Tephritidae) serta Serangannya terhadap Beberapa Galur dan



Varietas Tanaman Cabai (*Capsicum annum L.*) Di Desa Pancasari, Sukasada, Buleleng. *E-Jurnal Agroekoteknologi Tropika (Journal of Tropical Agroecotechnology)*, 5(1), 11–19. Retrieved from <https://ojs.unud.ac.id/index.php/JAT/article/view/18206>

Serit and Tan. (1990). Immature life table of natural population of *Dacus dorsalis* in a village ecosystem (pp. 305–330). *Tropical Pest Management*, 10, 36(3)

http://psasir.upm.edu.my/id/eprint/3198/1/Biology_of_Diachasmimorpha_longicaudata,.pdf

Siwi, S.S, Purnama Hidayat, S. (2006). Taksonomi dan Bioekologi Lalat buah penting *Bactrocera* spp. (Diptera: Tephritidae) di Indonesia. In *Balai Besar Penelitian dan Pengembangan Bioteknologi dan Sumberdaya Genetik Pertanian*.
<https://www.researchgate.net/publication/330102477%0ATaksonomi>

Solangi, B., R. Sultana, N. Ahmed, M. S. Waganand, & Abbassi, A. R. (2014). Seasonal monitoring of fruit fly, *Bactrocera zonata* (Saunders) and its parasitoids Trybliographa adaci on guava fruit from Sindh Pakistan. *J. Bio. Environ. Sci.*, 4(5), 222–232.
<http://www.innspub.net/wp-content/uploads/2014/05/JBES-Vol4No5-p222-232.pdf>

Suputa, Cahyaniati, Arminudin, A. T., Kustaryati, A., Railan, M., & Issusilaningtyas. (2007). *Pedoman Koleksi dan Preservasi Lalat Buah (Diptera : Tephritidae)*.
<https://www.researchgate.net/publication/321184130%0APedoman>

Suputa, Ahmad Taufiq Arminudin, Palupi Jatuasri, I. P. R. (2007). Tingkat Parasitasi *Fopius Arisanus* Hymenoptera: Braconidae) Pada Lalat Buah Belimbing Di Daerah Istimewa Yogyakarta. *Jurnal Perlindungan Tanaman Indonesia*, 13, 8. Retrieved from https://www.researchgate.net/publication/200642117_Effect_of_Relative_Humidity_on_Longevity_Ovarian_Maturation_and_Egg_Production_in_the_Olive_Fruit_Fly_Diptera_Tephritidae

Suputa, Y. Andi Trisyono, Edhi Martono, and S. S. S. (2010). Update on The Host Range of Different Species of Fruit Flies in Indonesia. *Jurnal Perlindungan Tanaman Indonesia*, 16(2), 62–75.
<https://doi.org/10.22146/jpti.11725>

Sushil Chhapekar, M. K. and N. R. (2016). Advances in Molecular Breeding of Pearl Millet. (January), 397–419. https://doi.org/10.1007/978-3-319-27090-6_15.
https://www.researchgate.net/publication/301545608_Advances



Tan, K., Nishida, R., & Toong, Y. (2002). Floral Synomone Of A Wild Orchid , *Bulbophyllum Cheiri* , Lures Bactrocera Fruit Flies For Pollination. *Journal Of Chemical Ecology*, 28(6). Retrieved From <https://link.springer.com/article/10.1023/A:1016277500007>

Vargas, R. I., Piñero, J. C., & Leblanc, L. (2015). An overview of pest species of Bactrocera fruit flies (Diptera: Tephritidae) and the integration of biopesticides with other biological approaches for their management with a focus on the pacific region. *Insects*, 6(2), 297–318. <https://doi.org/10.3390/insects6020297>

Vayssières, J. F., De Meyer, M., Ouagoussounon, I., Sinzogan, A., Adandonon, A., Korie, S., ... Goergen, G. (2015). Seasonal abundance of mango fruit flies (Diptera: Tephritidae) and ecological implications for their management in mango and cashew orchards in Benin (Centre & North). *Journal of Economic Entomology*, 108(5), 2213–2230. <https://doi.org/10.1093/jee/tov143>

Verghese, Abraham; DK Nagaraju, HS Madhura, PD Kamalajayanthi, S. (2006). *Wind speed as an independent variable to forecast the trap catch of the fruit fly (Bactrocera dorsalis) speed as an independent variable forecast the trap catch of the fruit fly (Bactrocera dorsalis)*. 76(July 2018).

Warduna, R, WT Retno, Desmawati, P Warastin, T Sulistyawati, dan CR Hidayat. (2015). Metode Pengamatan Organisme Pengganggu Tumbuhan Tanaman Hortikultura. Direktorat Jenderal Hortikultura. Jakarta.

Ye, H., & Liu, J. H. (2005). Population dynamics of the oriental fruit fly, *Bactrocera dorsalis* (Diptera: Tephritidae) in the Kunming area, southwestern China. *Insect Science*, 12(5), 387–392. <https://doi.org/10.1111/j.1005-295X.2005.00048.x>

Yones, M. S., Abd ElRahman., Abou Hadid, Arafat., Dahi (2011). Heat Unit Requirements for Development of the Pink Bollworm *Pectinophora gossypiella* (Saund.). *Egyptian Academic Journal of Biological Sciences. A, Entomology*, 4(1), 115–122. <https://doi.org/10.21608/eajbsa.2011.15176>