

- Alkhateeb and J. R. Connor, "The significance of ferritin in cancer: Anti-oxidation, inflammation and tumorigenesis," *Biochimica et Biophysica Acta (BBA) - Reviews on Cancer*, vol. 1836, no. 2, pp. 245–254, 2013.
- Andrews, N. C. "Disorders of iron metabolism," *The New England Journal of Medicine*, vol. 341, no. 26, pp. 1986–1995, 1999.
- Ashraf Mikhail, Christopher Brown, Jennifer Ann Williams, Vinod Mathrani, Rajesh Shrivastava, Jonathan Evans, Hayleigh Isaac and Sunil Bhandari "Renal association clinical practice guideline on Anaemia of Chronic Kidney Disease", BMC Nephrology 2017.
- Bailie GR, Larkina M, Goodkin DA, Li Y, Pisoni RL, Bieber B, et al, *Variation in intravenous iron use internationally and over time: the dialysis outcomes and practice patterns study (DOPPS)*. Nephrol Dial Transplant. 2013.
- Black, J.M, Hawks, J.H. 2005. *Medical Surgical Nursing Clinical Management for Positive Outcome 7th edition*. Philadelphia : W.B Saunders Company.
- Cepeda-Lopez, J. Allende-Labastida, A. Melse-Boonstra et al., "The effects of fat loss after bariatric surgery on inflammation, serum hepcidin, and iron absorption: A prospective 6-mo iron stable isotope study," *American Journal of Clinical Nutrition*, vol. 104, no. 4, pp. 1030–1038, 2016
- Cronin, R.E., Henrich, W.L. 2010. *Kt/V and The Adequacy of Hemodialysis*. <http://www.ncbi.nlm.nih.gov/pubmed/9807323>
- Dalimunthe N, *Peningkatan Nilai Parameter Status Besi Reticulocyte Hemoglobin Equivalent setelah pemberian suplemen besi intravena pada pasien hemodialisis reguler*. Medan: Universitas Sumatera Utara. 2012.
- Datz, E. Muller, and E. Aigner, "Iron overload and non-alcoholic fatty liver disease," *Minerva Endocrinologica*, vol. 42, no. 2, pp. 173–183, 2017.
- Daugirdas, J.T., Blake, P.G., Ing, T.S. 2007. *Handbook of Dialysis 4th edition*. Philadelphia: Lippincott.
- Descamps, Latscha B., Goldford, B., Nguyen, A. T., 1991. *Establishing the relationship between complement activation and stimulation of phagocyte oxidative metabolism in hemodialysed patients: a randomized prospective study*. Nephron, 59:279-285.

Dewi, I.G.A.P.A. 2010. *Hubungan antara Quick of Blood/Qb dengan Adekuasi Hemodialisis di Ruang HD BRSU Daerah Tabanan Bali*. Tesis. Tidak dipublikasikan.

Eknoyan, G., Levin, N.W., Eschbach, J.W. 2000. *KDOQI Clinical Practice Guidelines*. <http://www.asnjournals.org>

Erwinsyah (2009). *Hubungan antara Qb dengan penurunan kadar ureum dan kreatinin plasma pada pasien hemodialisis di RSUD Raden Mattaher Jambi*. Tesis. Tidak dipublikasikan.

Fairweather-Tait, A. A. Wawer, R. Gillings, A. Jennings, and P. K. Myint, "Iron status in the elderly," *Mechanisms of Ageing and Development*, vol. 136-137, pp. 22–28, 2014.

Fernandez-Rodriguez AM, Guindeo-Casasus MC, Molero-Labarta T, et al. "Diagnosis of iron deficiency in chronic renal failure". *Am J Kidney Dis*.1999.

Fishbane S, Kowalski EA, Imbriano LJ, Maesaka JK: The evaluation of iron status in hemodialysis patients. *J Am Soc Nephrol* 7:2654–2657, 1996

Forouhi NG, Harding AH, Allison M, Sandhu MS, Welch A, Luben R, et al. *Elevated serum ferritin levels predict newonset type 2 diabetes*, results from the EPIC-Norfolk prospective study. *Diabetologia* 2007;50:949–56.

Gatot, D. 2003. *Rasio Reduksi Ureum dalam Dialisis*. <http://library.usu.ac.id/download/fk/penyakitdalam-dairot%20gatot.pdf>

Guo, P.-C. Chen, and W.-S. Ko, "Status of essential trace minerals and oxidative stress in viral hepatitis C patients with nonalcoholicfatty liverdisease," *InternationalJournalofMedical Sciences*, vol. 10, no. 6, pp. 730–737, 2013.

Hasuike Y, Nonoguchi H, Tokuyama M, Ohue M, Nagai T, Yahiro M, et al, *Serum ferritin predicts prognosis in hemodialysis patients: the Nishinomiya study*. *Clin Exp Nephrol*. 2010;14:349–55

Holyoake, D. J. Stott, P. J. McKay, A. Hendry, J. B. MacDonald, and N. P. Lucie, "Use of plasma ferritin concentration to diagnose iron deficiency in elderly patients," *Journal of Clinical Pathology*, vol. 46, no. 9, pp. 857–860, 1993.

Indonesian renal registry. 2015. 8th Report Of Indonesian Renal Registry.

Iwanaga S, Sakano N, Taketa K, Takahashi N, Wang DH, Takahashi H, et al *Comparison of serum ferritin and oxidative stress biomarkers between Japanese workers with and without metabolic syndrome*. *Obes Res Clin Pract* 8: e271–282, 2014

Jeong Min Seong¹, Yo Sang Yoon, Kyu Su Lee, Nan Young Bae, Mi Young Gi, Hyun Yoon, *Gender difference in relationship between serum ferritin and 25-hydroxyvitamin D in Korean adults*, 2014

Jindal, K., Chan, C.T. 2006. *Hemodialysis adequacy in Adult*. Journal of The American Society of Nephrology. 2006: 17:4-7.

KDOQI guidelines on anaemia management: http://www2.kidney.org/professionals/KDOQI/guidelines_anemia/guide1.htm. Accessed 15 June 2017

K/DOQI. 2006. *Clinical practice guidelines for hemodialysis adequacy*. Maret 2, 2010. <http://www.kidney.org/Professionals/kdoqi>.

Kalantar-Zadeh K, Don BR, Rodriguez RA, Humphreys MH: *Serum ferritin is a marker of morbidity and mortality in hemodialysis patients*. *Am J Kidney Dis* 37:564–572, 2001

Kalantar-Zadeh K, Rodriguez RA, Humphreys MH: *Association between serum ferritin and measures of inflammation, nutrition and iron in haemodialysis patients*. *Nephrol Dial Transplant* 19:141–149, 2004

Kalantar-Zadeh K, Hoffken B, Wunsch H, Fink H, Luft FC, et al. “Diagnosis of iron deficiency anaemia in renal failure patients during the post erythropoietin”. *Am J Kidney Dis*. 1995.

Kee JL. Laboratory and diagnostic tests with nursing implication. In: Kapoh RP, Kurnianingsih S, Widyastuti P, Chyaningrum R, Rahayu S, editors *Bahasa Indonesia. Pedoman Pemeriksaan Laboratorium & Diagnostik (6th ed)*. Jakarta: EGC, 2013; p. 194.

Kowdley, P. Belt, L. A. Wilson et al., “Serum ferritin is an independent predictor of histologic severity and advanced fibrosis in patients with nonalcoholic fatty liver disease,” *Hepatology*, vol. 55, no. 1, pp. 77–85, 2012.

Li, F.K., Lai, K.N. 2000. *Current Issues in the Chronic Renal failure : Dialysis*. Havas Media

Martin LA, Critelli JW, Doster JA, Powers C, Purdum M, Doster MR, et al. *Cardiovascular risk: gender differences in lifestyle behaviors and coping strategies*. *Int J Behav Med* 20: 97–105. <https://doi.org/10.1007/s12529-011-9204-3> PMID: 22144263, 2013.

Mavromatidis, C. Fytil, P. Kynigopoulou, T. Fragia, and K. Sombolos, “Serum ferritin levels are increased in patients with acute renal failure,” *Clinical Nephrology*, vol. 49, no. 5, pp. 296–298, 1998.

Muhammad, A. and Sianipar, O. (2005), *Penentuan Defisiensi Besi Anemia Penyakit Kronis Menggunakan Peran Indeks sTfR-F*, Indonesian Journal of Clinical Pathology and Medical Laboratory, 12(1), pp. 9– 15. doi: 10.3390/s100706535

Moreno-Navarrete, M. Moreno, J. Puig et al., “Hepatic iron content is independently associated with serum hepcidin levels in subjects with obesity,” *Clinical Nutrition*, vol. 36, no. 5, pp. 1434–1439, 2017.

Moreno-Navarrete, G. Blasco, G. Xifra et al., “Obesity is associated with gene expression and imaging markers of iron accumulation in skeletal muscle,” *The Journal of Clinical Endocrinology & Metabolism*, vol. 101, no. 3, pp. 1282–1289, 2016.

Nakanishi T, Kuragana T, Nanami M, Otaki Y, Nonoguchi H, Hasuike Y. *Importance of ferritin for optimizing anemia therapy in chronic kidney disease*. Am J Nephrol. 2010;32:439-46.

National Kidney Foundation. 2001. *Guidelines for Hemodialysis adequacy*. <http://www.kidney.org/Professionals/kdoqi/>

Owen, W.F. 2000. *Hemodialysis Adequacy*. <http://www.oxfordjournals.org>.

Pernefri. 2003. *Konsensus Dialisis Perhimpunan Nefrologi Indonesia*. Jakarta

Porth, M.C., Matfin, G., 2009. *Pathophysiology Concepts of altered Health States*. 8th ed. Philadelphia : Lippincott Williams & Wilkins.

Pourfarzianu, V., Ghanbarpour, F., Nemati, E., Taheri, S., Einollahi, B. 2008. Laboratory variables and Treatment Adequacy in Hemodialysis Patient in Iran. *Saudi Journal of Nursing Scholarship Volume 42. Issue 3. Pages 242-249*.

Raku Son, Takuya Fujimaru, Takeshi Kimura, Fumika Taki, Miyuki Futatsuyama, Masahiko Nagahama, Masaaki Nakayama and Yasuhiro Komatsu, *Association between serum ferritin levels and clinical outcomes in maintenance hemodialysis patients: a retrospective single-center cohort study*, Renal Replacement Therapy (2019) 5:17.

Reis KA, Guz G, Ozdemir H, Erten Y, Atalay V, Bicik Z, et al, *Intravenous iron therapy as a possible risk factor for atherosclerosis in end-stage renal disease*. Int Heart J. 2005;46:255 –64. 20.

Roesli, R. 2005. *Bila ginjal Aus*. <http://www.solusi.kesehatan.com/penyakit-ginjal/bila-ginjal-aus.html>.

Salgia R. J. and K. Brown, “Diagnosis and management of hereditary hemochromatosis,” *Clinics in Liver Disease*, vol. 19, no. 1, pp. 187–198, 2015.

Santoso. 2010. *Gagal Ginjal Kronik*. <http://antiloans.org>

Sebastiani, B. Tyagarajan, F. Sun et al., "Age and sex distributions of age-related biomarker values in healthy older adults from the long life family study," *Journal of the American Geriatrics Society*, vol. 64, no. 11, pp. e189–e194, 2016.

Septiwi. C. 2010. *Hubungan antara adekuasi hemodialisis dengan kualitas hidup pasien hemodialisis di unit hemodialisis RS. Prof. Dr. Margono Soekarjo Purwokerto*. Tesis. Tidak dipublikasikan.

Simatupang, T. 2006. *Gangguan kardiovaskular pada penderita Ginjal*. <http://litbang.depkes.go.id/aktual/kliping/ginjal/250406.htm>

Suhardjono. 2014. *Gagal Ginjal Kronik* dalam Buku ajar ilmu penyakit dalam. *edisi ke-4. Jilid I*. Jakarta : Perhimpunan Dokter Spesialis Penyakit Dalam Indonesia.

Suharjono. 2010. *Penderita gagal Ginjal di Indonesia*. 222.ikcc.or.id/content.php?c=1&id=275

Sun-Ah You, Qing WangT, *Ferritin in atherosclerosis*, Center for Molecular Genetics, Department of Molecular Cardiology, ND4-38, Lerner Research Institute, and Center for Cardiovascular Genetics, The Cleveland Clinic Foundation, 9500 Euclid Ave., Cleveland, OH 44195, United States Received 6 December 2004.

Suwitra, Ketut. 2014. *Gagal Ginjal Kronik* dalam Buku ajar ilmu penyakit dalam. *edisi ke-4. Jilid I*. Jakarta : Perhimpunan Dokter Spesialis Penyakit Dalam Indonesia.

Stein, S. Connor, G. Virgin, D. E. H. Ong, and L. Pereyra, "Anemia and iron deficiency in gastrointestinal and liver conditions," *World Journal of Gastroenterology*, vol. 22, no. 35, pp. 7908–7925, 2016.

Teehan GS, Ruthazer R, Balakrishnan VS, Snyderman D, Jaber BL, *Iron storage indices and risk of bacterial infections in hemodialysis patients*, *Hemodial Int*. 2004;8:226–32.

Victor, H. Evgeniy, T. Gergana et al., "Serum hepcidin levels in multiple myeloma," *Clinical Laboratory*, vol. 63, no. 7, pp. 1273–1277, 2017.

Wasse. 2007. *Association of initial hemodialysis vascular Access with Quality of Life*. <http://cjasn.asnjournals.org>

WHO, *Serum Ferritin Concentrations for the Assessment of Iron Status and Iron Deficiency in Populations*. Vitamin and Mineral Nutrition Information System, World Health Organization, Geneva, 2011.

Yendriwati. *Status besi pada penderita gagal ginjal kronik dalam menentukan diagnosa anemia*. Jurnal Universitas Sumatera Utara. 2008.

Zhao, L. X. Zhang, Y. Shen, X. Fang, Y. Wang, and F. Wang, "Obesity and iron deficiency: a quantitative meta-analysis," *Obesity Reviews*, vol. 16, no. 12, pp. 1081–1093, 2015.