

- Ayrian, E., Kaye, A. D., Varner, C. L., Guerra, C., Vadivelu, N., Urman, R. D., Zelman, V., Lumb, P. D., Rosa, G., & Bilotta, F. (2015). *Effects of Anesthetic Management on Early Postoperative Recovery, Hemodynamics and Pain After Supratentorial Craniotomy*. Journal of Clinical Medicine Research. <https://doi.org/10.14740/jocmr2256w>
- Bala, I., Gupta, B., Vhardwaj, N., Ghai, B., & Khosla, V. K. (2006). Effect of scalp block on postoperative pain relief in craniotomy patients. *Anaesthesia and Intensive Care*, 34(2), 224–227.
- Bisri, T., & Yulianti, D. (2016). Anestesi untuk Operasi Tumor Supratentorial. In *Anestesi untuk Operasi Tumor Otak Supratentorial, Infratentorial* (1st ed., pp. 41–90). Fakultas Kedokteran Universitas Padjajaran.
- Castro, M. G., Cowen, R., Williamson, I. K., David, A., Jimenez-Dalmaroni, M. J., Yuan, X., Bigliari, A., Williams, J. C., Hu, J., & Lowenstein, P. R. (2003). Current and future strategies for the treatment of malignant brain tumors. *Pharmacology and Therapeutics*, 98(1), 71–108. [https://doi.org/10.1016/S0163-7258\(03\)00014-7](https://doi.org/10.1016/S0163-7258(03)00014-7)
- Charles, P. D., Cécile, L., Chantal, L., Jean-Michel, N., Karim, A., & Corinne, L. (2015). 0.5% levobupivacaine versus 0.5% ropivacaine: Are they different in ultrasound-guided sciatic block? *Saudi Journal of Anaesthesia*, 9(1), 3–8. <https://doi.org/10.4103/1658-354X.146250>
- Chowdhury, T., Garg, R., Sheshadri, V., Venkatraghavan, L., Bergese, S. D., Cappellani, R. B., & Schaller, B. (2017). Perioperative Factors Contributing the Post-Craniotomy Pain: A Synthesis of Concepts. *Frontiers in Medicine*, 4(March), 1–5. <https://doi.org/10.3389/fmed.2017.00023>
- Dunn, L. K., Naik, B. I., Nemergut, E. C., & Durieux, M. E. (2016). Post-Craniotomy Pain

<https://doi.org/10.1007/s11910-016-0693-y>

- Gélinas, C., & Johnston, C. (2007). Pain assessment in the critically ill ventilated adult: Validation of the critical-care pain observation tool and physiologic indicators. *Clinical Journal of Pain*, 23(6), 497–505. <https://doi.org/10.1097/AJP.0b013e31806a23fb>
- Guilfoyle, M. R., Helmy, A., Duane, D., & Hutchinson, P. J. A. (2013). Regional scalp block for postcraniotomy analgesia: A systematic review and meta-analysis. *Anesthesia and Analgesia*, 116(5), 1093–1102. <https://doi.org/10.1213/ANE.0b013e3182863c22>
- Inez M, M., Yulianti Bisri, D., & Adam, A. (2015). Insidensi Tumor Supratentorial berdasarkan Jenis dan Letaknya di RSUP Dr. Hasan Sadikin Tahun 2012-2013. *Jurnal Neuroanestesi Indonesia*, 4(3), 157–161. <https://doi.org/10.24244/jni.vol4no3.94>
- Karim, S. A., Shuer, L. M., & Chang, S. D. (2014). Craniotomy for Tumor. In R. A. Jaffe, C. A. Schmiesing, & B. Golianu (Eds.), *Anesthesiologist's Manual of Surgical Procedure, 5th Edition* (5th ed., pp. 29–34). Wolters Kluwer Health.
- Liisanantti, O., Luukkonen, J., & Rosenberg, P. H. (2004). High-dose bupivacaine, levobupivacaine and ropivacaine in axillary brachial plexus block. *Acta Anaesthesiologica Scandinavica*, 48(5), 601–606. <https://doi.org/10.1111/j.0001-5172.2004.00393.x>
- Malav, K., Singariya, G., Mohammed, S., Kamal, M., Sangwan, P., & Paliwal, B. (2018). Comparison of 0.5% ropivacaine and 0.5% levobupivacaine for sciatic nerve block using labat approach in foot and ankle surgery. *Turk Anesteziyoloji ve Reanimasyon Dernegi Dergisi*, 46(1), 15–20. <https://doi.org/10.5152/TJAR.2017.03411>
- Ostrom, Q. T., Gittleman, H., Truitt, G., Boscia, A., Kruchko, C., & Barnholtz-sloan, J. S. (2018). *Neuro-Oncology CBTRUS Statistical Report : Primary Brain and Other Central Nervous System Tumors Diagnosed in the United States in 2011-2015*. 20(S4), 1–86.

- Papangelou, A., Radzik, B. R., Smith, T., & Gottschalk, A. (2013). A review of scalp blockade for cranial surgery. *Journal of Clinical Anesthesia*, 25(2), 150–159.  
<https://doi.org/10.1016/j.jclinane.2012.06.024>
- Pin-on, P., & Punjasawaswong, Y. (2016). Effect of pre-incisional anterior scalp block on intraoperative opioid consumption in adult patients undergoing elective craniotomy to remove tumor: study protocol for a randomized double-blind trial. *Asia Pacific Clinical and Translational Nervous System Diseases*, 1(3), 131. <https://doi.org/10.4103/2468-5577.187079>
- Sastroasmoro, S., & Ismael, S. (2014). *Dasar Dasar Metodologi Penelitian Klinis* (5th ed.). Binarupa Aksara.
- Sessler, C. N., Grap, M. J., & Ramsay, M. A. (2008). Evaluating and monitoring analgesia and sedation in the intensive care unit. *Critical Care*, 12(Suppl 3), S2.  
<https://doi.org/10.1186/cc6148>
- Sripada, R., Garg, S., & Raeder, J. (2015). Regional anesthesia for ambulatory surgery. *Practical Ambulatory Anesthesia*, 8180(95), 61–78.  
<https://doi.org/10.1017/CBO9781107588219.007>
- Theerth, K. A., Sriganesh, K., & Reddy, K. M. (2018). *Analgesia Nociception Index-guided intraoperative fentanyl consumption and postoperative analgesia in patients receiving scalp block versus incision-site infiltration for craniotomy*. 84(12), 1361–1368.  
<https://doi.org/10.23736/S0375-9393.18.12837-9>
- Tuchinda, L., Somboonviboon, W., Supbornsug, K., Worathongchai, S., & Limutaitip, S. (2010). Bupivacaine scalp nerve block: Hemodynamic response during craniotomy, intraoperative and post-operative analgesia. *Asian Biomedicine*, 4(2), 243–251.
- Vadivelu, N., Kai, A. M., Tran, D., Kodumudi, G., Legler, A., & Ayrian, E. (2016). Options

<https://doi.org/10.2147/JPR.S85782>

Wardhana, A., & Sudadi, S. (2019). Scalp block for analgesia after craniotomy: A meta-analysis. *Indian Journal of Anaesthesia*, 63(11), 886–894.

[https://doi.org/10.4103/ija.IJA\\_315\\_19](https://doi.org/10.4103/ija.IJA_315_19)

Yang, X., Ma, J., Li, K., Chen, L., Dong, R., Lu, Y., Zhang, Z., & Peng, M. (2019). A comparison of effects of scalp nerve block and local anesthetic infiltration on inflammatory response , hemodynamic response , and postoperative pain in patients undergoing craniotomy for cerebral aneurysms : a randomized controlled trial. *BMC Anesthesiology*, 19:91, 1–11.