

## DAFTAR PUSTAKA

- Adawiyah, D., Wisetsombat, M., & Chueamchaitrakun, P. (2017). The Correlation Between Sensory and Physicochemical of Green Tea. *Jurnal Mutu Pangan*, 65-69.
- Al-Farsi, M., Alasalvar, C., Al-Abid, M., Al-Shoaily, K., Al-Amry, M., & Al-Rawahy, F. (2007). Compositional and Functional Characteristics of Dates, Syrups, and Their By-Products. *Food Chemistry* 104, 943-947.
- An, C., & Mou, Z. (2011). Salicylic Acid and Its Function in Plant Immunity. *J. Integr.Plant.Biol.*53, 412-428.
- Andersen, C. (2003). Source-sink Balance and Carbon Allocation Below Ground in Plants Exposed to Ozone. *New Phytol*, 2013-228.
- Anggai, R., Hasan, H., & Thomas, N. (2015). Preparations Microemulsion Formulation and Evaluation of Ethanol Extract Red Rice (*Oryza nivara*) As Antioxidants.
- Anggraini, T. (2017). *Proses dan Manfaat Teh*. Padang: Erka.
- Anjasari. (2016). Katekin Teh Indonesia : Prospek dan Manfaatnya. *Jurnal Kultivasi* 15 (2), 99-106.
- Anni, I. A., Saptiningsih, E., & Haryanti, S. (2013). Pengaruh Naungan Terhadap Pertumbuhan dan Produksi Tanaman Bawang Daun (*Allium fistulosum* L.) di Bandungan, Jawa Tengah. *Jurnal Biologi* 2(3), 31-400.
- Anonim. (2021). Understanding and Intepreting Lux Values. *Sensor API*, 1-107.
- AOAC. (2005). *Official Methods of Analysis. 18th edn.* Arlington, VA, USA: Association of Official Analytical Chemists.
- Apak, R., Güçlü, K., Demirata, B., Özyürek, M., Çelik, S., Bektaşoğlu, B., et al. (2007). Comparative Evaluation of Various Total Antioiidant Capacity Assays Applied to Phenolic Compounds with The CUPPRAC Assays. *Molecules*, 1496.
- Arumsari, K., Aminah, S., & Nurrahman. (2019). Aktivitas Antioksidan dan Sifat Sensoris Teh Celup Campuran Bunga Kecombrang, Daun Mint dan Daun Stevia. *Jurnal Pangan dan Gizi* 9(2), 79-93.

- Atmaja, M. P., Shabri, Harianto, S., Maulana, H., & Rohdiana, D. (2018). Karakteristik Fisik Tepung Teh Hijau yang Diproses Menggunakan Disc Mill dan Stone Mill. *J. Teknologi dan Industri Pangan*, 77-84.
- Atmaja, M., Shabri, Harianto, S., Maulana, H., & Rohdiana, D. (2018). Karakteristik Fisik Tepung Teh Hijau yang Diproses menggunakan Disc Mill dan Stone Mill. *J. Teknol. dan Industri Pangan vol.29(1)*, 77-84.
- Aventi. (2015). Penelitian Pengukuran Kadar Air Buah. *Seminar Nasional Cendekiawan*, 1-16.
- Azizah, D. N., & Faramayuda, F. (2014). Penetapan Kadar Flavonoid Metode AlCl<sub>3</sub> pada Ekstrak Metanol Kulit Buah Kakao (*Theobroma Cacao* L.). *Kartika Jurnal Ilmiah* 2(2), 24.
- Balasundram, N., Sundram, K., & Samman, S. (2006). Phenolic Compounds in Plants and Agri-industrial by Products : Antioxidant Activity, Occurrence, and Potential Uses. *Food Chemistry* 99 (1), 191-203.
- BPS. (2019). *Statistik Teh Indonesia* . Jakarta: Badan Pusat Statistik.
- Brighente, I., Dias, M., Verdi , L., & Pizzolati, M. (2007). Antioxidant Activity and Total Phenolic Content of Some Brazilian Spices. *Pharmaceutical Biology Vol.45(2)*, 156-161.
- Byers, T., & Guerrero, N. (1995). Epidemiologic Evidence for Vitamin C and Vitamin E in Cancer Prevention. *Am.J.Clin.Nutr.*62, 1385-1392.
- Cabrera, C., Artacho, R., & Gimenez, R. (2006). Beneficial Effects of Green Tea - A Review. *J. Am. Col. Nutr*, 79-99.
- Cahyani, D., & Rustanti, N. (2015). Pengaruh Penambahan Teh Hijau Terhadap Aktivitas Antioksidan dan Kadar Protein Minuman Fungsional Susu Kedelai dan Madu. *Journal of Nutrition College* 4(2), 394-399.
- Cheyner, V., Comte, G., Davies, K., Lattanzio, V., & Martens, S. (2013). Recent Advances on Their Biosynthesis, Genetics, and, Ecophysiology . *Plant Physiology and Biochemistry*, 1-20.
- Chu, D., & Juneja, L. (1997). *General Chemical Composition of Green Tea and Its Infusion. Chemistry and Applications of Green Tea*. New York: CRC New York.

- Chun, I. (2018, Juni 5). *How Does Shading Affect The Type and Quality of Tea*. Retrieved Agustus 17, 2021, from Yunomi: <https://yunomi.life/blogs/japanese-tea-guide/how-does-shading-affect-the-type-and-quality-of-tea>
- Chun, O., Kim, D., & Lee, C. (2003). Superoxide Radical Scavenging Activity of The Major Polyphenols in Fresh Plums. *J.Agric.Food.Chem* 51, 8067-8072.
- Clifford, A., & Cuppett, S. (2000). Review: Anthocyanins-nature, Occurrence and Dietary Burden. *J.Sci.Food.Agric.*80, 1063-1072.
- Cohen, S., & Kennedy, J. (2010). Plant Metabolism and The Environment : Implications for Managing Phenolics. *Crit Rev Food Sci* 50, 620-643.
- Colon, M., & Nerin, C. (2016). Synergistic, Antagonistic, Additive Interactions of Green Tea Polyphenols. *Eur Food Res Technol*, 211-220.
- Cominelli, E., Gusmaroli, G., Allegra, D., Galbiati, M., Wade, H. K., Jenkins, G. I., et al. (2008). Expression Analysis of Anthocyanin Regulatory Genes in Response to Different Light Qualities in *Arabidopsis thaliana*. *J. Plant Physiol*,165, 886-894.
- Cook, N., & Samman , S. (1996). Review : Flavonoids Chemistry, Metabolism, Cardioprotective Effects and Dietary Sources. *J.Nutri.Biochem* 7, 66-76.
- Dai, Y., Shen, Z., Liu, Y., Wang, L., Hannaway, D., & Lu. (2009). Effects of Shade Treatments on The Photosynthetic Capacity, Chlorophyll Fluorescence, and Chlorophyll Content of *Tetrastigma hemsleyanum* Diels et Gilg. *Environ. Exp. Bot*, 177-182.
- Dewi, K. (2008). Pengaruh Ekstrak Teh Hijau (*Camellia sinensis* var. *assamica*) Terhadap Penurunan Berat Badan. Kadar Trigliserida dan Kolesterol Total Pada Tikus Jantan Galur Wistar. *JKM* 7(2), 156-163.
- Dewi, P. P., Hidayat, R., & Permatasari, R. (2009). *Measurement of Antioxidant Capacity in the Commercial Tea and Correlation with the Total Phenol Content*. Bogor: Bogor Agricultural University.
- Fidrianny, I., Darmawati , A., & Sukrasno. (2014). Antioxydant Capacities from Different Polaities Extracts of Cucurbitaceae Leaves Using FRAP, DPPH Assay adn Correlation with Phenolic, Flavonoid, Carotenoid Content. *Int.J.Pharm.Sci* 6, 858-862.

- GFM. (2010, August). *Shading Corps*. Retrieved September 24, 2021, from <https://www.growingformarket.com/articles/Shading-Crops>
- Ghasemzadeh, A., Jaafar, H., & Rahmat, A. (2010). Antioxidant Activities, Total Phenolics and Flavonoids Content In Two Varieties of Malaysia Young Ginger (*Zingiber officinale roscoe*). *Mol.15*, 4324-4333.
- Graham, T. (1998). Flavonoid and Flavonol Glycoside Metabolism in *Arabidopsis*. *Plant Physiol.Biochem.36*, 135-144.
- Guenni, O., Seiter, S., & Figueroa, R. (2008). Growth Respons of Three *Brachiaria* Species to Light Intensity and Nitrogen Supply. *Trop.Grasslands*, 75-87 (42).
- Hajiaghaalipour, F., Sanusi, J., & Kanthimathi, M. (2016). Temperatue and Time of Sleeping Affect The Antioxidant Poperties of White, Green, and Black Tea Infusions. *J.Food Sci81*, 246-254.
- Hejinen, C., Haenen, G., Vanacker, F., Vjigh, W., & Bast, A. (2001). Flavonoids as Peroxynitrite Scavengers : The Role of The Hydroxyl Groups. *Toxi.In Vitro.15*, 3-6.
- Hidayat, T. (2008). *Karakteristik Fisik dan Organoleptik Tortilla Corn Chips dengan Penambahan Tepung Putih Telur sebagai Sumber Protein*. Bogor: Program Studi Teknologi Hasil Ternak. Fakultas Peternakan. Institut Pertanian Bogor.
- Hidayati, A. O., Lestariana, W., & Huriyati, E. (2012). Efek Ekstak Teh Hijau (*Camellia sinensis* (L.) O. Kuntze var. *assamica*) Terhadap Berat Badan dan Kadar Malondialdehid Wanita Overweight. *Jurnal Gizi Klinik Indonesia 9(1)*, 41-48.
- Hilal, Y., & Engelhardt, U. (2007). Characterisation of White Tea - Comparison to Green and Black Tea. *J. Consum. Prot. Food Saf* 2, 414-421.
- Hirai, M., Yoshikoshi, H., Kitano, M., Wakimizu, K., Sakaida, T., & Yoshioka. (2008). Production of Value-Added Crop of Green Tea in Summer. *International Workshop on Greenhouse Environmental Control and Crop Production in*, 411-417.
- Horie, H., Ema, K., & Sumikawa, O. (2017). Chemical Components of Matcha and Powdered Green Tea. *J. Cook.Sci 50*, 182-188.

- Islam, M., Kitamura, Y., Kokawa, M., & Fujii, S. (2020). Processing of Green Tea Pastes by Micro Wet Milling System : Influences on Physicochemical and Functional Properties. *Innivative Food Science and Emerging Technologies*, 2-12.
- Jaakola, L., Kärenlampi, S., Hohtola, A., & Määtä-Riihinen, K. (2004). Activation of Flavonoid Biosynthesis by Solar Radiation in Bilbery (*Vaccinium myrtillus* L.) leaves. *Planta* 218, 721-728.
- Jakubczyk, K., Kochman, J., Kwiatkowska, A., Kaldunska, J., Dec, K., Kawczuga, D., et al. (2020). Antioxidants Properties and Nutritional Composition of Matcha Green Tea. *Foods* 483, 2-10.
- Karori, S., Wachira, F., Wanyoko, J., & Ngure, R. (2007). Antioxidant Capacity of Diffeent Types of Tea Products. *African Journal of Biotechnology* 6, 2287-2296.
- Khaerunnisa, J., Purbaningtias, T., & Fauzi'ah, L. (2017). Determination of Total Phenolic with Different Tea Varieties by Spectrophotometric UV Visible. *The 2nd International Seminar on Chemical Education 2017*, (pp. 259-265). Indonesia.
- Khokkar, S., & Magnusdottir, S. (2002). Total Phenol, Catechin, and Caffeine Contents of Teas Commonly Consumed in The United Kingdom. *J Agr Food Chem*, 565-570.
- Ku, K., Choi, J., Kim, J., & Yoo , L. (2009). Metabolomics Analysis Reveals The Compositional Differences of Shade Grown Tea (*Camellia sinensis* L.). *J Agr Food Chem* 58, 418-426.
- Ku, K., Choi, J., Kim, J., & Yoo , L. (2009). Metabolomics Analysis Reveals The Compositional Differences of Shade Grown Tea (*Camellia sinensis* L.). *J Agr Food Chem* 58, 418-426.
- Landolt, W., Gunthardt-Georg, M., Pfenninger, I., Einig, W., Hampp, R., Maurer, S., et al. (1997). Effect of Fertilization on Ozone-induced Changes in The Metabolism of Birch (*Betula pendula*) Leaves. *New Phytol*, 389-397.
- Lattanzio, V., Cardinali, A., Ruta, C., Fortunato, I., Lattanzio, V., Linsalata, V., et al. (2009). Relationship of Secondary Metabolism to Growth in Oregano (*Origanum vulgare* L) Shoot Cultures Under Nutritional Stress. *Environt.Exp.Bot.*65, 54-62.

- Lee, K. I. (2003). Cocoa has More Phenolic Phytochemical and Higher Antioxdiant Capacity than Teas and Red Wine. *J.Agric.Food Chem* 51, 7292-7295.
- Lee, L.-S., Hea., C. J., Nari., S., Sang-Hee., K., Jong-Dae., P., Dae-Ja., J., et al. (2013). Metabolomic Analysis of the Effect of Shade Treatment on the Nutritional and Sensory Qualities of Green Tea. *Journal of Agricultural and Food Chemistry*, 332-338.
- Li, A., Li, S., Wu, X., Lu, H., Huang, M., Gu, R., et al. (2015). Influence of Light Intensity on the Yield and Quality of *Houttuynia cordata*. *Plant Production Science Vol 18, No 4*, 522-528.
- Lobos, G. A., Retamales, J. B., Hancock, J. F., Flore, J. A., Cobo, N., & Del Pozo, A. (2012). Spectral Irradiance, Gas Exchange Characteistics and Leaf Traits of *Vaccinium corymbosum* L. "Elliot" Grom Under Photo Selective Nets. *Environmental and Experimental Botany* 75, 142-149.
- Lorenzo, J., & Munekata, P. (2016). Phenolic Compounds of Green Tea : Health Benefits and Technological Application in Food. *Asia Pac J Trop Biomed Vol 6, No 8*, 709-719.
- Mahanta, P., & Baruah. (1992). *J.Sci Food Agric*, 59 : 21-26.
- Matthes, E., Douglas, A., Alan, D. P., & Cai, D. Y. (1997). Tea Chemistry. *Critical Reviews in Plant Sciences*, 415-480.
- Mishra, A., Tiwari, H., & Bhatt, R. (2010). Growth, Biomass Production and Photosynthesis of *Cenchrus ciliaris* L. Under *Acacia tortilis* (Forssk.) Hayne Based Silvopastoral Systems in Semi Arid Tropics. *Journal of Environmental Biology*, 987-993, 31(6).
- Mohotti, A. J., & Lawlor, D. W. (2002). Diurnal Variation of Photosynthesis and Photoinhibition in Tea: Effects of Irradiance and Nitrogen Supply During Growth in the Field. *J.Exp.Bot.*53, 313-322.
- Nadya, N. (2020, May 14). *Hari Matcha Nasional Mulai Tahun Ini Diperingati Setiap 14 Januari*. Retrieved Agustus 17, 2021, from Fimela: <https://www.fimela.com/lifestyle/read/4155725/hari-matcha-nasional-mulai-tahun-ini-diperingati-setiap-14-januari>
- Nazliniwaty, Hanum, T. I., & Laila, L. (2018). Antioxodant Activity of Green Tea (*Camellia sinensis* L. Kuntze) Ethanolic Extract using DPPH Method.

*International Conference of Science, Thenology, Engineering, Environmental, and Ramification Researches, 752-754.*

- Nishida, K. (2019, December 24). *Japanese Greentea Co.* Retrieved Agustus 15, 2021, from Japanese Green Tea and Health Blog: <https://www.japanesegreenteain.com/blogs/green-tea-and-health/everything-you-need-to-know-about-covering-process-for-matcha-and-gyokuro>
- Nishitani, E., & Sagesaka, Y. (2004). Simultaneous Determination of Catechins, Caffeine, and Other Phenolic Compounds in Tea Using New HPLC Method. *J. Food Compos.Anal* 17, 675-685.
- Nur, S., Sami, F., Wilda, R., Awaluddin, A., & Afsari, M. (2019). Korelasi antara Kadarr Total Flavonoid dan Fenolik dari Ekstrak dan Fraksi Daun Jati Putih (*Gmelina arborea* Roxb.) Terhadap Aktivitas Antioksidan. *Jurnal Farmasi Galenika : Galenika Hournal of Pharmacy* 5(1), 33-42.
- Osmond, C., Austin, M., Berry, J., Billings, W., Boyer, J., & Dacey, J. (1987). Stress Physiology and The Distributions of Plants. *BioScience* 37, 38-48.
- Pallas, J. E., Michel, B. E., & Harris, D. G. (1967). Photosynthesis, Transpiration, Leaf Temperature, and Stomatal Activity of Cotton Plants under Varying Wate Potentials. *Plant Physiol* 42, 76-88.
- Pastoriza, S., Mesias, M., Cabrera, C., & Rufian-Henares, J. (2017). Healthy Properties of Green and White Teas : An Update. *Food and Function* 8, 2650-2662.
- Pertiwi, F., & Made, N. (2018, Februari 4). *Apa Itu Matcha.* Retrieved Agustus 16, 2021, from Kompas: <https://travel.kompas.com/read/2018/02/04/120700027/apa-itu-matcha-?page=all>
- Pieterse, C., Leon-Reyes, A., Ent, V., & Wees, S. V. (2009). Networking by Small-molecule Hormones in Plant Immunity. *Nat Chem Biol* 5, 308-316.
- Pourmourad, F., Hosseinimehr, S., & Shahabimajd, N. (2006). Antioxodant Activity, Phenol and Flavonol and Flavonoid Contents of Some Selected Iranian Medicinal Plants. *African Journal of Biotechnology* 5(11), 1142-1145.



- PPTK. (2015, Mei 26). *Klon GB 1-11*. Retrieved Januari 10, 2022, from Pusat Penelitian Teh dan Kina: <https://www.gamboeng.com/pages/detail/2015/59/146>
- Prakash, A. (2001). Antioxidant Activity. *Analytical Progress*, 1-4.
- Probosari, E. (2019). Pengaruh Protein Diet terhadap Indeks Glikemik. *Journal of Nutrition and Health* 7 (1), 33-39.
- Pulitbang, P., & Balitbang, P. (2013). Teh Merah (*Camellia sinensis*) Hasil Eksplorasi di Kabupaten Wonosobo. *Warta Penelitian dan Pengembangan Tanaman Industri Vol.19, No.1*, 2-4.
- Puspitasari, M. (2019, September 19). *Perbedaan Teh Varietas Assamica dan Sinensis*. Retrieved Agustus 15, 2021, from Balai Penelitian Tanaman Industri dan Penyegar : Puslitbang Perkebunan - Badan Litbang Pertanian - Kementrian Pertanian: <http://balittri.litbang.pertanian.go.id/index.php/berita/info-teknologi/1024-perbedaan-teh-varietas-assamica-dan-sinensis>
- Rahayu, M. P., & Inanda, L. V. (2015). Penetapan Kadar Fenol Total Ekstrak Etil Asetat dan Fraksi Dichloromethan-etil Asetat Kulit Batang Mundu (*Garcinia dulcis*.Kurz). *Biomedika*, 38-44.
- Rahayu, R. (2016). *Penambahan Jenis dan Konsentrasi Bahan Pengisi Terhadap Formulasi Produk Banana Flakes*. Bandung: Program Studi Teknologi Pangan. Fakultas Teknik. Universitas Pasundan.
- Ramadan, P. (2015). *Knowing Antioxidants*. Yogyakarta: Graha Science.
- Rochana, A., Indriani, N. P., Ayuningsih, B., Hernaman, I., Dhalika, T., Rahmat, D., et al. (2016). Feed Forage and Nutrition Value at Altitude During the Dry Season in West Java. *Animal Production*, 85-93, 18(2).
- Rohdiana, D. (2015). Teh : Proses, Karakteristik, Komponen, dan Fungsionalnya. *FOODREVIEW INDONESIA VOL X/ NO.8*, 34-39.
- Roshanak, S., Rahimmalek, M., & Goli, S. (2016). Evaluation of Seven Different Drying Treatments in Respect to Total Flavonoid, Phenolic, Vitamin C Content, Chlorophyll, Antioxidant activity and Color of Green Tea (*Camellia sinensis* or *C. assamica*) Leaves. *J Food Sci Technol* 53(1), 721-729.



- Rusak, G., Komes, D., Likic', S., Horz'ic, D., & Kovac, M. (2008). Phenolic Content and Antioxidative Capacity of Green and White Tea Extracts Depending on Extraction Conditions and The Solvent Used. *Food Chemistry* 110(4), 852-858.
- Saijo, R. (1980). Effect of Shade Treatment on Biosynthesis of Catechins in Tea Plant. *Plant & Cell Physiology*, 989-998.
- Saijo, R. (2010). Effect of Shade Treatment on Biosynthesis of Catechins in Tea Plants. *Plant Cell Physiol.*58, 418-426.
- Sano, T., Horie, H., Matsunaga, A., & Hirono, Y. (2018). Effect of Shading Intensity on Morphological and Color Traits, and Chemical Components of New Tea (*Camellia sinensis* L.) shoots under direct covering cultivation. *J Sci Food Agric*.
- Saptari, T., Ningsih, T., Lohita, B., & Sayyidah, I. (2019). Kadar Fenolik dan Aktivitas Antioksidan Ekstrak Etanol Rumput Laut Coklat (*Padina australis*). *Fitofarmaka Vol.9(1)*, 1-8.
- Scharbert, S., Holzmann, N., & Hofmann, T. (2004). Identification of the Astringent Identification of The Astringent Human Bioresponse. *J.Agric.Food Chem.*52, 3498-3508.
- Schröder, L., Marahrens, P., Koch, J. G., Heidegger, H., Vilsmeier, T., Phan-Brehm, T., et al. (2019). Effects of Green Tea, Matcha Tea and Their Components Epigallocatechin gallate and quercetin on MCF-7 and MDA-MB-231 Breast Carcinoma Cells. *Oncol.Rep* 41, 387-396.
- Setyamidjaya, D. (2000). *Teh. Budi Daya dan Pengolahan Pascapanen*. Yogyakarta: Kanisius.
- Shahwar, D., Rehman, S., Ahmad, N., Ullah, S., & Raza, M. (2010). Antioxidant Activities of The Selected Plants from The Family Euphorbiaceae, Lauraceae, Malvaceae and Balsaminaceae. *African Journal of Biotechnology* 9, 1086-1096.
- Sivaci, A., & Duman, S. (2014). Evaluation of Seasonal Antioxidant Activity and Total Phenolic Compounds in Stems and Leaves of Some Almond (*Prunus amygdalus* L.) Varieties. *Biological Research*, 47-48.
- Sudaryat, Y., Kusmiyati, M., Pelangi, C. R., Rustamsyah, A., & Rohdiana, D. (2015). Antioxidant Activity of Ten Grades of Indonesia Black Tea

- (*Camellia sinensis* (L.)O.Kuntze) Liquor. *Research Journal of Tea and Quinine*, 95-100.
- Sugiyono. (2010). *Metode Penelitian Pendidikan Pendekatan Kuantitatif, Kualitatif, dan R&D*. Bandung: Alfabeta.
- Sulaiman, C., Sadashiva, C., George, S., Goplakrishnan, V., & Balachandran, I. (2013). Chromatographic Studies and in vitro Screening for Acetyl Cholinesterase Inhibition and Antioxidant Activity of three Acacia Species from South India. *Anal Chem Lett* 3 (2), 111-118.
- Syah, A. (2006). *Taklukkan Penyakit dengan Teh Hijau. Cet 1*. Jakarta: Penebar Swadaya.
- Taiz, L., & Zeiger. (2002). *Plant Physiology, 3rd ed*. Sunderland: Sinauer Associates, Inc.
- Tan, J., Engelhardt, U., Lin, Z., Kaiser, N., & Maiwald, B. (2017). Flavonoids, Phenolic Acids, Alkaloids, and Theanine in Different Types of Authentic Chinese White Tea Samples. *J. Food Compos* 57, 8-15.
- Teranishi, R., & Hornstein, I. (1995). Teranishi dan Hornstein International Vol 11, No. 3. *Marcel Decker*.
- Topuz, A., Dincer, C., Torun, M., Nadeem, H., Haznedar, A., & Ozdemir, F. (2014). Physicochemical Properties of Turkish Green Tea Powder : Effects of Shooting Period, Shading, and Clone. *Turk J Agric For* 38, 233-241.
- Unno, K., Furushima, D., Hamamoto, S., Iguchi, K., Yamada, H., Morita, A., et al. (2018). Stress-Reducing Function of Matcha Green Tea in Animal Experiments and Clinical Trials. *Nutrients* 10, 1468.
- USDA. (2014, - -). *Plants Database*. Retrieved January 10, 2021, from United States Departement of Agricuture : Natural Resources Conservation Service: <https://plants.usda.gov/core/profile?symbol=CASI16>
- Vison, J., Dabbagh, Y., Serry , M., & Jang, J. (1995). Plant Flavonoids, Especially Tea Flavonols, are Powerful Using An In Vitro Oxidation Model for Heart Disease. *J. Agric Food Chem* 43, 727-731.
- Wang, L., Micheletti, D., Palmer, J., Volz, R., & Lozano, L. (2011). High Temperature Reduces Apple Fruit Colour via Modulation of The Anthocyanin Regulatory Complex. *Plant Cell Environ* 34, 1176-1190.

- Wang, Y. S., Gao, L. P., Shan, Y., Liu, Y. J., Tian, Y. W., & Xia, T. (2012). Influence of Shade on Flavonoid Biosynthesis in Tea (*Camellia sinensis* (L.) O.Kuntze. *Scientia Horticulturae*, 8-16.
- Wei, K., Wang, L., Zhou, J., He, W., Zeng, J., Jiang, Y., et al. (2011). Catechin Content in Tea (*Camellia sinensis*) as Affected by Cultivar and Environment and Their Relation to Chlorophyll Contents. *Food Chemistry* 125, 44-48.
- Winarno, F. (1997). *Kimia Pangan dan Gizi*. Jakarta: Gramedia Pustaka Utama.
- Winarno, F. (2004). *Kimia Pangan dan Gizi*. Jakarta: Gramedia Pustaka Utama.
- Xiang, P., Zhu, Q., Tukhvatshin, M., Cheng, B., Tan, M., Liu, J., et al. (2021). Light Control of Cathecin Accumulation is Mediated by Photosynthetic Capacity in Tea Plant (*Camellia sinensis*). *BMC Plant Biol*, 21 :478.
- Xie, B., & Wang, H. (2006). Effects of Light Spectrum and Photoperiod on Contents of Flavonoid and Terpene in Leaves of *Ginkgo biloba* L. *J.Nanji.Fores.Uni*.30, 51-54.
- Young, D. R., & Smith, W. K. (1980). Influence of Sunlight on Photosynthesis, Water Relations, and Leaf Structure in the Understory Species *Arnica Cordifolia*. *Wiley on Behalf of the Ecological Society of America* 61(6), 1380-1390.
- Zaprometov, M. (1987). *Soviet Plant Physiol*, 34 : 561-572.
- Zhang, Q., Shi , Y., Ma, L., Yi, X., & Ruan, J. (2014). Metabolomic Analysis Using Ultra-Performance Liquid Chromatography-Quadrupole-Time of Flight Mass Spectrometry (UPLC-Q-TOF MS) Uncovers the Effects of Light Intensity and Temperature under Shading Treatments on the Metabolites in Tea. *PLOS One*, 1-10.
- Zhang, Z., Song, H., Peng, Z., Luo, Q., Ming, J., & Zhao, G. (2012). Characterization of Stipe and Cap Powders. *J Food Eng* 109, 406–413.
- Zhao, C., Li, C., Liu, S., & Yang, L. (2014). The Galloyl Cathecins Contributing to Main Antioxidant Capacity of Tea Made from *Camellia sinensis* in China. *The Scientific World Journal*, 1-11.