



DAFTAR PUSTAKA

- Al Hakimi, A. N. 2008. Effect of salicylic acid on biochemical changes in wheat plants under khat leaves residue. *Plant, Soil and Environment*, 54(7):288-293. DOI:[10.17221/418-PSE](https://doi.org/10.17221/418-PSE)
- Almansouri, M. J.M. Kinet, and S. Lutts. 2001. Effect of Salt and Osmotic Stresses on Germination in Durum Wheat (*Triticum durum* Desf.). *Plant and Soil*, 231(2): 243- 254. doi: [10.1023/A:1010378409663](https://doi.org/10.1023/A:1010378409663)
- Amador, B.M, E.T. Die'guez, J.L.G. Hernández, R.L. Aguilar, N.Y.A. Serrano, S.Z. Salgado, E.O.R. Puente, and C. Kaya. 2006. Effect of NaCl salinity in the genotypic variation of cowpea (*Vigna unguiculata*) during early vegetative growth. *Scientia Horticulturae* 108:423-431.
- An, C., and Mou, Z. 2011. Salicylic acid and its function in plant immunity. *J. Integ. Plant Biol.* 53, 412–428. doi: 10.1111/j.1744-7909.2011.01043.x
- Ashraf, M. 1989. The effect of NaCl on water relations, chlorophyll, and protein and proline content of two cultivars of blackgram (*Vigna mungo* L.). *Plant and soil* 119(2):205- 210.
- Augustin JM, Kuzina V, Andersen SB, Bak S. 2011. Molecular activities, biosynthesis and evolution of triterpenoid saponins. *Phytochemistry*. 72(6):435-57.
- Azam- Ali, S., E. Bonkoungou, C. Bowe, C. deKock, A. Godara, J. T. Williams. 2006. *Ber and Other Jujubes*. Southampton : International Centre for Underutilised Crops, p 2-3
- Azarmi, R., R.D. Taleshmikail, A. Gikloo. 2010. Effects of salinity on morphological and physiological changes and yield of tomato in hydroponic system. *J. Food. Agric. Environ.* 8: 573-576.
- Azooz, M.M., 2009. Salt stress mitigation by seed priming with salicylic acid in two faba bean genotypes differing in salt tolerance. *Int. J. Agric. Biol.*, 11: 343–350
- Barba-Espin G., M.J. Clemete-Moreno, S. Alvarez, M.F. Garcia-Legaz., J.A. Hernandez and P. Diaz- Vivancos. 2011. Salicylic acid negatively affects the response to salt stress in pea plants. *Plant Biol.*, 13 : 909- 917
- Barnett, N.M. and Naylor, A.W. 1966. Amino acid and protein metabolism in Bermuda grass during water stress. *Plant Physiol.* 41 : 1222- 1230
- Bissinger R, Modicano P, Alzoubi K, Honisch S, Faggio C, Abed M, Lang F. 2014. Effect of saponin on erythrocytes. *International Journal of*



Hematology. 100(1):51-59.

Blanch, G. P., M. C. Gomez-Jimenez, and M. L. R. del Castillo. 2020. Exogenous Salicylic Acid Improves Phenolic Content and Antioxidant Activity in Table Grapes. *Plant Foods for Human Nutrition*, 75(2):177-183.
[DOI:10.1007/s11130-019-00793-z](https://doi.org/10.1007/s11130-019-00793-z)

Chen, M. and Blankenship, R. E. 2011. Expanding the solar spectrum used by photosynthesis. *Trends in Plant Science*, 16: 427–431.

Dores, R G R., Sarah F G., Titiane V B., Maria C M F., Puliane M M., Thaiane C F. 2014. Phenolic Compounds, Flavonoids and Antioxidant Activity of Leaves, Flowers and Roots of White-Weed. *Hortic. Bras.* 3(4).

Efendi, M.Y. 2016. *Pengaruh Konsentrasi Asam Salisilat Terhadap Pertumbuhan Kacang Koro Pedang (Canavalia ensiformis L.) di Tanah Ultisol*. Universitas Pasir Pengaraian.

Einhellig, F.A. (1996) Interactions Involving Allelopathy in Cropping Systems. *Agronomy Jurnal*, 88: 886-893. doi:10.1007/s12185-014-1605

El- Housini E. A., M. A. Ahmed, M. S. Hassanein, dan M. M. Tawfik. 2014. Effect of Salicylic Acid (SA) on Growth and Quality of Stevia (Stevia rebaudiana Bert.) Under Salt Stress. *Agric. & Environ. Sci.*, 14(4): 275-281

El- tayeb, M.A. 2005. Response of barley grains to the interactive effect of salinity and salicylic acid. *Plant Growth*, 45 : 215-224

Elekofehinti OO. 2015. Saponins: Anti-diabetic principles from medicinal plants—A review. *Pathophysiology*. 22:95-103.
doi:10.1016/j.pathophys.2015.02.001

Encyclopaedia Britannica. 2022. Stomate, <https://britannica.com/science/stomate>, diakses pada 12/01/22

Friedman, M. 2006. Potato glycoalkaloids and metabolites: roles in the plant and in the diet. *J Agric Food Chem.* 54(23):8655-81.

Garg, N., Abdel-Aziz, S.M., & Aeron, A., 2016, Microbes in Food and Health, Springer, Switzerland 42-45.

Ginting, F.P., Y. Asbur, Y. Purwaningrum, M.S. Rahayu, Nurhayati. 2019. Pengaruh Stress Salinitas Terhadap Pertumbuhan, Komposisi Mineral, Kadar Prolin, Zat Antioksidan Kedelai. *Agriland*, 7(1): 48-54



Ginzberg I, Tokuhisa JG, Veilleux RE. 2009. Potato steroidal glycoalkaloids: biosynthesis and genetic manipulation. *Potato Res.* 52:1–15.

Güçlü-Ustündağ O, Mazza G. 2007. Saponins: properties, applications and processing. *Critical Reviews in Food Science and Nutrition.* 47(3):231-258. doi:10.1080/10408390600698197

Hadizadeh I, Peivastegan B, and Kolahi M. 2009. Antrifungal activity of nettle (*Urtica dioica L.*), colocynth (*Citrullus colocynthes L. Schrad.*), oleasnde (*Nerium oleander L.*) and konar (*Ziziphus spina-christi L.*) extracs on plant pathogenic fungi. *Pakistan journal of biological sciences*,12 (1) : 58-63.

Hakeem, K.R., F. Khan, R. Chandna, T.O. Siddiqui, and M. Iqbal. 2012. Genotypic variability among soybean genotypes under NaCl stress and proteome analysis of salt-tolerant genotype. *Appl. Biochem. Biotechnol.* 168:2309-2329.

Haryanti, S. 2010. Jumlah dan Distribusi pada Daun Beberapa Species Tanaman Dikotil dan Monokotil. *Buletin Anatomi dan Fisiologi*, 18 (2): 21- 28.

Hasanuzzaman, M., K. Nahar, Md. M. Alam, R. Roychowdhury, and M. Fujita. 2013. Physiological, Biochemical, and Molecular Mechanisms of Heat Stress Tolerance in Plants. *Int J Mol Sci.* 14(5): 9643- 9684. doi: [10.3390/ijms14059643](https://doi.org/10.3390/ijms14059643)

Hayat, S. et al. 2012. Role of proline under changing environments. *Plant Signaling & Behavior*, 7(11): 1456–1466.

Herrmann, K. M., and Weaver, L. M. 1999. The shikimate pathway. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 50: 473–503. doi: [10.1146/annurev.arplant.50.1.473](https://doi.org/10.1146/annurev.arplant.50.1.473)

Hesami, S. Nabizadeh, E. Rahimi, A. Rokhzadi. 2012. Effects of Salicylic acid levels and irrigation interval on growth and yield of coriander (*Coriandrum sativum*) in field condition. *Environmental and Experimental Biology*, 10 : 113-116.

Horváth, E., Pál, M., Szalai, G., Páldi, E., and Janda, T. 2007. Exogenous 4-hydroxybenzoic acid and salicylic acid modulate the effect of short-term drought and freezing stress on wheat plants. *Biol. Plant.* 51, 480–487. doi: [10.1007/s10535-007-0101-1](https://doi.org/10.1007/s10535-007-0101-1)

Hostettmann and Marston. 1995. *Saponins*. Cambridge University Press : New York.p 1-3

Hussein, M. M., L. K. Balbaa, and M. S. Gaballah. 2007. Salicylic Acid and Salinity Effects on Growth of Maize Plants. *Journal of Agriculture and*



Biological Sciences, 3(4): 321- 328

Hussen MH, El-sayed ME, & Said AA. 2010. Antihyperglycemic, antihyperlipidemic and antioxidant effects of *Ziziphus spina-christi* L. and *Ziziphus jujuba* in alloxan rats. International Journal of Pharmacology, 2 (5) : 5. DOI: [10.3923/ijp.2006.563.570](https://doi.org/10.3923/ijp.2006.563.570)

Ilmiah, H. H., T. R. Nuringtyas., L. H. Nugroho. 2018. Accumulation of Potential PhotoProtective Compound Groups in Mangrove (*Sonneratia caseolaris* (L.) Engler.) Leaves. *Pharmacogn J.*, 10(3):576-580.

Itkin M, Heinig U, Tzfadia O, Bhide AJ, Shinde B, Cardenas PD, Bocobza SE, Unger T, Malitsky S, Finkers R, Tikunov Y, Bovy A, Chikate Y, Singh P, Rogachev I, Beekwilder J, Giri AP, Aharoni A. 2013. Biosynthesis of antinutritional alkaloids in solanaceous crops is mediated by clustered genes. *Science*. 341(6142):175-9.

Kadam, P. and C. Pravin. 2010. Effect of NaCl salinity on stomatal density and stomatal behaviour of *Crotallaria*. *Bianano Frontier* 3(2):300-303.

Kaldehoff, R., A. Kolling, J. Meyers, U. Karmann, G. Roppel, and G. Richter. 1995. The blue light responsive AthH2 gene of *Arabidopsis thaliana* is primarily expresses in expanding as well as in differentiating cells and encodes a putative channel protein of the plasmalemma. *The Plant Journal* 7:87-95.

Kawano, T., Furuichi, T., and Muto, S. 2004. Controlled salicylic acid levels and corresponding signaling mechanisms in plants. *Plant Biotech.* 21: 319–335. doi: 10.5511/plantbiotechnology.21.319

Khairuna. 2019. Diktat Fisiologi Tumbuhan. Universitas Islam Negeri Sumatera Utara

Khan, M.I.R, M. Asgher, and N.A. Khan. 2014. Alleviation of salt-induced photosynthesis and growth inhibition by salicylic acid involves glycinebetaine and ethylene in mungbean (*Vigna radiata* L.). *Plant Physiol Biochem*. 80:67-74. doi: 10.1016/j.plaphy.2014.03.026.

Khan, W., B Prithiviraj, DL Smith. 2003. Photosynthetic responses of corn and soybean to foliar application of salicylates. *J. Plant Physiol.* 160: 485 –492.

Khan,M.I.R, M. Fatma, T.S Per, N. A. Anjum, and Nafees. 2015. Salicylic acid-induced abiotic stress tolerance and underlying mechanisms in plants. *Front. Plant Sci.* 30 (6):462. doi: 10.3389/fpls.2015.00462

Kurnia, T.D. dan Suprihati. 2013. Proline sebagai Penanda Ketahanan Kekeringan dan Salinitas pada Gandum. In Purnomo, D., ed. Prosiding Seminar



Nasional. Surakarta, 2013.

Kurniawati, S. et al. 2014. Pola Akumulasi Prolin dan Poliamin Beberapa Akses Tanaman Terung pada Cekaman Kekeringan. *J. Agron. Indonesia*, 42(2): 136-141.

Kusumiyati, T., M. Onggo, dan F.A. Habibah. 2017. Pengaruh Konsentrasi Larutan Garam NaCl terhadap Pertumbuhan & Kualitas Bibit Lima Kultivar Asparagus. *J. Hort. Indonesia*, 27(1) : 79-86.

Lauchli, A. and S. Grattan. 2014. Plant abiotic stress. *Elsevier*, 313-329

Lian, B., Zhou, X., Miransari, M. and Smith, D.L. (2000) Effects of Salicylic Acid on the Development and Root Nodulation of Soybean Seedling. *Journal of Agronomy and Crop Science*, 185, 187
[192.<http://dx.doi.org/10.1046/j.1439-037x.2000.00419.x>](http://dx.doi.org/10.1046/j.1439-037x.2000.00419.x)

Liang, X., L. Zhang, S. K. Natarajan, and D. F. Becker. 2013. Proline Mechanisms of Stress Survival. *Antioxid Redox Signal*, 19(9): 998–1011.
[doi: \[10.1089/ars.2012.5074\]\(https://doi.org/10.1089/ars.2012.5074\)](https://doi.org/10.1089/ars.2012.5074)

Lin D.R., Hu L.J., You H., Sarkar D., Xing B.S., Shetty K. Initial screening studies on potential of high phenolic-linked plantclonal systems for nitrate removal in cold latitudes. *J. Soils Sediment.* 2010;10:923–932.
[doi: 10.1007/s11368-010-0214-6](https://doi.org/10.1007/s11368-010-0214-6)

Mariaflaya.2018. Structure of Stomatal complex with open and closed stoma with titles, <https://www.istockphoto.com/id/vektor/struktur-kompleks-stomatal-dengan-stoma-terbuka-dan-tertutup-dengan-judul-gm1028226046-275636135>, diakses pada 12/01/22

Marjoni, R., Afrinaldi, A.D. Novita. 2015. Kandungan Total Fenol Dan Aktivitas Antioksidan Ekstrak Air Daun Kersen (*Muntingia calabura L.*). *Jurnal Kedokteran Yarsi*, 23 (3) : 187-196

Marrelli M, Conforti F, Araniti F, Statti GA. 2016. Effects of saponins on lipid metabolism: A review of potential health benefits in the treatment of obesity. *Molecules*. 21:1404. do i:10.3390/molecules21101404

Marschner, H., A. Kylin, and P.J.C. Kuiper. 1981. Differences in salt tolerance of three sugar beet genotypes. *Physiol. Plant* 51:234-238.

Martin, L. 2019. What are the roles of chlorophyll a b?.
<https://sciencing.com/what-are-the-roles-of-chlorophyll-a-b-12526386.html>, diakses pada 29/ 04/ 20

Masitoh, D. 2018. Pengaruh Konsentrasi Dan Waktu Pemberian SA (Salicylic Acid) Terhadap Pertumbuhan Dan Produksi Kedelai (*Glycine max (L.)*



Merrill) Pada Kondisi Tercekan Garam (NaCl). Universitas Jember. Halaman 10.

Michel GC, Nasseem ID, & Ismail F. 2011. Antidiabetik activity and stability study of the formulated leaf extract of *Ziziphus spina-christi* with the influence of seasonal variation. *Journal of Ethnopharmacology*, 133 (1); 53-62.

Mindari, W. 2009. Kesuburan Tanah Dan Pertumbuhan Tanaman. Surabaya : UPN Veteran Jawa Timur. Hal 1

Misra, N. and A.K. Gupta. 2005. Effect of salt stress on proline metabolism in two high yielding genotypes of green gram. *Plant Science*, 169(2):331-339.

Miura, K. and Y. Tada. 2014. Regulation of water, salinity, and cold stress responses by salicylic acid. *Frontiers in Plant Science*, 5(4) : 1-12.

Mohammadkhani,N. 2018. Effects Of Salinity On Phenolic Compounds In Tolerant And Sensitive Grapes. *Agriculture & Forestry*, 64(2) : 73-86, DOI: 10.17707/AgriculForest.64.2.05

Moses T, Papadopoulou KK, Osbourn A. 2014. Metabolic and functional diversity of saponins, biosynthetic intermediates and semi-synthetic derivatives. *Crit Rev Biochem Mol Biol*.49(6):439–462. doi:10.3109/10409238.2014.953628

Mustafa, N. R., Kim, H. K, Choi, Y. H., Erkelens, C., Lefeber, A. W., Spijksma, G., et al. 2009. Biosynthesis of salicylic acid in fungus elicited *Catharanthus roseus* cells. *Phytochemistry* 70: 532–539. doi: 10.1016/j.phytochem.2009.01.009

Nazar, R., S. Umar, N. A. Khan, O. Sareer. 2014. Salicylic acid supplementation improves photosynthesis and growth in mustard through changes in proline accumulation and ethylene formation under drought stress. *South African Journal of Botany*. 98: 84- 94.

Nugrahwati, Fauziah. 2016. *Uji Aktivitas Antipiretik Ekstrak Daun Bidara (Ziziphus spina-christi L.) terhadap Mencit Jantan (Mus Musculus)*. Undergraduate (S1) thesis, Universitas Islam Negeri Alauddin Makassar.

Nurmalasari, I. R. 2018. Kandungan Asam Amino Prolin Dua Varietas Padi Hitam Pada Kondisi Cekaman Kekeringan. *Gontor AGROTECH Science Journal*, 4 (1); 29- 43.

Ogihara, Y., Inoue, O., Otsuka, H., Kawai, K. I., Tanimura, T. and Shibata, S. (1976) Droplet counter current chromatography for the separation of plant products. *Journal of Chromatography*, 128(1): 218-223.



Orwa C, A Mutua, Kindt R , Jamnadass R, S Anthony. 2009. Agroforestry Database:a tree reference and selection guide version 4.0 <http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>, diakses pada 15/03/20

Palma, F. M. Lopez, N. A. Tejera, and C. Llunch. 2013. Salicylic acid improves the salinity tolerance of *Medicago sativa* in symbiosis with *Sinorhizobium meliloti* by preventing nitrogen fixation inhibition. *Plant Science*. 208C: 75- 82. doi : [10.1016/j.plantsci.2013.03.015](https://doi.org/10.1016/j.plantsci.2013.03.015)

Parashar, A. and S.K. Verma. 1993. Effect of gibberellic acid on chemical composition of wheat grown under different salinity levels. Paper presented at international conference on the Plant Physiology. Narendra Deva university og Agriculture and Technology (NDUAT) Kumarganj. Faizabad, India, 22-25.

Prihastati,E. 2012. Peranan Dan Pola Akumulasi Proline Tanaman Pada Adaptasi Cekaman Kekeringan. Seminar Nasional IX Pendidikan Biologi FKIP UNS.

Rao, K.V.M, A. S. Raghvendra. And K. and J.Reddy. 2006. *Physiology and Molecular Biologi of Stress Tolerance in Plants*. Dordrecht: Springer, p 2

Razmjoo, K., P. Heydarizadeh, M. R. Sabzalian. 2008. Effect of salinity and drought stresses on growth parameters and essential oil content of *Matricaria chamomila*. *International Journal of Agriculture and Biology*,10(4):1560-8530

Reece, J.B., L.A. Urry, M.L. Cain, S. A. Wasserman, P.V. Minorsky, R.B. Jackson. 2014. *Biologi*. Jakarta: Penerbit Erlangga

Rina, Y dan H. Syahbuddin. 2013. Zona Kesesuaian Lahan Rawa Pasang Surut Berbasis Keunggulan Kompetitif Komoditas. *Sepa*, 10(1): 103-117.

Rosmayati, N. Rahmawati, R. P. Astari, dan F. Wibowo. 2015. Analisa Pertumbuhan Vegetatif Kedelai Hibridisasi Genotipa Tahan Salin dengan Varietas Anjasmoro untuk Mendukung Perluasan Areal Tanam Di Lahan Salin. *Pertanian Tropik*, 2(2): 132-139.

Sakr, M. T., H. M. A. El- Salam, M. I. Atta, dan M. A. A. El- Aal. 2013. Alleviating The Harmful Effect of Salinity Stress on Soybean Plants by Using Some Promoters. *Plant Production*, 4(2): 205- 218

Sánchez, N. F., R. S. Coronado., B. H. Carlos & C. V. Canongo. 2019. Shikimic Acid Pathway in Biosynthesis of Phenolic

Setyorini dan Yusnawan. 2016. Peningkatan Kandungan Metabolit Sekunder Aneka Kacang sebagai Respon Cekaman Biotik. *Iptek Tanaman Pangan*,



11(2): 167- 174

- Shakirova, F.M., A. R. Sakhabutdinova, M. V. Bezrukova, R. A. Fatkhutdinova, dan D. R. Fatkhutdinova. 2003. Changes in the Hormonal Status of Wheat Seedlings Induced by Salicylic Acid and Salinity. *Plant Science*, 164: 317-322.
- Sharma, A., B. Shahzad., A. Rehman., R. Bhardwaj., M. Landi., B. Zheng. 2019. Response of Phenylpropanoid Pathway and the Role of Polyphenols in Plants under Abiotic Stress. *Molecules*, 24: 2452.
- Singh, A.K. and R.S. Dubey. 1995. Changes in chlorophyll a and b contents and activities of photosystem 1 and 2 in rice seedlings induced by NaCl. *Photosynthetica*, 31:489-499.
- Singh, P.K., dan S. Gautama. 2013. Role of Salicylic acid Treatment on Grain yield, Process of grain Growth, and Some of Chemical and Morphological Traits of Chamran Cultivar Wheat (*Triticum aestivum*). *Advances in Environmental Biology*, 7 (11): 3234- 3240
- Sivasankari, M.P. and A. Sankaravadivoo . 2017. Leaf Anatomical Studies of *Ziziphus mauritiana* Lam. *Int. J. Curr. Res. Biosci. Plant Biol.* 4(8), 73-79 . doi: <https://doi.org/10.20546/ijcrbp.2017.408.010>
- Sobir, Miftahudin dan S. Helmi. 2018. Respon Morfologi dan Fisiologi Genotipe Terung (*Solanum melongena* L.) terhadap Cekaman Salinitas. *J. Hort. Indonesia*, 9(2) : 131- 138.
- Souri and Tohidloo. 2019. Effectiveness of different methods of salicylic acid application on growth characteristics of tomato seedlings under salinity. *Chem. Biol. Technol. Agric.*, 6(26):1-7. <https://doi.org/10.1186/s40538-019-0169>
- Stepien, P. and G. Klobus. 2006. Water relations and photosynthesis in *Cucumis sativus* L. leaves under salt stress. *Biologia Plantarum* 50(4):610-616.
- Summart, J., P. Thanonkeo, S. Pannichajakul, P. Prapthepha, and MT. McManus. 2010. Effect of salt stress on growth, inorganic ion and proline accumulation in thai aromatic rice Khao Dawk Mali 105. *Callus Cultura Afr. J. Biotechnol.* 9(2):145-152.
- Swamy,S. 2022. Stomata : Definition, Diagrams, and Functions, <https://embibe.com/exams/stomata/>, diakses pada 13/01/22
- Tarigan, D. F. dan F. K. Wardana. 2020. Pertumbuhan Tanaman Akar Wangi (*Vetiveria zizanioides* L.) di Tanah Salin Dengan Perlakuan Asam Salisilat Dan Fungi Mikoriza Arbuskular. *Agrium*, 22 (3): 166-171.



The Editors of Encyclopaedia Britannica. 2020. Chlorophyll.

<https://www.britannica.com/science/chlorophyll>, diakses pada 29/04/20

Townsend, C.C., Guest, E., 1980. *Flora of Iraq*. Baghdad. Vol.4, Part One. pp.432-437.

Tripama,B. dan H. Hidayatullah. 2017. Respon Pertumbuhan dan Kandungan Fenol pada Tanaman Cengkeh (*Eugenia caryophyllus*) Terhadap Macam Media dan Cekaman NaCl. *Agritrop*, 15 (1) : 119- 130.

Tufaila, M., S. Alam, dan S. Leomo. 2014. Strategi Pengelolaan Tanah Marginal: Ikhtiar Mewujudkan Pertanian yang Berkelanjutan. Kendari: Unhalu Press.

USDA.2020.Classification Of *Ziziphus mauritina*,
<https://plants.usda.gov/java/ClassificationServlet?source=display&classid=ZIMA>, diakses pada 17/03/20

Vattakaven T, George R, Balasubramanian D, Réjou-Méchain M, Muthusankar G, Ramesh B, Prabhakar R (2016) India Biodiversity Portal: An integrated, interactive and participatory biodiversity informatics platform. *Biodiversity Data Journal* 4: e10279. <https://doi.org/10.3897/BDJ.4.e10279>"

Vattem D.A., Randhir R., Shetty K. Cranberry phenolics-mediated antioxidant enzyme response in oxidatively stressed porcine muscle. *Process Biochem.* 2005;40:2225–2238. doi: 10.1016/j.procbio.2004.09.001.

Velderrain-Rodríguez GR, Palafox-Carlos H, Wall-Medrano A, Ayala-Zavala JF, Chen CY, Robles-Sánchez M, Astiazaran-García H, Alvarez-Parrilla E, González-Aguilar GA. 2014. *Food Funct*, 5(2):189-97.

Verberne, M. C., Budi Muljono, R. A., and Verpoorte, R. 1999. "Salicylic acid biosynthesis," in *Biochemistry and Molecular Biology of Plant Hormones*, eds P. P. J. Hooykaas, M. A. Hall, and K. R. Libbenga. Amsterdam: Elsevier Science BV: 295–314. doi: 10.1016/S0167-7306(08)60493-7

Vincken JP, Heng L, De Groot A, Gruppen H. Saponins, classification and occurrence in the plant kingdom. *Phytochemistry*. 2007;68:275–97

Wahyuni, D. K., W. Ekasari, J. R. Witono, dan H. Purnobasuki. 2016. *Toga Indonesia*. Surabaya : Airlangga University Press, halaman 1

Wahyuni, R.D. 2014. Pengaruh Paklobutrazol Terhadap Pertumbuhan dan Kandungan Saponin Tanaman Binahong (*Anredera cordifolia* (Ten) Steenis). Yogyakarta : Universitas Gadjah Mada



- Weisany, W., Y. Sohrabi, Y. Heidari, A. Siosemardeh, and K. Ghassemi. 2011. Physiological responses of soybean (*Glycine max L.*) to zinc application under salinity. *Australian Journal of Crop Science*, 5(11):1441-1447
- Wijayanti, W., Taryono, Toekidjo. 2014. Keragaan 29 galur kacang tanah (*Arachis hypogaea L.*) pada kondisi salin. *Vegetalika*. 3: 40-51.
- Yassin, M., A. El Sabagh, M. S. Islam, and A. Hossain. 2019. Comparative performance of two bread wheat (*Triticum Aestivum L.*) genotypes under salinity stress. *Ecology and Environmental Research*, 17(2):5029 - 5041. DOI:[10.15666/aeer/1702_50295041](https://doi.org/10.15666/aeer/1702_50295041)
- Yuliani., F. Rachmadiarti., Dewi, S. K., & Asri, M. T. 2019. Total Phenolic And Flavonoid Contents of *Elephantopus scaber* and *Ageratum conyzoides* (Asteraceae) Leaves Extracts from Various Altitude Habitats. *Eco. Env. & Cons*, 25: S106-S11.
- Yuniati R. 2004. Penapisan Galur Kedelai *Glycine max* (L.) Merrill Toleran Terhadap NaCl Untuk Penanaman di Lahan Salin. *Makara Sains*. 8(1):21– 24.
- Zahra,S., B. Ami, V.S.M. Ali, Y. Ali, and Y. Mehdi. 2010. The Salicylic acid effect on the tomato (*Lycopersicum esculentum Mill.*) sugar, protein and proline contents under salinity stress (NaCl). *Journal of Biophysics and structural Biology*, 2(3): 35-41