

DAFTAR PUSTAKA

- Agatonovic-Kustrin, S., Morton, D.W., and Ristivojević, P., 2016. Assessment of antioxidant activity in Victorian marine algal extracts using high performance thin-layer chromatography and multivariate analysis. *J. Chromatogr. A* 1468, 228–235.
- Ah, W., N, D., 2014. Biolarvacidal activity of methanol extract of Kipahit leaves (*Tithonia diversifolia*) against larvae of *Chrysomya bezziana* fly. *Jurnal Ilmu Ternak Dan Veteriner*. 19.
- Al-Saeedi, A.H., Al- Ghafri, M.T.H., and Hossain, M.A., 2016. Comparative evaluation of total phenols, flavonoids content and antioxidant potential of leaf and fruit extracts of *Omani ziziphus jujuba* L. *Pac. Sci. Rev. Nat. Sci. Eng.*
- Amić, A., Marković, Z., Marković, J.M.D., Jeremić, S., Lučić, B., and Amić, D., 2016. Free radical scavenging and COX-2 inhibition by simple colon metabolites of polyphenols: A theoretical approach. *Comput. Biol. Chem.* 65, 45–53.
- Anagnostopoulou, M.A., Kefalas, P., Papageorgiou, V.P., Assimopoulou, A.N., and Boskou, D., 2006. Radical scavenging activity of various extracts and fractions of sweet orange peel (*Citrus sinensis*). *Food Chem.* 94, 19–25. doi:10.1016/j.foodchem.2004.09.047
- Athar, M. and Nasir, S.M., 2005. Taxonomic perspective of plant species yielding vegetable oils used in cosmetics and skin care products. *Afr.J.Biotechnol* 4: 36–44.
- Bharathi, E., and Jagadeesan, G., 2014. Antioxidant potential of hesperidin and ellagic acid on renal toxicity induced by mercuric chloride in rats. *Biomed. Prev. Nutr.* 4, 131–136.
- Bravo, K., Alzate, F., and Osorio, E., 2016. Fruits of selected wild and cultivated Andean plants as sources of potential compounds with antioxidant and anti-aging activity. *Ind. Crops Prod.* 85, 341–352.
- Chen, L.X., Hu, D.J., Lam, S.C., Ge, L., Wu, D., Zhao, J., Long, Z.R., Yang, W.J., Fan, B., and Li, S.P., 2016. Comparison of antioxidant activities of different parts from snow chrysanthemum (*Coreopsis tinctoria nutt.*) and identification of their natural antioxidants using high performance liquid chromatography coupled with diode array detection and mass spectrometry and 2,2'-azinobis(3-ethylbenzthiazoline-sulfonic acid)diammonium salt-based assay. *J. Chromatogr. A* 1428, 134–142.

- Creswell, C. J., Runquist, O. A., and Campbell, M. M., 1982. *Analisis Spektrum Senyawa Organik*. ITB, Bandung.
- Dungir, S.G., Katja, D.G., and Kamu, V.S., 2012. Aktivitas antioksidan ekstrak fenolik dari kulit buah manggis (*Garcinia mangostana* L.). *J. MIPA Unsrat Online* 1, 11–15.
- Erel, O., 2004. A novel automated direct measurement method for total antioxidant capacity using a new generation, more stable ABTS radical cation. *Clinical Biochemistry*, **37**: 277–285.
- Eryilmaz, S., Gül, M., İnkaya, E., İdil, Ö., and Özdemir, N., 2016. Synthesis, crystal structure analysis, spectral characterization, quantum chemical calculations, antioxidant and antimicrobial activity of 3-(4-chlorophenyl)-3a,4,7,7a-tetrahydro-4,7-methanobenzo[d]isoxazole. *J. Mol. Struct.* 1122, 219–233.
- Fakhlai, R., Khoo, H.E., Sia, W.C.M., and Yim, H.S., 2015. Extraction and evaluation of antioxidant potential in rambutan rind as food waste. *ResearchGate* 7, 35–44.
- Fawole, O.A., and Opara, U.L., 2016a. Stability of total phenolic concentration and antioxidant capacity of extracts from pomegranate co-products subjected to in vitro digestion. *BMC Complement. Altern. Med.* 16.
- Fidrianny, I., Sukowati, A., and Sukrasno, S., 2015. Invitro antioxidant activities of various leaves extracts from five varieties of rambutan (*Nephelium lappaceum* L.) and its correlation with total flavonoid, phenolic, carotenoid content. *Asian J. Pharm. Clin. Res.* 8, 139–143.
- Fidriany, I., Ruslan, K., and Saputra, J., 2013. Antioxidant Activities of Different Polarity Extracts from Cashew (*Anacardium occidentale* L.) Leaves and Isolation of Antioxidant Compound. *J. Med. Planta* 2.
- Floegel, A., Kim, D.-O., Chung, S.-J., Koo, S.I., and Chun, O.K., 2011. Comparison of ABTS/DPPH assays to measure antioxidant capacity in popular antioxidant-rich US foods. *J. Food Compost Anal.* , 24: 1043–1048.
- Gritter, R., Bobbit, J. M., Schwarting, A. E., 1991. *Pengantar Kromatografi*, Kedua. ed.
- Gunenc, A., HadiNezhad, M., Farah, I., Hashem, A., and Hosseinian, F., 2015. Impact of supercritical CO₂ and traditional solvent extraction systems on the extractability of alkylresorcinols, phenolic profile and their antioxidant activity in wheat bran. *J. Funct. Foods* 12, 109–119.

- Harbone, J. B., 1973. Metode Fitokimia; *Penuntun Cara Modern Menganalisis Tumbuhan*. ITB, Bandung.
- He, J.-M., Zhang, and S.-Y., and Mu, Q., 2016. Online-storage recycling counter-current chromatography for preparative isolation of naphthaquinones from *Arnebia euchroma* (Royle) Johnst. *J. Chromatogr. A* 1464, 79–86.
- Hendayana, S., Karohman, A., Sumarna, A. A., and Supriatna, A., 1994. *Kimia Analitik Instrumen*, kesatu. ed. IKIP Semarang Press, Semarang.
- Hoffman, E. D., Charette, J., and Stroobant, V., 1996. Mass Spectrometry. *John Wiley & Sons*, Unites Stated of America.
- Hofmann, T., Nebehaj, E., and Albert, L., 2016. Antioxidant properties and detailed polyphenol profiling of European hornbeam (*Carpinus betulus* L.) leaves by multiple antioxidant capacity assays and high-performance liquid chromatography/multistage electrospray mass spectrometry. *Ind. Crops Prod.* 87, 340–349.
- Islam, T., Yu, X., and Xu, B., 2016. Phenolic profiles, antioxidant capacities and metal chelating ability of edible mushrooms commonly consumed in China. *LWT - Food Sci. Technol.* 72, 423–431.
- Jabbar, A., Ali, A., Tawab, A., and Iqbal, M., 2014. Fatty Acid Profiling of Lipid A Isolated from Indigenous *Salmonella Typhi* Strain by Gas Chromatography Mass Spectrometry. *J. Chem. Soc. Pak.* 36.
- Jayaprakasha, G.K., Girenavar, B., and Patil, B.S., 2008. Radical scavenging activities of Rio Red grapefruits and Sour orange fruit extracts in different in vitro model systems. *Bioresour. Technol.* 99, 4484–4494.
- Jo atilde o, da R.L.N., Amanda, D. de A. uacute jo U. ocirc a, Priscila, A. de M., Clovis, M. ecirc do B.F., Juciara, C.G.T. oacute rio, Alexandre, G. da S., Rafael, M.X., M aacute rcia, V. da S., and Maria, T. dos S.C., 2016. Phytochemical screening, total phenolic content and antioxidant activity of some plants from Brazilian flora. *J. Med. Plants Res.* 10, 409–416.
- Kamaludin, N.H.I., Mun, L., and Sa'adi, R.A., 2016. Evaluation of antioxidant activity of some tropical fruit peel extracts: Extraction conditions optimization of rambutan peel extract. *ResearchGate* 11, 1623–1631.
- Karla Bigetti Guergoletto, Alessandra Yuri Tsuruda, Elisa Yoko Hirooka, Elvis Peboni Martins, Jean Clovis Bertuol de Souza, Kátia Sivieri, Salvador Massaguer Roig, and Sandra Garcia, 2012. Dried Probiotics for Use in Functional Food Applications. *INTECH Open Access Publisher*.

- Kikuzaki, H., Hisamoto, M., Hirose, K., Akiyama, K., and Taniguchi, H., 2002. Antioxidant properties of ferulic acid and its related compounds. *J.Agric Food Chem*, 50: 2161–2168.
- Kristatnti, R. E., 2012. 'Isolasi dan Elusidasi Struktur Senyawa Antioksidan dan Penghambat Xantin Oksidase dari Buah Andaliman (*Zanthoxylum acanthopodium* DC.)', , *Tesis*, . Universitas Indonesia, Jakarta.
- La, S., Sia, C.M., GAb, A., PNa, O., and Yim, H.S., 2013. The effect of extraction conditions on total phenolic content and free radical scavenging capacity of selected tropical fruits' peel. *Health (N. Y.)* 4, 80–102.
- Lim, T.K., 2013. *Nephelium lappaceum* L., in: Edible Medicinal And Non-Medicinal Plants. *Springer*, Dordrecht, pp. 62–71.
- Ling, L.T., Palanisamy, U.D., and Cheng, H.M., 2010. Prooxidant/Antioxidant Ratio (ProAntidex) as a Better Index of Net Free Radical Scavenging Potential. *Molecules* 15, 7884–7892.
- Liyana-Pathirana, C.M., and Shahidi, F., 2007a. Antioxidant and free radical scavenging activities of whole wheat and milling fractions. *Food Chem.* 101, 1151–1157.
- Liyana-Pathirana, C.M., and Shahidi, F., 2007b. The antioxidant potential of milling fractions from breadwheat and durum. *J. Cereal Sci.* 45, 238–247.
- Mahisanunt, B., Na Jom, K., Matsukawa, S., and Klinkesorn, U., 2016. Solvent fractionation of rambutan (*Nephelium lappaceum* L.) kernel fat for production of non-hydrogenated solid fat: Influence of time and solvent type. *J. King Saud Univ.*
- Mikusanti, E., and Hotdelina, S., 2012. Aktivitas Antioksidan dan Sifat Kestabilan Warna Campuran Ekstrak Etil Asetat Kulit Buah Manggis (*Garcinia mangostana* L.) dan Kayu Secang (*Caesalpinia sappan* L.). *JPS* 15.
- Mulja, M., and Suharma, 1995. Analisis Instrumental. Airlangga University Press, Surabaya. Perera, A., Appleton, D., Ying, L.H., Elendran, S., Palanisamy, U.D., 2012. Large scale purification of geraniin from *Nephelium lappaceum* L. rind waste using reverse-phase chromatography. *Sep. Purif. Technol.* 98, 145–149. <https://doi.org/10.1016/j.seppur.2012.06.019>

- Mohammed, Y.H., Ghaidaa, J.M., and Imad, H.H., 2016. Analysis of bioactive chemical compounds of *Nigella sativa* using gas chromatography-mass spectrometry. *J.Pharmacogn.Phytother.* 8,8–24. <https://doi.org/10.5897/JPP2015.0364>
- NIST Mass spectrometry Data Center, 2014. *Bis(2-ethylhexyl) phthalate* [WWW Document]. URL <http://webbook.nist.gov/cgi/cbook.cgi?ID=C117817&Mask=200> (accessed 7.10.17).
- Omena, C.M.B., Valentim, I.B., Guedes, G. da S., Rabelo, L.A., Mano, C.M., Bechara, E.J.H., Sawaya, A.C.H.F., Trevisan, M.T.S., da Costa, J.G., Ferreira, R.C.S., Sant'Ana, A.E.G., and Goulart, M.O.F., 2012. Antioxidant, anti-acetylcholinesterase and cytotoxic activities of ethanol extracts of peel, pulp and seeds of exotic Brazilian fruits. *Food Res. Int.* 49, 334–344.
- Palanisamy, U., Cheng, H.M., Masilamani, T., Subramaniam, T., Ling, L.T., and Radhakrishnan, A.K., 2008. Rind of the rambutan, *Nephelium lappaceum* L., a potential source of natural antioxidants. *Food Chem.* 109, 54–63.
- Perera, A., Appleton, D., Ying, L.H., Elendran, S., and Palanisamy, U.D., 2012. Large scale purification of geraniin from *Nephelium lappaceum* L.rind waste using reverse-phase chromatography. *Sep. Purif. Technol.* 98, 145–149.
- Permatasar, L., and Rohman, A., 2016. 2,2'-diphenil-1-picrylhydrazil (DPPH) Radical Scavenging Activity of Extracts and Fractions of Rambutan (*Nephelium lappaceum* L.) Peel. *Res. J. Phytochem.* 10, 75–80.
- Phongpaichit, S., Nikom, J., Rungjindamai, N., Sakayaroj, J., HutadilokTowatana, N., Rukachaisirikul, V., and Kirtikara, K. (2007). Biological activities of extracts from endophytic fungi isolated from garcinia plants. *FEMS I.Med. Microbiology*, **51**, 517-525.
- Priestman, T.J., 2008. Cancer chemotherapy in *clinical practice*. Springer, New York ; London.
- Ragae, S., Abdelaal, E., and Noaman, M., 2006. Antioxidant activity and nutrient composition of selected cereals for food use. *Food Chem.* 98, 32–38.
- Rahayu, S.E., Sri, R., Lestari, Wulandari, N., and Maslikah, S., 2015. Effect of Polyphenol from Rambutan Peel Extract on Serum Lipidand Protein Profileof Visceral Fat on Normal and Obesity Rat Model. *ResearchGate* 8, 974–4304.
- Rohman, A., and Riyanto, S., 2005. Daya antioksidan ekstrak etanol Daun Kemuning (*Murraya paniculata* (L) Jack) secara in vitro. *Maj. Farm. Indones.* 16, 136–140.

- Rohman, A., and Riyanto, S., Utari, D., 2006. Aktivitas antioksidan, kandungan fenolik total dan kandungan flavonoid total ekstrak etil asetat buah Mengkudu serta fraksi-fraksinya. *Maj. Farm. Indones.* 17, 136–142.
- Smith, R., 2003. Before the injection - Modern methods of sample preparation for separation techniques. doi:10.1016/S0021-9673(03)00511-9
- Safdar, M.N., Kausar, T., Jabbar, S., Mumtaz, A., Ahad, K., and Saddozai, A.A., 2016. Extraction and quantification of polyphenols from kinnow (*Citrus reticulata* L.) peel using ultrasound and maceration techniques. *J. Food Drug Anal.*
- Samuagam, L., Sia, C.M., Akowuah, G.A., Okechukwu, P.N., and Yim, H.S., 2015. In vivo antioxidant potentials of rambutan, mangosteen, and langsung peel extracts and effects on liver enzymes in experimental rats. *ResearchGate* 24.
- Sideney, B.O., Francini, Y.K., and Shaiana, P.M., 2015. Antioxidant activity, total phenolic and flavonoids contents in *Stachytarpheta cayennensis*, (Rich.) Vahl. (Verbenaceae). *J. Med. Plants Res.* 9, 569–575.
- Silva, K.D.R.R., and Sirasa, M.S.F., 2016. Antioxidant properties of selected fruit cultivars grown in Sri Lanka. *Food Chem.*
- Silva, P.J., and Prather, K.A., 2000. Interpretation of Mass Spectra from Organic Compounds in Aerosol Time-of-Flight Mass Spectrometry. *Anal. Chem.* 72, 3553–3562. <https://doi.org/10.1021/ac9910132>
- Siregar, M., 2006. Species Diversity of Local Fruit Trees in Kalimantan: Problems of Conservation and Its Development. *Biodiversitas J. Biol. Divers.* 7, 94–99.
- Stahl, E., 1969. Thin Layer Chromatography. *Springer*, New York.
- Soeng, S., 2015. Antioxidant and hypoglycemic activities of extract and fractions of Rambutan seeds (*Nephelium lappaceum* L.). *Biomed. Eng.* 1, 6.
- Sudjadi, 1988. Metode Pemisahan. *Kanisius*, Yogyakarta.
- Sudjadi, 1983. Penentuan Struktur Senyawa Organik. Ghalia Indonesia, Jakarta Timur.
- Sudjono, T.A., Hidayati, A.L., and Suhendi, A., others, 2014. Pengujian Daya Antioksidan dari Beberapa Ekstrak Kulit Buah Asli Indonesia dengan Metode FTC.
- Sun, L., Zhang, H., and Zhuang, Y., 2012. Preparation of free, soluble conjugate, and insoluble-bound phenolic compounds from peels of rambutan (*Nephelium lappaceum* L.) and evaluation of antioxidant activities in vitro. *J. Food Sci.* 77, C198–204.

- Thitilertdecha, N., Teerawutgulrag, A., Kilburn, J.D., and Rakariyatham, N., 2010. Identification of Major Phenolic Compounds from *Nephelium lappaceum L.* and Their Antioxidant Activities. *Molecules* 15, 1453–1465.
- Thitilertdecha, N., and Rakariyatham, N., 2011. Phenolic content and free radical scavenging activities in rambutan during fruit maturation. *Sci. Hortic.* 129, 247–252. doi:10.1016/j.scienta.2011.03.041
- Thitilertdecha, N., Teerawutgulrag, A., and Rakariyatham, N., 2008. Antioxidant and antibacterial activities of *Nephelium lappaceum L.* extracts. *LWT - Food Sci. Technol.* 41, 2029–2035.
- Tjandra, O., Rusliati R, T., and Zulhipri, Z., others, 2011. Uji Aktivitas Antioksidan dan Profil Fitokimia Kulit Rambutan Rapih (*Nephelium lappaceum L.*).
- Tristantini, D., Ismawati, A., Pradana, B.T., and Jonathan, J.G., 2016. Pengujian Aktivitas Antioksidan Menggunakan Metode DPPH pada Daun Tanjung (*Mimusops elengi L*), in: *Seminar Nasional Teknik Kimia Kejuangan*. p. G1.
- Van Der Werf, R., Dal, S., Le Grandois, J., Aoude-Werner, D., Digel, F., Ennahar, S., Sigrist, S., and Marchioni, E., 2015. Determination of active radical scavenging compounds in polar fruit and vegetable extracts by an on-line HPLC method. *LWT - Food Sci. Technol.* 62, 152–159.
- Vemula, H., Kitase, Y., Ayon, N.J., Bonewald, L., and Gutheil, W.G., 2017. Gaussian and linear deconvolution of LC-MS/MS chromatograms of the eight aminobutyric acid isomers. *Anal. Biochem.* 516, 75–85.
- Williams, D. H., and Fleming, I., 1997. *Spectroscopic Methods In Organic Chemistry*. McGraw-Hill Publishing Company, England.
- Wootton-Beard, P.C., Moran, A., and Ryan, L., 2011. Stability of the total antioxidant capacity and total polyphenol content of 23 commercially available vegetable juices before and after in vitro digestion measured by FRAP, DPPH, ABTS and Folin–Ciocalteu methods. *Food Research International*, 44: 217–224.
- Yu, E.A., Kim, G.-S., Jeong, S.W., Park, S., Lee, S.J., Kim, J.H., Lee, W.S., Bark, K.-M., Jin, J.S., and Shin, S.C., 2014. Flavonoid profile and biological activity of Korean citrus varieties (II): Pyunkyul (*Citrus tangerina Hort. ex Tanaka*) and overall contribution of its flavonoids to antioxidant effect. *J. Funct. Foods* 6, 637–642.

- Yuslianti, E.R., Bachtia, B.M., Suniart, D.F., and Sutjiat, A.B., 2015. Antioxidant Activity of Rambutan Honey: The Free Radical-Scavenging Activity in vitro and Lipid Peroxidation Inhibition of Oral Mucosa Wound Tissue in vivo. *Research Gate* 9, 284–292.
- Zhang, H., Liu, R., and Tsao, R., 2016. Anthocyanin-rich phenolic extracts of purple root vegetables inhibit pro-inflammatory cytokines induced by H₂O₂ and enhance antioxidant enzyme activities in Caco-2 cells. *J. Funct. Foods* 22, 363–375.