

BIBLIOGRAPHY

- Agusrly, C., Sungkar, T. and Siregar, G. A. (2020) 'Relationship Between Staging and Carcinoembryonic Antigen Serum Levels In Colorectal Cancer Patients', *Journal of Endocrinology, Tropical Medicine, and Infectious Disease (JETROMI)*, 2(3), pp. 118–123. doi: 10.32734/jetromi.v2i3.3959.
- Baran, B. *et al.* (2018) 'Difference Between Left-Sided and Right-Sided Colorectal Cancer: A Focused Review of Literature', *Gastroenterology Research*, 11(4), pp. 264–273. doi: 10.14740/gr1062w.
- Beauchemin, N. and Arabzadeh, A. (2013) 'Carcinoembryonic antigen-related cell adhesion molecules (CEACAMs) in cancer progression and metastasis', *Cancer and Metastasis Reviews*, 32(3–4), pp. 643–671. doi: 10.1007/s10555-013-9444-6.
- Candrawati, O., Eko Broto Hari Utomo, B. and Sofi, L. (2018) 'Correlation of neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio, lymphocyte-to-monocyte ratio and carcinoembryonic antigen level in colorectal cancer', *JKKI: Jurnal Kedokteran dan Kesehatan Indonesia*, 9(2), pp. 82–88. doi: 10.20885/JKKI.VOL9.ISS2.ART4.
- canreg.fk.ugm.ac.id – Web Lembaga Civitas Akademik UGM* (no date). Available at: <https://canreg.fk.ugm.ac.id/> (Accessed: 21 December 2021).
- Chiang, S. F. *et al.* (2012) 'Can neutrophil-to-lymphocyte ratio predict the survival of colorectal cancer patients who have received curative surgery electively?', *International journal of colorectal disease*, 27(10), pp. 1347–1357. doi: 10.1007/S00384-012-1459-X.
- Colloca, G. A., Venturino, A. and Guarneri, D. (2020) 'Different variables predict the outcome of patients with synchronous versus metachronous metastases of colorectal cancer', *Clinical & translational oncology : official publication of the Federation of Spanish Oncology Societies and of the National Cancer Institute of Mexico*, 22(8), pp. 1399–1406. doi: 10.1007/S12094-019-02277-7.
- Elferink, M. A. G. *et al.* (2015) 'Metachronous metastases from colorectal cancer: a population-based study in North-East Netherlands', *International Journal of Colorectal Disease*, 30(2), pp. 205–212. doi: 10.1007/s00384-014-2085-6.
- Filip, S. *et al.* (2020a) 'Distant Metastasis in Colorectal Cancer Patients-Do We Have New Predicting Clinicopathological and Molecular Biomarkers? A Comprehensive Review', *International journal of molecular sciences*, 21(15), pp. 1–24. doi: 10.3390/IJMS21155255.
- Filip, S. *et al.* (2020b) 'Distant metastasis in colorectal cancer patients—do we have new predicting clinicopathological and molecular biomarkers? A comprehensive review', *International Journal of Molecular Sciences*, 21(15), pp. 1–24. doi: 10.3390/ijms21155255.
- GLOBOCAN 2020: New Global Cancer Data | UICC* (no date). Available at:

<https://www.uicc.org/news/globocan-2020-new-global-cancer-data> (Accessed: 21 December 2021).

Goldstein, M. and Mitchell, E. P. (2005) 'Carcinoembryonic Antigen in the Staging and Follow-up of Patients with Colorectal Cancer', *Cancer Investigation*, 23(4), pp. 338–351. doi: 10.1081/cnv-200058878.

Gusti, I. *et al.* (2018) 'HUBUNGAN ANTARA KADAR CARCINOEMBRYONIC ANTIGEN (CEA) DAN STADIUM KANKER KOLOREKTAL DI RSUP SANGLAH TAHUN 2016-2017', *E-Jurnal Medika Udayana*, 7(12), pp. 2303–1395. Available at: <https://ojs.unud.ac.id/index.php/eum/article/view/45062> (Accessed: 20 December 2021).

He, W. *et al.* (2013) 'Initial neutrophil lymphocyte ratio is superior to platelet lymphocyte ratio as an adverse prognostic and predictive factor in metastatic colorectal cancer', *Medical Oncology*, 30(1). doi: 10.1007/s12032-012-0439-x.

Hung, H. Y. *et al.* (2011) 'Effect of preoperative neutrophil-lymphocyte ratio on the surgical outcomes of stage II colon cancer patients who do not receive adjuvant chemotherapy', *International journal of colorectal disease*, 26(8), pp. 1059–1065. doi: 10.1007/S00384-011-1192-X.

Kamiyama, H. *et al.* (2014a) 'Molecular biomarkers for the detection of metastatic colorectal cancer cells', *World Journal of Gastroenterology: WJG*, 20(27), p. 8928. doi: 10.3748/WJG.V20.I27.8928.

Kamiyama, H. *et al.* (2014b) 'Molecular biomarkers for the detection of metastatic colorectal cancer cells', *World Journal of Gastroenterology: WJG*, 20(27), p. 8928. doi: 10.3748/WJG.V20.I27.8928.

Kato, T. *et al.* (2016) 'Tumor size is an independent risk predictor for metachronous colorectal cancer', *Oncotarget*, 7(14), pp. 17896–17904. doi: 10.18632/oncotarget.7555.

Kim, H. *et al.* (2019) 'Preoperative neutrophil-lymphocyte ratio and CEA is associated with poor prognosis in patients with synchronous colorectal cancer liver metastasis', *Annals of Surgical Treatment and Research*, 96(4), pp. 191–200. doi: 10.4174/astr.2019.96.4.191.

Kim, H. J. *et al.* (2008) 'Noninvasive molecular biomarkers for the detection of colorectal cancer', *BMB reports*, 41(10), pp. 685–692. doi: 10.5483/BMBREP.2008.41.10.685.

Laubert, T. *et al.* (2013) 'Aneuploidy and elevated CEA indicate an increased risk for metachronous metastasis in colorectal cancer', *International journal of colorectal disease*, 28(6), pp. 767–775. doi: 10.1007/S00384-012-1625-1.

Luo, D. *et al.* (2021) 'The correlation between tumor size, lymph node status, distant metastases and mortality in rectal cancer patients without neoadjuvant therapy', *Journal of Cancer*, 12(6), p. 1616. doi: 10.7150/JCA.52165.

- Mallappa, S. *et al.* (2013) 'Preoperative neutrophil to lymphocyte ratio >5 is a prognostic factor for recurrent colorectal cancer', *Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland*, 15(3), pp. 323–328. doi: 10.1111/CODI.12008.
- Maradjabessy, F. *et al.* (2018) 'Hubungan Nilai Carcinoembryonic Antigen dengan Kejadian Metastasis Karsinoma Kolorektal di RSUP Prof. Dr. R. D. Kandou Manado', *Jurnal Biomedik (Jbm)*, 10(1), pp. 49–54. doi: 10.35790/jbm.10.1.2018.19002.
- McNally, G. E., Lloyd, D. M. and Grondona, J. P. (2015) 'Carcinoembryonic Antigen as a Prognostic Factor in Colorectal Cancer with Liver Metastases', *Journal of Cancer Therapy*, 06(12), pp. 1035–1044. doi: 10.4236/jct.2015.612113.
- Özgehan, G. *et al.* (2014) 'Neutrophil-lymphocyte ratio as a predictive factor for tumor staging in colorectal cancer', *Turkish Journal of Medical Sciences*, 44(3), pp. 365–368. doi: 10.3906/sag-1305-33.
- Schmoll, H. J. *et al.* (2012) 'Esmo consensus guidelines for management of patients with colon and rectal cancer. A personalized approach to clinical decision making', *Annals of Oncology*, 23(10), pp. 2479–2516. doi: 10.1093/ANNONC/MDS236.
- Takagawa, R. *et al.* (2008) 'Preoperative serum carcinoembryonic antigen level as a predictive factor of recurrence after curative resection of colorectal cancer', *Annals of surgical oncology*, 15(12), pp. 3433–3439. doi: 10.1245/S10434-008-0168-8.
- Thomas, P., Forse, R. A. and Bajenova, O. (2011) 'Carcinoembryonic antigen (CEA) and its receptor hnRNP M are mediators of metastasis and the inflammatory response in the liver', *Clinical & experimental metastasis*, 28(8), pp. 923–932. doi: 10.1007/S10585-011-9419-3.
- Tsai, P. L. *et al.* (2016) 'Neutrophil-lymphocyte ratio and CEA level as prognostic and predictive factors in colorectal cancer: A systematic review and meta-analysis', *Journal of cancer research and therapeutics*, 12(2), pp. 582–589. doi: 10.4103/0973-1482.144356.
- Walsh, S. R. *et al.* (2005) 'Neutrophil-lymphocyte ratio as a prognostic factor in colorectal cancer', *Journal of surgical oncology*, 91(3), pp. 181–184. doi: 10.1002/JSO.20329.
- Wiratkapun, S. *et al.* (2001) 'High preoperative serum carcinoembryonic antigen predicts metastatic recurrence in potentially curative colonic cancer: Results of a five-year study', *Diseases of the Colon and Rectum*, 44(2), pp. 231–235. doi: 10.1007/BF02234298.