

DAFTAR PUSTAKA

- Asdie, R.H., Subronto, Y.W., Kurniawan, J., 2020. Persistent lymphopenia in septic patients at Dr. Sardjito General Hospital, Yogyakarta. *J. the Med. Sci. (Berkala Ilmu Kedokteran)* 52, 37–49. <https://doi.org/10.19106/jmedsci005204202003>
- Assinger, A., Schrottmaier, W.C., Salzmann, M., Rayes, J., 2019. Platelets in sepsis: An update on experimental models and clinical data. *Front. Immunol.* 10, 1–19. <https://doi.org/10.3389/fimmu.2019.01687>
- Banerjee, D. and Opal, S.M. (2017). Age, exercise, and the outcome of sepsis. *Critical Care*, 21(1).
- Chen, Y., Feng, F., Li, M., Yuan, J.J., Chang, X.N., Wei, B.H., Du, H., Dong, C.M., 2020. Relationship between platelet/lymphocyte ratio and prognosis of patients with septic acute kidney injury: A pilot study. *J. Chinese Med. Assoc.* 83, 1004–1007. <https://doi.org/10.1097/JCMA.0000000000000404>
- Chou, E.H., Mann, S., Hsu, T.-C., Hsu, W.-T., Liu, C.C.-Y., Bhakta, T., Hassani, D.M. and Lee, C.-C. (2020). Incidence, trends, and outcomes of infection sites among hospitalizations of sepsis: A nationwide study. *PLOS ONE*, [online] 15(1), p.e0227752. <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0227752>
- De Pablo, R., Monserrat, J., Prieto, A., Alvarez-Mon, M., 2014. Role of Circulating Lymphocytes in Patients with Sepsis. *Biomed Res. Int.* 2014. <https://doi.org/10.1155/2014/671087>
- Djordjevic, D., Rondovic, G., Surbatovic, M., Stanojevic, I., Udovicic, I., Andjelic, T., Zeba, S., Milosavljevic, S., Stankovic, N., Abazovic, D., Jevdjic, J. and Vojvodic, D. (2018). Neutrophil-to-Lymphocyte Ratio, Monocyte-to-Lymphocyte Ratio, Platelet-to-Lymphocyte Ratio, and Mean Platelet Volume-to-Platelet Count Ratio as Biomarkers in Critically Ill and Injured Patients: Which Ratio to Choose to Predict Outcome and Nature of Bacteremia? *Mediators of Inflammation*, 2018, pp.1–15.
- Fleischmann, C., Scherag, A., Adhikari, N.K.J., Hartog, C.S., Tsaganos, T., Schlattmann, P., Angus, D.C. and Reinhart, K. (2016). Assessment of Global Incidence and Mortality of Hospital-treated Sepsis. Current Estimates and Limitations. *American Journal of Respiratory and Critical Care Medicine*, 193(3), pp.259–272.
- Fujishima, S., 2016. Organ dysfunction as a new standard for defining sepsis. *Inflamm. Regen.* 36, 1–6. <https://doi.org/10.1186/s41232-016-0029-y>
- Hajian-Tilaki, K. (2013). Receiver Operator Characteristic Analysis of Biomarkers Evaluation in Diagnostic Research. *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH*.

- Hedi, B., Fitri, E.Y. and Hikayati (2017). Faktor-Faktor yang Berhubungan dengan Kejadian Systemic Inflammatory Response Syndrome di Ruang ICU RSUD Lahat. *Jurnal Keperawatan Sriwijaya*, 4(1).
- Kempker, J.A. and Martin, G.S. (2016a). The Changing Epidemiology and Definitions of Sepsis. *Clinics in Chest Medicine*, 37(2), pp.165–179.
- Liu, X., Shen, Y., Wang, H., Ge, Q., Fei, A., Pan, S., 2016. Prognostic Significance of Neutrophil-to-Lymphocyte Ratio in Patients with Sepsis: A Prospective Observational Study. *Mediators Inflamm.* 2016. <https://doi.org/10.1155/2016/8191254>
- Martin-Loeches, I., Guia, M.C., Valleccoccia, M.S., Suarez, D., Ibarz, M., Irazabal, M., Ferrer, R. and Artigas, A. (2019). Risk factors for mortality in elderly and very elderly critically ill patients with sepsis: a prospective, observational, multicenter cohort study. *Annals of Intensive Care*, 9(1).
- Martins, E.C., Silveira, L. da F., Viegas, K., Beck, A.D., Fioravanti Júnior, G., Cremonese, R.V. and Lora, P.S. (2019). Neutrophil-lymphocyte ratio in the early diagnosis of sepsis in an intensive care unit: a case-control study.
- Murphy, K., Weaver, C., 2017. Janeway's Immunobiology, 9th ed. Garland Science, Taylor & Francis Group, LLC, New York.
- Ni, J., Wang, H., Li, Y., Shu, Y. and Liu, Y. (2019). Neutrophil to lymphocyte ratio (NLR) as a prognostic marker for in-hospital mortality of patients with sepsis. *Medicine*, 98(46), p.e18029.
- Novosad, S.A., Sapiano, M.R.P., Grigg, C., Lake, J., Robyn, M., Dumyati, G., Felsen, C., Blog, D., Dufort, E., Zansky, S., Wiedeman, K., Avery, L., Dantes, R.B., Jernigan, J.A., Magill, S.S., Fiore, A. and Epstein, L. (2016). Vital Signs: Epidemiology of Sepsis: Prevalence of Health Care Factors and Opportunities for Prevention. *Wilderness & Environmental Medicine*, 22(1), p.95.
- Pc De Jager, C., Tl Van Wijk, P., Mathoera, R., De Jongh-Leuvenink, J., Van Der Poll, T. and Wever, P. (2010). *Lymphocytopenia and neutrophil-lymphocyte count ratio predict bacteremia better than conventional infection markers in an emergency care unit.*
- Pepper, D.J., Sun, J., Welsh, J., Cui, X., Suffredini, A.F. and Eichacker, P.Q. (2016). Increased body mass index and adjusted mortality in ICU patients with sepsis or septic shock: a systematic review and meta-analysis. *Critical Care*, 20(1).
- Riché, F., Gayat, E., Barthélémy, R., Le Dorze, M., Matéo, J. and Payen, D. (2015). Reversal of neutrophil-to-lymphocyte count ratio in early versus late death from septic shock. *Critical Care*, [online] 19(1).
- Rudd, K.E., Johnson, S.C., Agesa, K.M., Shackelford, K.A., Tsoi, D., Kievlan, D.R., Colombara, D.V., Ikuta, K.S., Kissoon, N., Finfer, S., Fleischmann-Struzek, C., Machado, F.R., Reinhart, K.K., Rowan, K., Seymour, C.W., Watson, R.S., West, T.E., Marinho, F., Hay, S.I. and Lozano, R. (2017). *Global, regional,*

- and national sepsis incidence and mortality, 1990–2017: analysis for the Global Burden of Disease Study. The Lancet*, 395(10219), 200–211 / 10.1016/S0140-6736(19)32989-7.
- Sato, T., Kudo, D., Kushimoto, S., Hasegawa, M., Ito, F., Yamanouchi, S., Honda, H., Andoh, K., Furukawa, H., Yamada, Y., Tsujimoto, Y., Okuyama, M. and Kobayashi, M. (2021). Associations between low body mass index and mortality in patients with sepsis: A retrospective analysis of a cohort study in Japan. *PLOS ONE*, [online] 16(6), p.e0252955.
- Shannon, O., 2020. The role of platelets in sepsis. *Res. Pract. Thromb. Haemost.* 5, 27–37. <https://doi.org/10.1002/rth2.12465>
- Shen, Y., Huang, X., Zhang, W., 2019. Platelet-to-lymphocyte ratio as a prognostic predictor of mortality for sepsis: Interaction effect with disease severity - A retrospective study. *BMJ Open* 9, 1–7. <https://doi.org/10.1136/bmjopen-2018-022896>
- Sharma, M., Jain, N., Sinha, N., Kaushik, R., Jash, D. and Chaudhry, A. (2018). Diagnostic and Prognostic Role of Neutrophil-to-Lymphocyte Ratio in Early and Late Phase of Sepsis. *Indian Journal of Critical Care Medicine*, [online] 22(9), pp.660–663.
- Singer, M., Deutschman, C.S., Seymour, C.W., Shankar-Hari, M., Annane, D., Bauer, M., Bellomo, R., Bernard, G.R., Chiche, J.-D., Coopersmith, C.M., Hotchkiss, R.S., Levy, M.M., Marshall, J.C., Martin, G.S., Opal, S.M., Rubenfeld, G.D., Van Der Poll, T., Vincent, J.-L., Angus, D.C., 2016. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3) HHS Public Access. *JAMA* 315, 801–810. <https://doi.org/10.1001/jama.2016.0287>
- Susanto, J., Pradian, E., Zulfariansyah, A., 2019. Correlation Between Neutrophil-lymphocyte Count Ratio and Procalcitonin in Sepsis and Septic Shock Korelasi antara Rasio Hitung Neutrofil-Limfosit dan Prokalsitonin pada Pasien Sepsis dan Syok Sepsis. *Maj. Kedokt. Bandung* 51.
- Wang, S., Liu, X., Chen, Q., Liu, C., Huang, C. and Fang, X. (2017). The role of increased body mass index in outcomes of sepsis: a systematic review and meta-analysis. *BMC Anesthesiology*, [online] 17(1).
- Wartenberg, K.E., Stoll, A., Funk, A., Meyer, A., Schmidt, J.M. and Berrouschot, J. (2011). Infection after Acute Ischemic Stroke: Risk Factors, Biomarkers, and Outcome. *Stroke Research and Treatment*, 2011, pp.1–8.
- Whiles, B., Deis, A., Miller, P. and Simpson, S. (2016). Comorbid Conditions Predict Outcomes in Patients With Severe Sepsis.
- World Health Organization: WHO (2020). *Sepsis*. [online] Who.int. Available at: <https://www.who.int/news-room/fact-sheets/detail/sepsis> [Diakses 4 Mar. 2021].
- Zhang, F., Liu, A.L., Gao, S., Ma, S., Guo, S. Bin, 2016. Neutrophil dysfunction in

sepsis. Chin. Med. J. (Engl). 129, 2741–2744. <https://doi.org/10.4103/0366-6999.193447>

Zou, K.H., O'Malley, A.J. and Mauri, L. (2007). Receiver-Operating Characteristic Analysis for Evaluating Diagnostic Tests and Predictive Models. *Circulation*, 115(5), pp.654–657.