

DAFTAR PUSTAKA

- Abd El-Samad, H.M., Shaddah, M.A.K., & Barakat, N. (2011). Improvement of plants salt tolerance by exogenous application of amino acids. *J. Med. Plant Res.* 5:5692-5699.
- Abdelhamid, M.T., Rady, M., Osman, A., & Abdalla, M. (2013). Exogenous application of proline alleviates salt-induced oxidative stress in *Phaseolus vulgaris* L. *Plants. J. Hortic Sci Biotechnol.* 88: 439-446.
- Afzal, I., Butt, A., Ur Rehman, H., Basra, S.M.A., & Afzal, A. (2012). Alleviation of salt stress in fine aromatic rice by seed priming. *Aust. J. Crop Sci.* 6(10): 1401-1407.
- Argawal, P., Agarwal, P.K., & Gohil, K. (2018). Transcription factor-based genetic engineering for salinity tolerance in crops. *Salinity Responses and Tolerance in Plant.* 1:185-211.
- Aghbolaghi, M.A., & Sedghi, M. (2013). The effect of halo-and hydro-priming on germination characteristics of millet seeds under salinity stress. *Cercetari Agronomice in Moldova.* 57(2): 42-51.
- Amirjani, M.R. (2011). Effect of salinity stress on growth, sugar content, pigments and enzyme activity of rice. *Intern J. Bot.* 7(1): 73-81.
- Aggarwal, M. et al. (2011). Exogenous proline application reduces phytotoxic effects of selenium by minimising oxidative stress and improves growth in bean (*Phaseolus vulgaris* L.) seedlings. *Biol. Trace Element Res.* 140: 354-367.
- Ansari, O., Choghazardi, H.R., Sharif, Z. F., & Nazarli, H. (2012). Seed reserve utilization and seedling growth of treated seeds of mountain ray (*Secale montanum*) as affected by drought stress. *Cercetari Agronomice in Moldova.* 2(150): 43-48.
- Ashraf, M., Athar, H.R., Harris, P.J.C., & Kwon, T.R. (2008). Some prospective strategies for improving crop salt tolerance. *Adv Agron.* 45-59.
- Balai Besar Penelitian dan Pengembangan Bioteknologi dan Sumberdaya Genetik Pertanian. (2009). Beras hitam, pangan berkhasiat yang belum populer. *Warta Penelitian dan Pengembangan Pertanian.* 13(2): 1-2.
- Bhatt, T., Sharma, A., Puri, S., & Minhas, A.P. (2020). Salt tolerance mechanisms and approaches: future scope of halotolerant genes and rice landraces. *Rice Sci.* 27(5): 368-383.
- Binzel, M.L. et al. (1987). Solute accumulation in tobacco cells adapted to NaCl. *Plant Physiol.* 84(4): 1408-1415.
- Celik, O., & Atak, C. (2011). Evaluation of proline accumulation and $\Delta 1$ -pyrroline-5-carboxylate synthetase (P5CS) gene expression during salinity stress in two soybean (*Glycine max* L. Merr) varieties. *Pol. J. Environ. Stud.* 12(3): 559-564.
- Cha-Um, S., & Kirdmanee, C. (2009). Effect of salt stress on proline accumulation, photosynthetic ability and growth characters in two maize cultivars. *Pak. J. Bot.* 87-98.
- Chen, K., & Arora, R. (2013). Priming memory invokes seed stress-tolerance. *Environ. Exp. Bot.* 94: 33-45.

- Chen, Z. et al. (2007). Compatible solute accumulation and stress-mitigating effects in barley genotypes contrasting in their salt tolerance. *J.Exp Botany*. 15(16): 4245-4255.
- Delavari, P.M. et al. (2010). The effect of salicylic acid on some of biochemical and morphological characteristic of *Ocimum basilicum* under salinity stress. *Aust. J. Basic. Appl. Sci.* 4:4832-4845.
- Dong, W., Liu, X., Li, D., Gao, T., & Song, Y. (2018). Transcriptional profiling reveals that a MYB transcription factor *MsMYB4* contributes to the salinity stress response of alfalfa. *PLoS ONE*. 13(9): 1-18.
- Dorna, H., Li, W., & Szopinska, D. (2014). The effect of priming on germination and vigour of pansy (*Viola x wittrockiana* Gams.) seeds. *Acta Sci. Pol., Hortorum Cultus*. 13 (6): 15-29.
- Ganie, S.A., Molla, K.A., Henry, R.K., Bhat, K.V., & Mondal, T.P. (2019). Advance in understanding salt tolerance in rice. *Theor Appl Genet*. 1-20.
- Gujjar, R.S, & Akhtar, M. (2014). Transcription factors in abiotic stress tolerance. *Ind. J. Plant Physiol*. 19(4): 306-316.
- Hamidi, R., Pirasteh-Anosheh, H., & Izadi, M. (2013). Effect of seed halo-priming compared with hydro-priming on wheat germination and growth. *Int J Agron Plant Prod*. 4(7): 1611-1615.
- Hadinezhad, P., Payamenur, V., Mohamadi, J., & Ghaderifar, F. (2013). The effect of priming on seed germination and seedling growth in *Quercus castaneifolia*. *Seed Sci. Technol*. 41: 121-124.
- Hidayah, A. 2020. *Toleransi Padi Hitam (Oryza sativa Linn.) Cempo Ireng Pendek Terhadap Cekaman Salinitas Berdasarkan Level Ekspresi Gen Transporter dan Metabolit Compatible Solute*. Tesis. Sekolah Pascasarjana Universitas Gadjah Mada. Yogyakarta. p. 1-86.
- Hoang, X.L., Hai Nhi, D.N., Anh Thu, N.B., Thao, N.P., & Tran, L-S.P. (2013). Review article: transcription factors and their roles in signal transduction in plants under abiotic stresses. *Curr Gen*. 18: 483-497.
- Hoang, T.M.L. et al. (2016). Improvement of salinity stress tolerance in rice: challenge and opportunities. *Agron*. 6: 1-23.
- Hoque, M.A., Banu, M.N.A., Okuma, K., & Nakamura, Y. (2007). Exogenous proline and glycinebetaine increase NaCl-induced ascorbate-glutathione cycle enzyme activities, and proline improves salt tolerance more than glycinebetaine in tobacco bright yellow-2-suspension-cultured cells. *J. Plant Physiol*. 164: 1457-1468.
- Hossain, M.R. (2013). *Salinity tolerance and transcriptomics in rice*. Thesis. University of Birmingham. England.
- Hur, J., Kung, K., Lee, C-H., & An, G. (2004). Stress-inducible *OsP5CS2* gene is essential for salt and cold tolerance in rice. *Plant Sci*. 1133: 83-94.

- Hussain, S. et al. (2017). Effects of salt stress on rice growth, development characteristics, and the regulating ways: A review. *J. Integr. Agric.* 16(11): 2357-2374.
- Hyun, J.W., & Chung, H.S. (2004). Cyanidin and malvidin from *Oryza sativa* cv. Heugjinjubyeo mediate cytotoxicity against human monocytic leukemia cells by arrest of G(2)/M phase and induction of apoptosis. *J. Agric. Food Chem.* 52: 2213-2217.
- Ibrahim, E.A. (2016). Seed priming to alleviate salinity stress in germinating seeds. *J. Plant Physiol.* 1-15.
- Jahan, F., Bhusan, D., Jahiruddin, M., Murata, Y., & Hoque, M.A. (2018). Improvement of salinity tolerance in rice during boro season by proline application. *Progress. Agric.* 29(4): 295-303.
- Jha, S. (2019). Transgenic Approaches for Enhancement of Salinity Stress Tolerance in Plants. Chapter 14. Springer Nature Singapore Pte Ltd. S.P.Singh et al. (eds.), Molecular Approaches in Plant Biology and Environmental Challenges, Energy, Environment, and Sustainability. 1-59.
- Joseph, E.A., Radhakrishnan, V.V., & Mohanan, K.V. (2015). A study on accumulation of proline-an osmoprotectant amino acid under salt stress in some native rice cultivars of North Kerala, India. *Univ J. Agr. Res.* 3(1): 15-22.
- Joshi, N., Jain, A., & Arya, K. (2013). Alleviation of salt stress in *Curcuma sativa* L. through seed priming with calcium chloride. *Indian J Appl Res.* 3(11):22-25.
- Jyotsna, V., & Srivastava, A.K. (1998). Physiological basis of salt stress resistance in pigeon pea (*Cajanus cajan* L.). *Plant Physiol Biochem.* 2:89-94.
- Julkowska, M.M., & Testerink, C. (2015). Turning plant signalling and growth to survive salt. *Trend in Plant Sci.* 20(9): 586-594
- Karolinoerita, V., & Yusuf, W.A. (2020). Salinisasi Lahan dan Permasalahannya di Indonesia. *J Sumberdaya Lahan.* 14(2): 91-99
- Khong, G.N. et al. (2013). Modulating rice stress tolerance by transcription factor. *Biotechnol. Genet Eng. Rev.* 325-403.
- Kordrostami, M & Rabiei, B. (2019). *Salinity Stress Tolerance in Plants: Physiological, Molecular, and Biotechnological Approaches*. Springer.
- Kristantini, S. Widyayanti, Sutarno, & Sudarmaji. (2009). Keragaman genetik lima kultivar lokal padi beras hitam asal Yogyakarta berdasarkan sifat morfologi. *Prosiding Seminar Nasional Sumber Daya Genetik Pertanian.* 90-100.
- Kristantini, Taryono, Basunanda, P., & Murti, R.H. (2014). Keragaman genetik kultivar padi beras hitam lokal berdasarkan penanda mikrosatelit. *J. AgroBiogen.* 10(2): 69-76.
- Kurniasih, N.S. (2018). *Karakterisasi padi (Oryza sativa L.) lokal berpigmen di Indonesia berbasis SNPs (Single Nucleotide Polymorphism) dan morfologi*. Tesis Program Studi Biologi. Fakultas Biologi, Universitas Gadjah Mada. Yogyakarta. 1-63.

- Lafitte, H.R., Ismail, A., & Bennet, J. (2004). *Abiotic stress tolerance in rice Fore Asia progress and the future*. International Rice Research Institute. Metro Manila, Philippines.
- Lamke, J., & Baurle, I. (2017). Review: epigenetic and chromatin-based mechanisms in environmental stress adaptation and stress memory in plants. *Genom Biol.* 18(124): 1-11.
- Li, J., Han, G., Sun, C., & Sui, N. (2019). Research advances of MYB transcription factors in plant stress resistance and breeding. *Plant Signal and Behav.* 14(8): 1-10.
- Liu, L., Du, H., & Tang, X. (2008). The roles of MYB transcription factors on plant defense responses ananism. *J Hereditas.* 30: 1265-1271.
- Livak, K.J., & Schmittgen, T.D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the $2^{\Delta\Delta CT}$ method. *Methods.* 25(4): 402-408.
- Maggio, A. et al. (2002). Does proline accumulation play an active role in stress-induced growth reduction. *Plant J.* 31:699-712.
- Mansuri, R.M., Shobbar, Z-S., Jelodar, N.B., & Ghaffari, M.R. (2019). Disecting molecular mechanism underlying salt tolerance in rice: a comparative transcriptional profiling of the contrasting genotypes. *Rice.* 12: 1-13
- Marthandan, V. et al. (2020). Seed priming: a feasible strategy to enhance drought tolerance in crop plants. *International Journal of Molecular Sciences.* 8258: 1-23.
- Mishra, A., & Bhakti, T. (2017). Halophytes: potential resources for salt stress tolerance genes and promoters. *Plant Sci.* 8(219): 1-8
- Mums, R., & Tester, M. (2008). Mechanism of salinity tolerance. *Annual Review of Plant Biology.* 59(1): 651-681
- Munns, R. (2011). Plant adaptation to salt and water stress. Differences and commonalities. *Adv. Bot. Res.* 57: 1-32.
- Nam, S. et al. (2006). Antioxidative activities of bran extract from twenty one pigmented rice cultivars. *Food Chem.* 94: 613-620.
- Nasri, N. et al. (2011). The effect of osmopriming on germination, seedling growth and phosphatase activities of lettuce under saline condition. *Afr. J. Biotechnol.* 10: 14366-14372.
- Nguyen, H.T.T., Bhowmik, S.D., Long, H., Cheng, Y., & Mundree, S. (2021). Rapid accumulation of proline enhances salinity tolerance in Australian wild rice *Oryza australiensis* Domin. *Plants.* 10: 1-16.
- Nisak, R.R. (2019). *Respon Biokimia dan Ekspresi Gen OsSOS1, OsNHX1 dan OsHKT1;1 terhadap cekaman salinitas pada padi (Oryza saruva L. 'Merah') Kalimantan Selatan*. Tesis. Sekolah Pascasarjana Universitas Gadjah Mada. Yogyakarta. 1-113.
- Olieveira, A.B., Gomes-Filho, E. Eneas-Filho, J., Prisco, J.T., & Alencar, N.L.M.A. (2012). Seed priming effects on growth, lipid peroxidation, and activity of ROS scavenging enzymes in NaCl-stressed sorghum seedlings from aged seeds. *J. Plant Interact.* 7:151-159.

- Parvaiz, A., & Satyawati, S. (2008). Salt stress and phyto-biochemical responses of plants. *Plant Soil Environ.* 54(3): 89-99.
- Rachman, A., I.G.M., Subiksa, & Wahyunto. (2017). Perluasan Areal Tanaman Kedelai ke Lahan Suboptimal. *Kedelai:: Teknik Produksi dan Pengembangan.* 185-204.
- Ranganayakulu, G.S., Veeranagamallaiah, G., & Sudhakar, C. (2013). Effect of salt stress on osmolyte accumulation in two groundnut cultivars (*Arachis hypogaea* L.) with contrasting salt tolerance. *Afr. J. Plan. Sci.* 7(12): 586-592.
- Razzaque, S. et al. (2019). Gene expression analysis associated with salt stress in a reciprocally crossed rice population. *Sci. Rep.* 9: 1-17.
- Reddy, A.S., Ali, G.S., Celesnik, H., & Day, I.S. (2011). Coping with stresses: role of calcium and calcium/calmodulin-regulated gene expression. *Plant Cell.* 23(6): 2010-2032.
- Roy, S., & Chakraborty, U. (2014). Salt tolerance mechanism in salt tolerant grasses (STGs) and their prospect in cereal crop improvement. *Bot. Stud.* 55:31.
- Safitri, H., Purwoko, B.S., Dewi, I.S., & Ardie, S.W. (2017). Salinity tolerance of several rice genotype at seedling stage. *Indones J. Agric. Sci.* 18(2): 63-68.
- Santos, A.P. et al. (2011). Transcription regulation of abiotic stress response in rice: a combined action of transcription factors and epigenetic mechanisms. *OMICS: Int. J. Integr. Biol.* 15(12): 839-857.
- Sarwar, M., Amjad, M., & Ayyub, C.M. (2017). Alleviation of salt stress in cucumber (*Cucumis sativus* L.) through seed priming with triacontanol. *Int. J. Agric. Biol.* 19: 771-778.
- Shahid, M.A. et al. (2020). Insights into the physiological and biochemical impacts of salt stress on plant growth and development. *Agron.* 10: 1-34.
- Shiddique, A., Kandpla, G., & Kumar, P. (2018). Proline accumulation and its defensive role under diverse stress condition in plants: an overview. *J. Pure Appl. Microbiol.* 12(3):1655- 1659.
- Singh, M. Kumar, J., Singh, V.P., & Prasad, SM. (2014). Proline and salinity tolerance in plants. *Biochem.Pharmacol.* 3(6): 1-2.
- Sivritepe, N., Sivritepe, H.O., & Eris, A. (2003). The effect of NaCl priming on salt tolerance in melon seedling grown under saline conditions. *Sci Hortic.* 97: 229-237.
- Smart, R.E., & Bingham, G.E. (1974). Rapid estimates of relative water content. *Plant Physiol.* 53: 258-260.
- Srivastava, A.K., Kumar, J.S., & Suprasanna, P. (2021). Seed primeomics : plant memorize their germination under stress. *Biol. Rev.* 1-22.
- Tabassum, T., Farooq, M., Ahmad, R., Zohaib A., & Wahid, A. (2017). Seed priming and transgenerational drought improves tolerance against salt stress in bread wheat. *Plant Physiol Biochem.* 118: 362-369.
- Tananuwong, K., & Tewaruth, W. (2010). Extraction and application of antioxidant from black glutinous rice. *LWT-Food Sci. Technol.* 43(3): 476-481.

- Tang, Y. et al. (2019). Overexpression of a MYB family gene, *OsMYB6*, increases drought and salinity stress tolerance in transgenic rice. *Plant Sci.* 10(168): 1-12.
- Theerakalpisut, P., Kanawapee, N., & Panwong, B. (2016). Seed priming alleviated salt stress effects on rice seedlings by improving Na^+/K^+ and maintaining membrane integrity. *Int. J. Plant. Biol.* 7: 53-60.
- Tsuda, T., Horio, F., and Osawa, T. (2002). Cyanidin 3-O-beta-D-glucoside supresses nitric oxide production during a zymosan treatment in rats. *J. Nutr. Sci. Vitaminol.* 48(4): 305-310.
- Tsuda, T., Horio, F., Uchida, K., Aoki, H., & Osawa, T. (2003). Dietary cyanidin 3-O-beta-D-glucoside-rice purple corn color prevent pbesity and ameliorates hyperglycemia in mice. *J. Nutr.* 133(7): 2125-2130.
- Tuteja, N. (2007). Mechanism of high salinity tolerance in plants. *Meth. Enzymol.* 428: 419-438.
- USSL. 2005. George E, Brown, Jr. Salinity Laboratory. Riverside, CA, USA:USDA-ARS. <http://www.ussl.ars.usda.gov>
- Vaktabhai, C.K. & Kumar, S. (2017). Seedling invigouration by halo priming in tomato against salt stress. *J. Pharmacogn. Phytochem.* 6(6): 716-722.
- Xiong, H. et al. (2014). Overexpression of *OsMYB48-1*, a novel MYB-related transcription factor, enhances drought and salinity tolerance in rice. *PLoS ONE.* 9(3): 1-13.
- Yang, A., Dai, X., & Zhang, W-H. (2012). A R2R3-type MYB gene, *OsMYB2*, is involved in salt, cold, and dehydration tolerance in rice. *J. Exp. Bot.* 63(7): 2541-2556.
- Yawadio, R., Tanimori, S., & Morita, N. (2007). Identification of phenolic compounds isolated from pigmented rices and their adolase reductase inhibitory activities. *Food Chem.* 101: 1616-1625.
- Zarattini, M., & Forlani, G. (2017). Toward unveiling the mechanisms for transcriptional regulation of proline biosynthesis in the plant cell response to biotic and abiotic stress conditions. *Plant Sc.* 8: 1-8.
- Zhang, J.L., & Shi, H. (2013). Physiological and molecular mechanisms of plant salt tolerance. *Photosynth. Res.* 115: 1-22.
- Zhu, A. et al. (2015). The R2R3-type MYB gene *OsMYB91* has a function in coordinating plant growth and salt stress tolerance in rice. *Plant Sci.* 236:146-156.