

REFERENCES

- Agusrly, C., Sungkar, T. and Siregar, G. A. (2020) 'Relationship Between Staging and Carcinoembryonic Antigen Serum Levels In Colorectal Cancer Patients', *Journal of Endocrinology, Tropical Medicine, and Infectious Disease (JETROMI)*, 2(3), pp. 118–123. doi: 10.32734/jetromi.v2i3.3959.
- Agusrly, C., Sungkar, T. and Siregar, G.A., 2020. Comparison of Carcinoembryonic Antigen Serum Levels in Colorectal Cancer Patients with Different Histopathological Grades. *Majalah Kedokteran Bandung*, 52(2), pp.87-91.
- Asad-Ur-Rahman, F. and Saif, M. W. (2016) 'Elevated Level of Serum Carcinoembryonic Antigen (CEA) and Search for a Malignancy: A Case Report', *Cureus*, 8(6), pp. 8–11. doi: 10.7759/cureus.648.
- Aykan, N.F., Artac, M. and Özatlı, T., 2018. Body Mass Index and Colorectal Cancer. In *Body-mass Index and Health*. IntechOpen.
- Balta, A.Z., Özdemir, Y., Sücüllü, İ., Derici, S.T., Bağcı, M., Demirel, D. and Akın, M.L., 2014. Can horizontal diameter of colorectal tumor help predict prognosis?. *Turkish Journal of Surgery/Ulusal cerrahi dergisi*, 30(3), p.115.
- Cancerresearchuk.org. 2019. TNM Staging | Bowel Cancer | Cancer Research UK. [online] Available at: <<https://www.cancerresearchuk.org/about-cancer/bowelcancer/stages-types-and-grades/TNM-staging>>
- Candrawati, O., Utomo, B., and Sofi'I I. (2018) 'Correlation of Neutrophil-to-Lymphocyte Ratio, Platelet-to-Lymphocyte ratio, Lymphocyte-to-Monocyte

Ratio and Carcinoembryonic Antigen Level in Colorectal Cancer', Indonesian Journal of Medicine and Health, 9(2), pp. 82-88. doi: 10.20885/JKKI.Vol9.Iss2.art4.

Cdc.gov. 2021. What Is Colorectal Cancer? | CDC. [online] Available at: <https://www.cdc.gov/cancer/colorectal/basic_info/what-is-colorectal-cancer.htm>

Cdc.gov. 2021. *Colorectal Cancer Screening Tests* | CDC. [online] Available at: <https://www.cdc.gov/cancer/colorectal/basic_info/screening/tests.htm> [Accessed 5 January 2022].

Coussens, L. M., & Werb, Z. (2002). Inflammation and cancer. *Nature*, 420(6917), 860–867. <https://doi.org/10.1038/nature01322>

Dai, W., Li, Y., Meng, X., Cai, S., Li, Q. and Cai, G., 2017. Does tumor size have its prognostic role in colorectal cancer? Re-evaluating its value in colorectal adenocarcinoma with different macroscopic growth pattern. *International journal of surgery*, 45, pp.105-112.

Doleman, B., Mills, K.T., Lim, S., Zelhart, M.D. and Gagliardi, G., 2016. Body mass index and colorectal cancer prognosis: a systematic review and meta-analysis. *Techniques in coloproctology*, 20(8), pp.517-535.

Euro.who.int. n.d. *Body mass index - BMI*. [online] Available at: <<https://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi>> [Accessed 9 January 2022].

Freeman, H. J. (2013) 'Early stage colon cancer', *World Journal of Gastroenterology*, 19(46), pp. 8468–8473. doi: 10.3748/wjg.v19.i46.8468.

Global Cancer Observatory. 2012. Cancer Fact Sheets: Colorectal Cancer. [online] Available at: <<https://gco.iarc.fr/today/data/pdf/fact-sheets/cancers/cancer-fact-sheets6.pdf>>

Haggar, F. A. and Boushey, R. P. (2009) 'Colorectal cancer epidemiology: Incidence, mortality, survival, and risk factors', *Clinics in Colon and Rectal Surgery*, 22(4), pp. 191–197. doi: 10.1055/s-0029-1242458.

Hall, C. et al. (2019) 'A review of the role of carcinoembryonic antigen in clinical practice', *Annals of Coloproctology*, 35(6), pp. 294–305. doi: 10.3393/ac.2019.11.13.

Hammarström, S., 1999, April. The carcinoembryonic antigen (CEA) family: structures, suggested functions and expression in normal and malignant tissues. In *Seminars in cancer biology* (Vol. 9, No. 2, pp. 67-81). Academic Press.

Horton, K.M., Abrams, R.A. and Fishman, E.K., 2000. Spiral CT of colon cancer: imaging features and role in management. *Radiographics*, 20(2), pp.419-430.

InformedHealth.org [Internet]. Cologne, Germany: Institute for Quality and Efficiency in Health Care (IQWiG); 2006-. Signs of colorectal cancer. 2006 Feb 14 [Updated 2017 Jan 26]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK279199/>

Jogja Cancer Registry. 2021. RKBR Maret 2021. [online] Available at: <https://canreg.fk.ugm.ac.id/laporan-data/registrasi-kanker-berbasis-rumah-sakit-dr-sardjito-fkkmk-ugm/rkbr-maret-2021/>

Kijima, S., Sasaki, T., Nagata, K., Utano, K., Lefor, A.T. and Sugimoto, H., 2014. Preoperative evaluation of colorectal cancer using CT colonography, MRI, and PET/CT. *World journal of gastroenterology: WJG*, 20(45), p.16964.

Kornprat, P., Pollheimer, M.J., Lindtner, R.A., Schlemmer, A., Rehak, P. and Langner, C., 2011. Value of tumor size as a prognostic variable in colorectal cancer: a critical reappraisal. *American journal of clinical oncology*, 34(1), pp.43-49.

Kubo, H. et al. (2016) 'The Prognostic Value of Preoperative Neutrophil-to-Lymphocyte Ratio in Colorectal Cancer'. doi: 10.1007/s00268-016-3595-x.

Ho, C. H., Yu, Y. Bin and Wu, P. H. (2008) 'The prevalence of iron deficiency anemia and its clinical implications in patients with colorectal carcinoma', *Journal of the Chinese Medical Association*, 71(3), pp. 119–122. doi: 10.1016/S17264901(08)70002-9.

Kuipers, E.J., Grady, W. M., Lieberman, D., Sseufferlein, T., Sung, J. J., Boelens, P. G., Vam de Velde, C. J., & Watanabe, T. (2015). 'Colorectal Cancer', *Nature Reviews. Disease Primers*, 1, 15065. <https://doi.org/10.1038/nrdp.2015.65>

Kumar, V., Abbas, A. and Aster, J., 2014. *Robbins & Cotran Pathologic Basis of Disease*. 9th ed. Elsevier, p.811.

Kwon, H.C., Kim, S.H., Oh, S.Y., Lee, S., Lee, J.H., Choi, H.J., Park, K.J., Roh, M.S.,

Kim, S.G., Kim, H.J. and Lee, J.H., 2012. Clinical significance of preoperative neutrophil-lymphocyte versus platelet-lymphocyte ratio in patients with operable colorectal cancer. *Biomarkers*, 17(3), pp.216-222.

Lalosevic, M. S. et al. (2017) 'Can preoperative CEA and CA19-9 serum concentrations suggest metastatic disease in colorectal cancer patients?', *Hellenic Journal of Nuclear Medicine*, 20(1), pp. 41–45. doi: 10.1967/s002449910505.

Lee, J.H. and Lee, S.W., 2017. The roles of carcinoembryonic antigen in liver metastasis and therapeutic approaches. *Gastroenterology research and practice*, 2017.

Li, J. et al. (2016) 'TNM Staging of Colorectal Cancer Should be Reconsidered According to Weighting of the T Stage', *Medicine (United States)*, 95(6), pp. 1–8. doi: 10.1097/MD.0000000000002711.

Li, X., An, B., Ma, J., He, B., Qi, J., Wang, W., Qin, C. and Zhao, Q., 2019. Prognostic value of the tumor size in resectable colorectal cancer with different primary locations: a retrospective study with the propensity score matching. *Journal of Cancer*, 10(2), p.313.

Liang, Y., Li, Q., He, D., Chen, Y. and Li, J., 2021. Tumor size improves the accuracy of the prognostic prediction of T4a stage colon cancer. *Scientific Reports*, 11(1), pp.1-11.

Lim, J.U., Lee, J.H., Kim, J.S., Hwang, Y.I., Kim, T.H., Lim, S.Y., Yoo, K.H., Jung, K.S., Kim, Y.K. and Rhee, C.K., 2017. Comparison of World Health Organization and Asia-Pacific body mass index classifications in COPD patients. *International journal of chronic obstructive pulmonary disease*, 12, p.2465.

Maeyama, Y., Mitsuyama, K., Noda, T., Nagata, S., Nagata, T., Yoshioka, S., Yoshida, H., Mukasa, M., Sumie, H., Kawano, H. and Akiba, J., 2018. Prediction of colorectal tumor grade and invasion depth through narrow-band imaging scoring. *World journal of gastroenterology*, 24(42), p.4809.

Martins, E. C. et al. (2019) 'Neutrophil-lymphocyte ratio in the early diagnosis of sepsis in an intensive care unit: A case-control study', *Revista Brasileira de Terapia Intensiva*, 31(1), pp. 63–70. doi: 10.5935/0103-507X.20190010.

Mo, C. ju et al. (2020) 'Diagnostic value of platelet-lymphocyte ratio and hemoglobinplatelet ratio in patients with rectal cancer', *Journal of Clinical Laboratory Analysis*, 34(4), pp. 1–7. doi: 10.1002/jcla.23153.

National Cancer Institute. n.d. *NCI Dictionary of Cancer Terms*. [online] Available at: <<https://www.cancer.gov/publications/dictionaries/cancer-terms/def/histologic-grade>> [Accessed 9 January 2022].

National Cancer Institute. n.d. *NCI Dictionary of Cancer Terms*. [online] Available at: <<https://www.cancer.gov/publications/dictionaries/cancer-terms/def/comorbidity>> [Accessed 9 January 2022].

ÖZGEHAN, G., Kahramanca, Ş., Kaya, I.O., Bilgen, K., Bostanci, H., Güzel, H., KÜÇÜKPINAR, T.H. and Kargici, H., 2014. Neutrophil-lymphocyte ratio as a predictive factor for tumor staging in colorectal cancer. *Turkish journal of medical sciences*, 44(3), pp.365-368.

Pedrazzani, C., Mantovani, G., Fernandes, E., Bagante, F., Salvagno, G.L., Surci, N., Campagnaro, T., Ruzzenente, A., Danese, E., Lippi, G. and Guglielmi, A., 2017. Assessment of neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio and platelet count as predictors of long-term outcome after R0 resection for colorectal cancer. *Scientific reports*, 7(1), pp.1-10.

Polat, E., Duman, U., Duman, M., Atici, A.E., Reyhan, E., Dalgic, T., Bostanci, E.B. and Yol, S., 2014. Diagnostic value of preoperative serum carcinoembryonic antigen and carbohydrate antigen 19-9 in colorectal cancer. *Current Oncology*, 21(1), pp.1-7.

Raehaan, N.R., Nurulita, A. and Arif, M., 2016. Carcinoembryonic antigen (CEA) di kanker kolorektal. *Indonesian Journal of Clinical Pathology and Medical Laboratory*, 20(3), pp.192-196.

Rafiyath, S., 2021. *Colon Cancer Staging: TNM Classification for Colon Cancer*. [online] Emedicine.medscape.com. Available at: <<https://emedicine.medscape.com/article/2006674-overview>> [Accessed 5 January 2022].

Rawla, P., Sunkara, T. and Barsouk, A. (2019) ‘Epidemiology of colorectal cancer: Incidence, mortality, survival, and risk factors’, *Przegląd Gastroenterologiczny*, 14(2), pp. 89–103. doi: 10.5114/pg.2018.81072.

Rosen RD, Sapra A. TNM Classification. [Updated 2020 Apr 16]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK553187/>

Shibutani, M. et al. (2013) ‘A high preoperative neutrophil-to-lymphocyte ratio is associated with poor survival in patients with colorectal cancer’, *Anticancer Research*, 33(8), pp. 3291–3294.

Siregar, G. A. and Sibarani, H. (2019) ‘Comparison of carcinoembryonic antigen levels among degree of differentiation and colorectal cancer’s location in Medan’, *Open Access Macedonian Journal of Medical Sciences*, 7(20), pp. 3447–3450. doi: 10.3889/oamjms.2019.442.

Somartini, J., Parewangi, A.L., Nu’man, A.S., Akil, F., Benyamin, F., Harjianti, T., Bakri, S., Kasim, H. and Seweng, A., 2019. Profile of CEA Levels in Colorectal Carcinoma patients: A Descriptive Analytic with Cross-Sectional Design. *International Journal of Medical Reviews and Case Reports*, 3(1), pp.5-9.

Stojkovic Lalosevic, M. et al. (2019) ‘Combined Diagnostic Efficacy of Neutrophil-to-Lymphocyte Ratio (NLR), Platelet-to-Lymphocyte Ratio (PLR), and Mean Platelet Volume (MPV) as Biomarkers of Systemic Inflammation in the

Diagnosis of Colorectal Cancer’, *Disease markers*, 2019, p. 6036979. doi: 10.1155/2019/6036979.

Su, B.B., Shi, H. and Wan, J., 2012. Role of serum carcinoembryonic antigen in the detection of colorectal cancer before and after surgical resection. *World journal of gastroenterology: WJG*, 18(17), p.2121.

The Global Cancer Observatory. 2020. Colon. [online] Available at:<<https://gco.iarc.fr/today/data/factsheets/cancers/8-Colon-fact-sheet.pdf>>

Tong, G. et al. (2018) ‘The role of tissue and serum carcinoembryonic antigen in stages I to III of colorectal cancer—A retrospective cohort study’, *Cancer Medicine*, 7(11), pp. 5327–5338. doi: 10.1002/cam4.1814.

Topdagi, O. and Timuroglu, A. (2018) ‘Evaluation of the relationship between carcinoembryonic antigen and TNM stage in colorectal cancer’, *Eurasian Journal of Medicine*, 50(2), pp. 96–98. doi: 10.5152/eurasianjmed.2018.17093.

Tsai, P. L. et al. (2016) ‘Neutrophil-lymphocyte ratio and CEA level as prognostic and predictive factors in colorectal cancer: A systematic review and meta-analysis’, *Journal of Cancer Research and Therapeutics*, 12(2), pp. 582–589. doi: 10.4103/09731482.144356.

White, A., Ironmonger, L., Steele, R.J., Ormiston-Smith, N., Crawford, C. and Seims, A., 2018. A review of sex-related differences in colorectal cancer incidence, screening uptake, routes to diagnosis, cancer stage and survival in the UK. *BMC cancer*, 18(1), pp.1-11.

Who.int. 2002. *WHO | Gender and human rights*. [online] Available at: https://www.who.int/reproductivehealth/topics/gender_rights/sexual_health/en/#:~:text=Sex%20refers%20to%20the%20biological,humans%20as%20males%20and%20females.> [Accessed 9 January 2022].

Wilson, M. J. et al. (2017) ‘Long-term prognostic value of preoperative anemia in patients with colorectal cancer: A systematic review and meta-analysis’, *Surgical Oncology*, 26(1), pp. 96–104. doi: 10.1016/j.suronc.2017.01.005.

Wilson, M. J. et al. (2017) ‘The role of preoperative iron deficiency in colorectal cancer patients: prevalence and treatment’, *International Journal of Colorectal Disease*, 32(11), pp. 1617–1624. doi: 10.1007/s00384-017-2898-1.

Wu, S. and Gu, W. (2020) ‘Association of T Stage and Serum CEA Levels in Determining Survival of Rectal Cancer’, *Frontiers in Medicine*, 6(January), pp. 1–7. doi: 10.3389/fmed.2019.00270.

wu, Y. et al. (2018) ‘Prediction of colorectal tumor grade and invasion depth through narrow-band imaging scoring’, *World Journal of Gastroenterology*, 24(42), pp. 4809–4820. doi: 10.3748/wjg.v24.i42.4809.

Xia, L. J. et al. (2020) ‘Significance of neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio, lymphocyte-to-monocyte ratio and prognostic nutritional index for predicting clinical outcomes in T1-2 rectal cancer’, *BMC Cancer*, 20(1), pp. 1–11. doi: 10.1186/s12885-020-6698-6.