UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Agati G, Azzarello E, Pollastri S, Tattini, M. 2012. Flavonoids as antioxidants in plants: location and functional significance. Plant Science Vol. 196: 67-76

Alén R. 2000. Structure and chemical composition of wood. In: Stenius P. (Ed.) Papermaking Science and Technology Page 11–57. Forest Products Chemistry Department. Helsinki.

Anouhe J S, Niamké F B, Faustin M, Virieux D, Pirat J, Adima A, Coulibaly S K, Amusant N. 2018. The role of extractives in the natural durability of the heartwood of *Dicorynia guianensis* Amsh: New insights in antioxydant and antifungal properties. Annals of Forest Science Vol. 75: 15-25

Aregay N, Belew D, Zenebe A, Haile M, Gebresamuel G, Girma A. 2021. Tree age and harvesting season affected physicochemical and bioactive compounds of elite type of gunda gundo orange (*Citrus* Spp) in the Northern Ethiopia. Fruit Science Vol. 20: 1-14

Aribowo T. 2016. Karakteristik sifat fisis dan keawetan kayu jati (*Tectona grandis* L.f.) cepat tumbuh terpadatkan. Skripsi Fakultas Kehutanan IPB. Bogor

Arief I I, Suryati T, Afiyah D N, Wardhani D P. Chemical and organoleptic of beef sausages with teak leaf extract (*Tectona grandis*) addition as preservative and Natural Dye. International Food Research Journal Vol. 21: 2033-2042

Arisandi R, Ashitani T, Takahashi K, Marsoem S N, Lukmandaru G. 2019. Lipophilic extractives of the wood and bark from (*Eucalyptus pellita*) F. Muell grown in Merauke, Indonesia. Wood Chemistry and Technology Vol. 40: 146-154

Ashwell G. 1957. Colorimetric analysis of sugars. Methods in Enzymology Vol. 3: 73–105

Ayadi N, Lejeune F, Charrier F, Charrier B, Merlin A. 2003. Color Stability of Heat-Treated During Artificial Weathering. Holz als Roh- und Werkstoff Vol. 61: 221–226.

Ayton J, Mailer R J, Haigh A, Tronson D, Conlan D. 2007. Quality and Oxidative Stability of Australian Olive Oil According to Harvest and Irrigation. Food Lipids Vol. 14: 138–156

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Baillères P H, Durand P Y. 2000. Non-destructive techniques for wood quality assessment of plantation-grown teak. Bois Forêts Tropica Vol. 263:17–29

Barbaroux C, Breda N. 2002. Contrasting distribution and seasonal dynamics of carbohydrate reserves in stem wood of adult ring-porous sessile oak and diffuse-porous beech trees. Tree Physiology Vol. 22: 1201–1210

Barbaroux C, Bréda N, Dufrêne E. 2003. Distribution of above-ground and below-ground carbohydrate reserves in adult trees of two contrasting broad-leaved species (*Quercus petraea*) and (*Fagus sylvatica*). New Phytologist Vol. 157: 605–615

Basri E, Wahyudi I. 2013. Sifat dasar kayu jati plus Perhutani dari berbagai umur dan kaitannya dengan sifat dan kualitas pengeringan. Jurnal Penelitian Hasil Hutan Vol. 31: 93-102.

Basri E, Yuniarti K, Wahyudi I, Damayanti R. 2015. Effects of girdling on wood properties and drying characteristics of Acacia mangium. Tropical Forest Science. 27: 498-505.

BeMiller J N. 1989.Carbohydrats. In: Rowe J.W. (eds) Natural products of woody plants. Springer Series in Wood Science. Berlin

Bertaud F, Holmbom B. 2004. Chemical composition of earlywood and latewood in Norway spruce heartwood, sapwood and transition zone wood. Wood Science Technology Vol. 38: 245–256

Bhat K M, Florence E J M. 2003. Natural Decay Resistance of Juvenile Teak Wood Grown in High Input Plantations. Holzforschung 57: 453-455

Bhat K M, Priya P B. 2004. Influence of provenance variation on wood properties of teak from the Western Ghat region in India. IAWA J Vol. 25:273–282

Bianchi S, Kroslakova I, Janzon R, Mayer I, Saake B, Pichelin F. 2015. Characterization of condensed tannins and carbohydrates in hot water bark extracts of European softwood species. Phytochemistry Vol. 120: 53-61

Blainski A, Lopes G C, Mello J C P. 2013. Application and Analysis of the Folin Ciocalteu Method for the Determination of the Total Phenolic Content from *Limonium Brasiliense* L. Molecules Vol. 18: 6852-6865.

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Bowyer J L, Shmulsky R, Haygreen J G. 2007. Forest product and wood science an introduction fifth edition. Blackwell Publishing Professional. Iowa.

Brown H P, Panshin A J, Forsaith C C. 1994. Textbook of wood technology. Volume I. McGraw-Hill Book Company Inc. 4th Edition. New York

Burtin P J C, Charpentier J P, Janin G. 1998. Natural wood coloring process in *Juglans* sp (*J nigra*, *J regia* and *J nigra* 23 x *J regia*) depends on native phenolic compounds accumulated in the transition zone between sapwood and heartwood. Trees Vol. 12: 258-264

Bustan A, Avni A, Lavee, S, Zipori I, Yeselson Y, Schaffer A A, Riov Joseph, Dag A. 2011. Role of carbohydrate reserves in yield production of intensively cultivated oil olive *(Olea europaea* L.) trees. Tree Physiology Vol. 31: 519–530

Caron A, Clemens M A, Barry G, Michael C J. 2013. Distribution of extractives in sitka spruce (*Picea sitchensis*) grown in the northern UK. Wood Product Vol. 71: 697–704

Chagas S F, Evangelista W V, Silva J C, Carvalho A M M L. 2014 Propriedades da madeira de teca de diferentes idades e oriundas de desbaste. Ciência da Madeira Vol. 5: 138-150

Chapin F S, Schulze E, Mooney H A. 1990. The ecology and economics of storage in plants. Annual Review of Ecology and Systematics Vol. 21: 423–447

Chen C M. 1970. Effect of extractive removal on adhesion and wettability of some tropical woods. Forest Product Vol. 20: 36–41

Chikezie P C, Ibegbulem C O, Mbagwu F N. 2015. Bioactive principles from medicinal plants. Phytochemistry Vol. 9: 88–115.

Choi H R, Choi J S, Han Y N, Bae S J, Chung H.Y. 2002. Peroxynitrite scavenging activity of herb extracts. Phytother Res Vol. 16: 364-367.

Chow P S, Landhausser S M. 2004. A method for routine measurements of total sugar and starch content in woody plant tissues. Tree Physiology Vol. 24: 1129 –1136

Chow P, Nakayama F S, Blahnik B, Youngquist J A, Coffelt T A. 2008. Chemical constituents and physical properties of guayule wood and bark. Industrial Crops and Products Vol. 28: 303–308

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Connor A M, Chad E F, Peter A A. 2005. Genotypic and enviromental variation in antioxidant activity and total phenolic content among blackbeery and hybridberry cultivars. American Society of Horticultural Science. Vol. 130: 527–533

Croft K D. 1998. The chemistry and biological effects of flavonoids and phenolic acids. Annals of The New York Academi of Science Vol. 854: 435-442.

Da Costa E W B, Rudman F.J G. 1958. Investigations on the durability of *Tectona grandis*. Empire Forestry Review Vol. 37: 291-298.

Damayanti R. 2010. Struktur makro, mikro, dan ultramikroskopik kayu Jati Unggul Nusantara dan kayu jati konvensional. Tesis Fakultas Kehutanan Institut Pertanian Bogor. Bogor

Damayanti R, Ozarska B, Pandit I K N, Febrianto F, Gustan P. 2018. Wood properties of 5-years-old fast grown teak. Wood research Vol. 9: 29-34

Damesin C, Lelarge C. 2003. Carbon isotope composition of current-year shoots from (*Fagus sylvatica*) in relation to growth, respiration and use of reserves. Plant Cell and Environtment Vol. 26: 207–219

Dedrie M, Jacquet N, Bombeck P, Hébert J, Richel A. 2015. Oak barks as raw materials for the extraction of polyphenols for the chemical and pharmaceutical sectors: A regional case study. Industrial Crops and Products Vol. 70: 316–321

De la Rosa L A, Moreno-Escamilla J O, Rodrigo-García J, Alvarez-Parrilla E. 2019. Phenolic compounds. Postharvest Physiology and Biochemistry of Fruits and Vegetables Vol. 2: 253–271

Del Rio D, Rodriguez-Mateos A, Spencer J P E, Tognolini M, Borges G A. 2013. Dietary (Poly) phenolics in human health: structures, bioavailability, and evidence of protective effects against chronic diseases. Antioxidants and Redox Signaling Vol. 18: 1818–1892

Domingues R M A, Sousa G D A, Freire C S R, Silvestre A J D, C Pascoal Neto 2010. *Eucalyptus globulus* biomass residues from pulping industry as a source of high value triterpenic compounds. Industrial Crops and Products Vol. 31: 65–70

Donnelly D M X. 1975 Neoflavonoids. In: Harborne B, Mabry T J, Mabry H (eds) The flavonoids. Chapman and Hall. London

Dróżdż P, Pyrzynska K. 2018. Assessment of polyphenol content and antioxidant activity of oak bark extracts. European Journal of Wood and Wood Products Vol. 76: 793–795

Dubois M, Gilles K A, Hamilton J K, Rebers P A, Smith F. 1956. Colorimetric method for determination of sugars and related substances. Anal Chem 28:350–356

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Dunisch O, Richter H G, Koch G. 2010. Wood properties of juvenile and mature heartwood in *Robinia pseudoacacia* L. Wood Science and Technology Vol. 44: 301-313.

Efansyah M N, Djofrie M H B, Limbong W H. 2011. Prospek usaha bagi hasil penanaman Jati Unggul Nusantara (studi kasus Koperasi Wanabhakti Nusantara di Kabupaten Bogor). Tesis Fakultas Kehutanan IPB. Bogor

Ekman R, Holmbom B. 2000. Thechemistry of wood resin. In: Back E L, Allen L H (Eds.) PitchControl Wood Resin and Deresination. GA: Tappi Press. Atlanta

Fatimah S, Susannto M, Lukmandaru G. 2013. Studi komponen kimia kayu *Eucalyptus pellita* F. Muell dari pohon plus hasil uji keturunan generasi kedua di Wonogiri Jawa Tengah. Jurnal Ilmu Kehutanan. Vol. 7: 1-13

Fengel D, Wegener G. 1989. Kayu: Kimia, ultrastuktur, reaksi-reaksi. Gadjah Mada University Press (diterjemahkan oleh Sastrohamidjojo, H.). Yogyakarta.

Ferreira J P A, Miranda I, Gominho J, Pereira H. 2015. Selective fractioning of *Pseudotsuga menziesii* bark and chemical characterization in view of an integrated valorization. Industrial Crops and Products Vol. 74: 998–1007

Fitri A, Toharmat T, Astuti D A, Tamura H. 2015. The Potential use of secondary metabolites in *Moringa oleifera* as an antioxidant source. Media Peternakan Vol. 38: 169-175

Gan R Y, Chan C L, Yang Q Q, Li H B, Zhang D, Ge Y Y, Corke H. 2019. Bioactive compounds and beneficial functions of sprouted grains. Sprouted Grains Vol. 1: 191–246

Gao H, Shupe T F, Eberhardt T, Hse C Y. 2006. Antioxidant activity of extracts from the wood and bark of Port-Orford cedar. Wood Science Vol. 53: 147-152

Genet H, Breda N, Dufrene E. 2010. Age-related variation in carbon allocation at tree and stand scales in beech (*Fagus sylvatica* L.) and sessile oak (*Quercus petraea* (Matt.) Liebl.) using a chronosequence approach. Tree Physiology Vol. 30: 177–192

Gominho J, Figuera J, Rodrigues C J. 2001. Within tree variations of heartwood, extracives and wood density in the eucalyptus hibrid urograndis. Wood and Fiber Science Vol. 33: 3-8

Gruber A, Pirkebner D, Oberhuber W, Wieser G. 2011. Spatial and seasonal variations in mobile carbohydrates in *Pinus cembrain* the timberline ecotone of the Central Austrian Alps. European Journal of Forest Research Vol. 130: 173–179

Gust J, Suwalski J. 1994. Use of mössbauer spectroscopy to study reaction products of

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

polyphenols and iron compounds. Corrosion Vol. 50: 355–365

Hall M, Hansen N W, Rudra A B. 1973. The effect of species, age and wood characteristics on eucalypt kraft pulp quality. Appita Vol. 26: 348 – 354

Harborne J B. 1980. Secondary plant products: encyclopedia of plant physiology. New Series Volume 8. Phytochemistry Vol. 19: 2803–2804

Hart J H, Shrimpton D M. 1979. Role of stilbenes in resistance of wood to decay. Phytopathology Vol. 69: 1138-1143

Hart J H. 1989. The Role of wood Exudate and extractives in protecting wood from decay. In: Rowe J.W. (eds) Natural products of woody plants. Springer Series in Wood Science. Berlin

Hata K, Sogo M. 1952. The difficulty of sulfite digestion of pine heartwood. Japan Forest Society Vol. 35: 74-77

Haupt M, Leithoff H, Meier D, Puls J, Richter H G, Faix O. 2003. Heartwood extractives and natural durability of plantation-grown teakwood (*Tectona grandis* L.) – a case study. Holz Roh Werkst Vol. 61: 473–474

Heim K E, Tagliaferro A R, Bobilya D J. 2002. Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. Nutrition Biochemistry Vol. 13: 572-584.

Hidayati F, Marsoem S N. 2010. Anatomi dan sifat fisika kayu jati unggul (*Tectona grandis* L.f.) umur 5 tahun yang tumbuh di Gunung Kidul pada berbagai laju pertumbuhan. Tesis Fakultas Kehutanan Universitas Gadjah Mada. Yogyakarta

Hidayati F, Fajrin I T, Ridho M R, Nugroho W D, Marsoem S N, Na'iem M. 2016. Sifat fisika dan mekanika kayu jati unggul mega dan kayu jati konvensional yang ditanam di Hutan Pendidikan Wanagama, Gunungkidul, Yogyakarta. Jurnal Ilmu Kehutanan Vol. 10: 98-107

Higgins H G. 1984. Pulp and paper. In: Hillis W E, Brown A G (eds) Eucalypts for wood production. CSIRO Academic Press. New York

Hillis W E. 1971. Distribution, properties and formation of Ssome wood extractives. Wood Science and Technology Vol. 5: 272—289

Hillis W E. 1987. Heartwood and tree exudates. Springer. New York

Hillis W E, Sumimoto M. 1989. Effect of extractives on pulp. In: Rowe J W (eds) Natural products of woody plants. Springer Series in Wood Science. Berlin

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Hillis W E, Hasegawa M. 1963. The formation of polyphenols in trees I. Administration of c glucose and subsequent distribution of radioactivity. Phytochemistry Vol. 2: 195–199

Howard E T. 1972. Physical and chemical properties of slash pine tree parts. Wood Science Vol. 5: 312 – 317

Hoch G, Richter A, Korner C. 2003. Non-structural carbon compounds in temperate forest trees. Plant, Cell and Environment Vol. 26: 1067–1081

Hse C, Kuo M. 1988. Influence of extractives on wood gluing and finishing. Forest Products Journal Vol. 38: 52-56

Ihda F V. 2019. Kadar ekstraktif dan sifat warna dari jenis permudaan jati. Skripsi Fakultas Kehutanan UGM. Yogyakarta

Irawati D, Azwar N R, Syafii W, Artika I M. 2009. Pemanfaatan serbuk kayu untuk produksi etanol dengan perlakuan pendahuluan delignifikasi menggunakan jamur *Phanerochaete chrysosporium*. Jurnal Ilmu Kehutanan Vol. 3: 13-22.

Kabtni S, Sdouga D, Bettaib R I. 2020. Influence of climate variation on phenolic composition and antioxidant capacity of Medicago minima populations. Scientific Report Vol. 10**:** 82-93

Kadambi K. 1972. Silviculture and management of teak. Stephen F Austin State University. Texas

Kampe A, Magel E. 2013. New insights into heartwood and heartwood formation: Cellular aspects of wood formation. Springer. Berlin

Kannan S, Kelly R G. 1996. The role of dihydroxybenzenes and oxygen on the corrosion of steel in black liquor. Corrosion Science Vol. 38: 1051–1069

Kaosa-ard A. 1981. Teak (*Tectona grandis* Linn. f)- its natural distribution and related factors. Natural His Siam Society Vol. 29: 55-74.

Kaškonienė V, Maruška A, Kornyšova O, Charczun N, Ligor M, Buszewski B. 2009. Quantitave and qualitative determination of phenolic compound in honey. Chemine Technologija Vol. 3: 74-80

Kininmonth J A. 1986. Wood from fast-grown, short-rotation trees. Proceedings 18th IUFRO World Congress. Division 5 Forest Products. Ljubljana.

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Klein T, Hoch G, Yakir D, Korner C. 2014. Drought stress, growth and nonstructural carbohydrate dynamics of pine trees in a semi-arid forest. Tree Physiology Vol. 34: 981–992

Kokutse A D, Stokes A, Bailleres H, Kokou K, Baudasse C. 2006. Decay resistance of Togolese teak (*Tectona grandis* L.) heartwood and relationship with colour. Trees Vol. 20: 219–223.

Kollert W, Kleine M. 2017. The global teak study: Analysis, evaluation and future potential of teak resources. International Union of Forestry Organizations. IUFRO World Series Volume 36. Viena

Korner C. 2003. Carbon limitation in trees. Journal of Ecology Vol. 91: 4–17

ozlowski T T. 1992. Carbohydrate sources and sinks in woody plants. The Botanical Review Vol. 58: 107–222

Krisdianto, Sumarni G. 2006. Perbandingan persentase volume teras kayu jati cepat tumbuh dan konvensional umur 7 tahun asal Penajam, Kalimantan Timur. Jurnal Penelitian Hasil Hutan Vol. 24: 385-394

Król A, Amarowicz R, Weidner S. 2014. Changes in the composition of phenolic compounds and antioxidant properties of grapevine roots and leaves (*Vitis vinifera* L.) under continuous of long-term drought stress. Acta Physiologiae Plantarum Vol. 36: 1491–1499

Kubczak M, Khassenova A B, Skalski B, Michlewska S, Wielanek M, Aralbayeva A. N, Murzahkmetova M K, Zamaraeva M, Sklodowska M, Bryszewska M I M. 2020. Bioactive compounds and antiradical activity of the *Rosa canina* L. leaf and twig extracts. Agronomy Vol. 10: 1-20

Ku C S, Jang J P, Mun S P. 2007. Exploitation of polyphenol-rich pine barks for antioxidant activity. Wood Science Vol. 53: 524-528.

Labosky P. 1979. Chemical constituents of four southern pine parks. Wood Science 12: 80–85.

Lachowicz H, Hanna W, Rafal W, Magdalena S. 2019. The effect of tree age on the chemical composition of the wood of Silver Berch (*Betula pendula* Roth) in Poland. Wood Science and Technology Vol. 53: 1135–1155

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Laks P E. 1991. Chemistry of bark. In: Hon, D N S, Shiraishi N (ed), Wood and cellulosic chemistry. NY: Marcel Dekker Inc. New York

Lamounier K C, Cunha L C S, de Morais S A L, de Aquino F J T, Chang R E, do Nascimento R T, de Souza M G M, Martins C H G, Cunha W R. 2012. Chemical analysis and atudy of phenolics, antioxidant activity, and antibacterial effect of the wood and bark of *Maclura tinctoria* (L.) D. Don ex Steud. Evidence-based Complementary and Alternative Medicine Vol. 2012: 1-8

Lattanzio V, Kroon P A, Quidheau S, Treutter D. 2008. Recent advances in polyphenol research plant phenolics secondary metabolites with diverse functions. Blackwell Publishing. Foggia

Lelis R. 1992. Zum verhalten von splint-und kernholz der kiefer bei der verleimung mit harnstofformaldehydharzen (UF-harzen) und diisocyanat-klebstoffen (PMDI). Magisterarbeit an der Forstlichen Fakultät der Georg-August-Universität Göttingen. Göttingen

Le Normand M, Ulrica E, Holmbom B, Monica E K. 2012. Hot-water extraction and characterization of spruce bark non-cellulosic polysaccharides. Nordic Pulp and Paper Research Journal Vol. 27: 18–023

Li-Paul H. 1989. Low temperature stress physiology in crops. CRC Press Inc. Florida

Lloret F, Sapes G, Rosas T, Galiano L, Saura-Mas S, Sala A, Martínez-Vilalta J. 2018. Non-structural carbohydrate dynamics associated with drought-induced die-off in woody species of a shrubland community. Annals of Botany Vol. 121: 1383–1396

Lukmandaru G, Takahashi K. 2008 Variation in the natural termite resistance of teak (*Tectona grandis* Linn.fil.) wood as a function of tree age. Annals of Forest Science Vol. 65: 708

Lukmandaru G. 2009a. Pengukuran kadar ekstraktif dan sifat warna pada kayu teras jati doreng (*Tectona grandis*). Jurnal Ilmu Kehutanan Vol. 3: 67-73

Lukmandaru G. 2009b. Sifat kimia dan warna kayu teras jati pada tiga umur berbeda. Tropical Wood Science and Technology Vol. 7: 1-8

Lukmandaru G, Takahashi K. 2009. Radial distribution of quinone in plantation teak (*Tectona grandis* L.f.). Annals of Forest Science Vol. 66: 605

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Lukmandaru G. 2011. Variability in the natural termite resistance of plantation teak wood and its relations with wood extractive content and color properties. Indonesian Journal of Forestry Research Vol. 8:17–31

Lukmandaru G. 2013. The natural resitance of teak wood grown in community forest. Ilmu dan Teknologi Kayu Tropis Vol. 11: 131-139

Lukmandaru G. 2016. Hubungan kadar ekstraktif dengan sifat warna kayu teras jati. Jurnal Penelitian Hasil Hutan Vol. 34: 207 - 216.

Lukmandaru G. 2015. Antitermitic activiy of the bark extracs of teak. Proceeding of The Fifth International Symposium of Idonesia Wood Research Society.

Luo Z B, Calfapietra C, Liberloo M, Scarascia-mugnozza G, Polle A. 2006. Carbon partitioning to mobile and structural fractions in poplar wood under elevated CO2 (Euroface) and N fertilization. Global Change Biology Vol. 12: 272–283

Magel E, Einig W, Hampp, R. 2000. Carbohydrates in trees in carbohydrate reserves in plants - Synthesis and regulation**.** Elsivier Science. Ludhiana

Martawijaya A, Kartasujana I, Kadir K, Prawira S.A. 2005. Atlas kayu Indonesia jilid I. Badan Penelitian dan Pengembangan Kehutanan. CV. Miranti. Bogor

Masendra, Ashitani T, Takahashi K, Susanto Mudji, Lukmandaru, G. 2019. Hydrophilic extracts of the bark from six pines species. Korean Wood Science Technology Vol. 47: 80-89

Matangaran J R, Romadoni A. 2012. Limbah pemanenan jati di Bayuwangi Jawa Timur. Perennial Vol. 8: 88.

Maulida F, Meiganati K B, Maslahat M. 2020. Komponen kimia kayu trubusan Jati Unggul Nusantara (JUN) pada bagian pangkal, tengah, ujung. Sains Natural UNB Vol. 10: 55-60

Mawardi P. 2012. Kaya dari investasi jati barokah. PT Agro Media Pustaka. Jakarta

McDonald M, Mila I, Scalbert A. 1996. Precipitation of metal ions by plant polyphenols: optimal conditions and origin of precipitation. Agriculture Food Chemistry Vol. 44: 599-606

McGinnis G D, Parikh, S. 1975. The chemical constituents of loblolly pine. Wood Science Vol. 7: 295–297

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Meiganati K B. 2017. Riap pertumbuhan jati unggul nusantara rotasi kedua di kebun percobaan Cogreg Universitas Nusa Bangsa. Nusa Sylvae Vol. 17: 101 - 105

Metsamuuronen S, Siren H. 2019. Bioactive phenolic compounds, metabolism and properties: a review on valuable chemical compounds in Scots pine and Norway spruce. Phytochem Review Vol. 18: 623-664

Midgley S J, Stevens P R, Arnold R J. 2017. Hidden assets: Asia's smallholder wood resources and their contribution to supply chains of commercial wood. Australian Forestry Vol. 80: 10–25

Migita N Y, Yonezawa, Kondo T. 1968. Wood chemistry (in Japanese). Kyoritsu Shupan. Tokyo

Miranda I, Souca V, Pereira H. 2011. Wood properties of teak (*Tectona grandis*) from a mature unmanaged stand in East Timor. Wood Science Vol. 57: 171-178

Miranda I, Pereira H. 2002. The variation of chemical composition and pulping yield with age and growth factors in young *Eucalyptus globulus*. Wood Fiber Science Vol. 34:140–145

Morais M C, Helena P. 2012. Variation of extractives content in heartwood and sapwood of *Eucalyptus globulus* trees. Wood Science and Technology Vol. 46:709–719.

Mota G S, Sartori C J, Miranda I, Quilhó T, Mori F A, Pereira H. 2017. Bark anatomy, chemical composition and ethanol-water extract composition of *Anadenanthera peregrina* and *Anadenanthera colubrina*. Plos One Vol. 12: 1-14

Moya R, Ledezma V A. 2003. Effect of plantation spacing on physical properties of teakwood along the stem (in Spanish). Madera Bosques Vol. 9: 15–27

Narayanamurthi D, George J, Pant H C, Singh J. 1962. Extractives in teak. Sylvae geneticae Vol. 11: 57-63.

Negi A, Nimisha S, Reetika P, Singh M F. 2012. Determination of total phenolic content of stem the stem bark of *Bauhinia variegata* Linn; an Approach to Standardization. Pharma Research Vol. 7: 16 - 22

Neiva D M, Araujo S, Gominhoa J, Carneiro A D C, Pereira H. 2018. Potential of *Eucalyptus Globulus* industrial bark as a biorefinery feedstock: Chemical and fuel characterization. Industrial Crops and Products Vol. 123: 262–270

Newell E A, Mulkey S S, Wright J S. 2002. Seasonal patterns of carbohydrate storage

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

in four tropical tree species. Oecologia Vol. 131: 333–342

Niamké B F, Amusant N, Kokutse A D, Chaix G, Charpentier J-P, Adima A A, Kati Coulibaly S, Jay-Allemand C. 2010. Radial distribution of non-structural carbohydrates in Malaysian teak. Biological and Chemical Science Vol. 4: 710-720

Niamké B F, Amusant N, Charpentier J-P, Chaix G, Baissac Y, Boutahar N, Adima A A, Kati-Coulibaly S, Jay-Allemand C. 2011. Relationships between biochemical attributes (non-structural carbohydrates and phenolics) and natural durability against fungi in dry teak wood (*Tectona grandis* L.). Annals Forest Science Vol. 68: 201-211

Niamke B F, Adima A A, Seraphin K, Amusant N, Jay-Allemand C. 2018. Heartwood formation process in teak (*Tectona grandis* L): fate of non-structural carbohydrates and characterization of forsythoside B. International Journal of Biological and Chemical Sciences Vol. 12: 1102-1112

Niinemets Ü. 2010. Responses of forest trees to single and multiple environmental stresses from seedlings to mature plants: Past stress history, stress interactions, tolerance and acclimation. Forest Ecology and Management Vol. 260: 1623–1639

Nisula L. 2018. Wood extractives in conifer : A study of stemwood and knots of indutrially important species. Abo Academy Unviversity Press. Tavastgatan

Noel J P, Austin M B, Bomati E K. 2005. Structure–function relationships in plant phenylpropanoid biosynthesis. Plant Biology Vol. 8: 249–253

Noreen H, Semmar N, Farman M, McCullagh J S O. 2017. Measurement of total phenolic content and antioxidant activity of aerial parts of medicinal plant *Coronopus didymus*. Asian Pacific Journal of Tropical Medicine Vol. 10: 792-801

Oberhuber W, Swidrak I, Pirkebner D, Gruber A. 2011. Temporal dynamics of nonstructural carbohydrates and xylem growth in (*Pinus sylvestris*) exposed to drought. Canadian Journal of Forest Research Vol. 41: 1590–1597

Panshin A.J, Zeeuw C D, Brown H P. 1964. Textbook of wood technology. Volume I-structure, identification, uses, and properties of the commercial woods of the United States. McGraw-Hill Book Company.

Payá M, Halliwell B, Hoult J R S. 1992. Interactions of a series of coumarins with reactive oxygen species. Scavenging of superoxide, hypochlorous acid and

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

hydroxyl radicals. Biochemistry and Pharmacology Vol. 44: 205-214

Pereira D M, Patricia V, Jose A P, Paula B A. 2009. Fenolics : From chemistry to biology. Molecules Vol. 14: 2202-2211

Pereira H. 1986. Variability in the chemical composition of plantation eucalypts (*Eucalyptus globulus*). Wood and Fiber Science Vol. 20: 82 - 90

Pérez L D, Kanninen M. 2003. Heartwood, sapwood and bark content, and wood dry density of young and mature teak (*Tectona grandis*) trees grown in Costa Rica. Silva Fenn Vol. 37: 45–54

Pérez L D, Kanninen M. 2005. Effect of thinning on stem form and wood characteristics of teak (*Tectona grandis*) in a humid tropical site in Costa Rica. Silva Fenn Vol. 39: 217–225

Piper F I, Fajardo A. 2011. No evidence of carbon limitation with tree age and height in *Nothofagus pumilio* under Mediterranean and temperate climate conditions. Annals of Botany Vol. 108(5), 907–917

Piper F I. 2020. Decoupling between growth rate and storage remobilisation in broadleaf temperate tree species. Functional Ecology Vol.34: 1180-1192

Plavcová L, Jansen S. 2015. The role of xylem parenchyma in the storage and utilization of nonstructural carbohydrates: Functional and Ecological Xylem Anatomy. Springer. Cham

Posch B, Wegener G, Grosser D, Wagner L. 2004. Physikalische und mechanische Untersuchungen an Teakholz (*Tectona grandis* L.f.) aus Plantagen in Panama. Holz Roh Werkst Vol. 63: 31–35

Pratiwi D D. 2018. Komposisi kimia ekstraktif kulit luar dan dalam kayu *Araucaria cunninghamii, Araucaria papuana, Agathis loranthifolia* dan *Podocarpus imbricartus.* Skripsi Fakultas Kehutanan UGM. Yogyakarta

Purba B A V. 2019. Kandungan dan aktivitas antioksidan ekstraktif kayu tiga klon hibrid akasia (*Acacia mangium x Acacia auriculiformis*). Skripsi Fakultas Kehutanan UGM. Yogyakarta

Putri K. 2009. Pengelolaan hutan berbasis masyarakat tersertifikasi di Kabupaten Konawe Selatan , Sulawesi Tenggara. Skripsi Fakultas Kehutanan IPB. Bogor

Putro G S, Marsoem S N, Sulistyo J, Hadiwinoto S. 2020. Sifat kayu jati unggul nusantara (*Tectona grandis* L.f) pada tiga kelas diameter pohon. Pemuliaan

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Tanaman Hutan Vol 14: 9 - 18.

Qiu H, Ru L, Ling L. 2019. Analysis of chemical composition by aceton and the chromatic aberration of teak (*Tectona grandis* L.f) from China. Molecules Vol. 24: 1-11

Reddy C V K, Sreeramulu D, Raghunath M. 2010. Antioxidant activity of fresh and dry fruits commonly consumed in India. Food Research International Vol. 43: 285–288.

Richardson A D, Carbone M S, Keenan T F, Czimczik C I, Hollinger D Y, Murakami P, Schaberg P G, Xu X. 2013. Seasonal dynamics and age of stemwood nonstructural carbohydrates in temperate forest trees. New Phytologist Vol. 197: 850–861

Rizanti D E, Wayan D, Beatrice G, Andre M, Stephane D, Hubert C, Christiane G, Eric Gelhaye, Phila R, Rita K S, Syafii W, Rozi M, Philippe G. 2018. Comparison of teak wood properties according to forest management: Short versus long rotation. Annals of Forest Science Vol. 75: 39

Roffael E. 2016. Significance of wood extractives for wood bonding. Applied Microbiol Biotechnology Vol. 100: 1589–1596

Rosales-Castro M, Honorato-Salazar J A, Reyes-Navarrete M. G, González-Laredo R F. 2015. Antioxidant phenolic compounds of ethanolic and aqueous extracts from pink cedar (*Acrocarpus fraxinifolius*) whight & arn bark at two tree ages. Wood Chemistry and Technology Vol. 35: 270–279

Rosell J A. 2019. Bark in woody plants: understanding the diversity of a multifunctional structure. National Laboratory of Sustainability Sciences, Institute of Ecology University Nacional Autónoma de México. Mexico

Rosell J A, Piper F I, Jiménez-Vera C, Vergílio P C B, Marcati C R, Castorena M, Olson M. E. 2020. Inner bark as a crucial tissue for non-structural carbohydrate storage across three tropical woody plant communities. Plant, Cell & Environment Vol. 44: 156-170

Rudman P, Da Costa EWB. 1959. Variation in extractive content and decay resistance of *Tectona grandis*. Inst Wood Science Vol. 3: 33–42

Routa J, Perttu A, Antti A. 2017. Wood extractives of finnish pine, spruce and birch – availability and optimal sources of ompounds: A Literature Review. Juvenes. Helshinki.

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Rowell R M, Roger P, Mandla A T. 2012. Wood chemistry and wood composites. CRC Press. London

Sakai K. 2001. Chemistry of bark in wood and cellulosic chemistry: Second edition, revised and expanded. Marcel Decker Inc. New York

Sala A, Hoch G. 2009. Height-related growth declines in ponderosa pine are not due to carbon limitation. Plant, Cell and Environtment Vol. 32: 22–30

Sandermann W, Dehn U V. 1951. Einfluß chemischer Faktoren auf die Festigkeitseigenschaften zementgebundener Holzwolleplatten. Holz Roh Werkst Vol. 9: 97–101

Santana W M S, Natalino C, Arantes M D C, Trugilho P F. 2012. Effect of age and diameter class on the properties of wood from clonal eucalyptus. Cerne, Lavras Vol. 18: 1-8

Sarawong C, Schoenlechner R, Sekiguchi K, Berghofer E, Nguyen P K W. 2014. Effect of extrusion cooking on the physicochemical properties, resistant starch, phenolic content and antioxidant capacities of green banana flour. Food Chemistry Vol. 143: 33–39

Savage J. A, Clearwater M. J, Haines D. F, Klein T, Mencuccini, M, Sevanto S, Turgeon Robert, Zhang C. 2016. Allocation, stress tolerance and carbon transport in plants: how does phloem physiology affect plant ecology? Plant, Cell & Environment Vol. 39: 709–725

Scheffer T C, Cowling E B. 1966. Natural resistance of wood to microbial deterioration. Annal Review Phytopathology Vol. 4: 147-170

Schofield P, Mbugua D M, Pell A N. 2001. Analysis of condensed tannins: A review. Anim. Feed ScienceTechnology Vol. 91: 21–40

Silverio F O, Barbisa L C A, Maltha C R A, Silvestre A J D, Veloso D P, Gomide J L. 2007. Lipophilic Extractives of *E. urogandis*. Bioresources Vol. 2(2): 157-168.

Simard S, Giovannelli A, Treydte K, Traversi M L, King G M, Frank D, Fonti P. 2013. Intra-annual dynamics of non-structural carbohydrates in the cambium of mature conifer trees reflects radial growth demands. Tree Physiology Vol. 33: 913-923

Singh R, Singh S, Kumar S, Arora S. 2007. Studies on antioxidant potential of methanol extract/fractions of Acacia auriculiformis A. Cunn. Food Chemistry Vol. 103: 505–511

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Singleton V. L, Orthofer R, Lamuela-Raventós R. M. 1999. Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. Methods in Enzymology Vol. 299: 152-178

Sjostrom E. 1995. Kimia kayu, dasar-dasar dan penggunaan edisi kedua. Diterjemaahkan oleh Sastrohamidjojo, H. Terjemaahan dari: Wood chemistry, fundamentals and application second edition. Gadjah Mada University Press. Yogyakarta.

Someya S, Yoshiki Y, Okubo K. 2002. Antioxidant compounds from bananas (Musa Cavendish). Food Chemistry Vol. 79: 351–354

Street R J. 1962. Exotic forest trees in the British Commonwealth. Clarendon Press. Oxford

Sturgeon R J. 1990. Monosaccharides. In methods in plant biochemistry. Carbohydrates. Ed. P.M. Dey. Academic Press. London

Sukardi, Mulyarto A R, Safera W. 2007. Optimasi waktu ekstraksi terhadap kandungan tanin pada bubuk ekstrak daun jambu biji (*Psidium folium*) serta biaya produksinya. Jurnal Teknologi Pertanian Vol. 8: 88 - 94

Sumarna Y. 2003. Budi daya jati. Penebar Swadaya. Jakarta

Supriono B, Setyaningsih L. 2012. Pertumbuhan Tanaman Jati Unggul Nusantara dengan Pola Agroforestry Umur Lima Tahun. Jurnal Sains Natural Universitas Nusa Bangsa Vol. 2: 179-185

Susrusa I K B, Ardhana I P G. 2007. Kelayakan finansial budidaya tanaman jati (*Tectona grandis* Linn) varietas unggul di Kabupaten Buleleng. Sosial Ekonomi Pertanian Vol. 10: 189-198

Swan B. 1968. Seasonal variations in the extractives of spruce wood and sulfite pulps. Sven Papperstidn Vol. 71: 436–440.

Swason J W, Cordingly. 1959. Surface chemical studies on pitch. 11. The mechanism of the loss of absorbancy and development of self-sizing in paper made from wood pulp. TAPPI Vol. 42: 812-819

Syafii W. 2000. The basic properties of Indonesia teakwood at various age classes. Proceeding of the Third International Wood Science Symposium. Kyoto

Taylor A, Cooper P. 2002. The effect of stem girdling on wood quality. Wood and Fiber Science Vol. 34: 212-220.

Thulasidas PK, Bhat KM 2007 Chemical extractive compounds determining the

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

brown-rot decay resistance of teak wood. Holz Roh Werkst Vol. 65: 121–124

Umezawa T. 2001. Chemistry of extractives in wood and cellulosic chemistry: Second edition, revised and expanded. Marcel Decker Inc. New York

Valentão P, Fernandes E, Carvalho F, Andrade P.B, Seabra R.M, Bastos M.L. 2002. Antioxidative properties of cardoon (*Cynara cardunculus* L.) infusion against superoxide radical, hydroxyl radical and hypochlorous acid. Agriculture Food Chemistry Vol. 50: 4989-4993

Valentão P, Fernandes E, Carvalho F, Andrade P.B, Seabra R.M, Bastos M.L. 2003. Hydroxyl radical and hypochlorous acid scavenging activity of small centaury (*Centaurium erythraea*) infusion. A comparative study with green tea (*Camellia sinensis*). Phytomedicine Vol. 10: 517- 522.

Van-Sumere C.F. 1989. Phenols and phenolic acids. In: Methods in Plant Biochemistry, Vol. 1. Plant Phenolics (eds P.M. Dey & J.B. Harborne). Academic Press. London.

Vázquez-León L. A, Páramo-Calderón D E, Robles-Olvera V J, Valdés-Rodríguez O A, Pérez-Vázquez A, García-Alvarado M. A, Rodríguez-Jimenes G. C. 2017. Variation in bioactive compounds and antiradical activity of *Moringa oleifera* leaves: influence of climatic factors, tree age, and soil parameters. European Food Research and Technology Vol. 243: 1593-1608

Vermerris W, Nicholson R. 2006. Chemical Properties of Phenolic Compounds. Phenolic Compound Biochemistry. Springer. Florida

Villagomez H Z, Peterson D M, Herrin L, Young R A. 2005. Antioxidant activity of different component of pine species. Holzforschung, Vol. 59: 156–162

Vora D.J.D, 2017. Insight into the biochemical link between biodiversity and nutraceuticals. IOSR Journal of Environmental Science, Toxicology and Food Technology Vol. 11: 22–25

Vuolo M. M, Lima V. S, Maróstica Junior M. R. 2019. Phenolic compounds: structure, classification, and antioxidant power. bioactive compounds Vol. 1: 33–50

Wahyudi I, Arifien A F. 2005. Perbandingan struktur anatomis, sifat fisis, dan sifat mekanis kayu jati unggul dan kayu jati konvensional. Jurnal Ilmu & Teknologi Kayu Tropis Vol. 3: 9-15.

Wahyudi I, Priadi T, Rahayu I S. 2014a. Karakteristik dan sifat-sifat dasar kayu jati unggul umur 4 dan 5 tahun asal Jawa Barat. Jurnal Ilmu Pertanian Indonesia Vol. 19: 50-56.

Wahyudi I, Sinaga DKD, Muhran, Jasni LB. 2014b. Pengaruh jarak tanam terhadap pertumbuhan pohon dan beberapa sifat fisis-mekanis kayu jati cepat tumbuh.

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Jurnal Ilmu Pertanian Indonesia Vol. 19: 204-210.

Waliszewska B, Wlodzimierz P, Magdalena Z, Agata S W, Hanna W, Agniezka S D. 2015. The diversification of chemical compotition of pine wood depending on tree age. Forestry and Wood Technology Vol. 91: 182-187

Wardani B W, Budi S. 2009. Pertumbuhan tanaman jati dari berbagai ras lahan di Pulau Muna. Jurnal Penelitian Hutan Tanaman Vol. 6: 63 – 71

White K. J. 1991. Teak: some aspects of research and development. Publication 1991/17. FAO Regional Office for Asia and the Pacific (RAPA). Bangkok

Windeisen E, Klassen A, Wegener G. 2003. On the chemical characterisation of plantation teakwood from Panama. Holz Roh Werkst Vol. 61: 416–418

Withouck H, Boeykens A, Luyten W, Lavigne R, Wagemans J, Broucke M V. 2019. Phenolic composition, antimicrobial and antioxidant properties of belgian apple wood extracts. Journal of Biologically Active Products from Nature Vol. 9: 24–38

Woo W L, Watson P, Mansfield S D. 2005. The effect of pine mountain bettle attack on lodgepole pine wood morphology and chemistry: implication for wood and fiber quality. Wood and Fiber Science Vol. 37: 112–126

Woodruff D R, Meinzer F C. 2011. Water stress, shoot growth and storage of non-structural carbohydrates along a tree height gradient in a tall conifer. Plant, Cell Environtment Vol. 34: 1920–1930

Würth M K R, Peláez-Riedl S, Wright S J, Körner C. 2004. Non-structural carbohydrate pools in a tropical forest. Oecologia Vol. 143: 11–24

Yang C S, Landau J M, Huang M T, Newmark H L. 2001. Inhibition of carcinogenesis by dietary polyphenolic compounds. Annual Review Nutrition Vol. 21: 381-406.

Yeh T F, Braun J L, Goldfarb B, Chang H, Kadla J F. 2006. Morphological and chemical variations between juvenile wood, mature wood, and compression wood of loblolly pine (*Pinus taeda* L.). Holzforschung Vol. 60: 1–8

Yongram C, Sungthong B, Puthongking P, Weerapreeyakul N. 2019. Chemical composition, antioxidant and cytotoxicity activities of leaves, bark, twigs and oleo-resin of *Dipterocarpus alatus*. Molecules Vol. 24: 3083–3090

Yu L W Y, Jin J W, Gangcheng J Q, Wang X. 2020. Effects of tree age on the chemical compositions and antioxidant activities of ‘Coratina’ and ‘Koroneiki’ olive oils from youth trees in China. Jiangnan University. Wuxi

Zelinka S L, Stone D S. 2011. The effect of tannins and pH on the corrosion of steel in wood extracts Vol. 62: 739–744

UNIVERSITAS
GADJAH MADA
Kadar Ekstraktif Kulit dan Kayu Jati Unggul Nusantara (JUN) Umur 6 dan 8 Tahun dari BKPH Yogyakarta
FAIS RAHMAN, Prof. Dr. Ganis Lukmandaru, S.Hut., M.Agr.,
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Zhang H, Wang C, Wang X. 2014. Spatial variations in non-structural carbohydrates in stems of twelve temperate tree species. Trees Vol. 28: 77–89

Zhang Q, Jia X, Shao M, Ma C. 2018. Unfolding non-structural carbohydrates from Sapling to dying black locust on China's Loess Plateau. Journal of Plant Growth Regulation Vol. 37: 794–802

Zobel B J, Buijtenen J P. 1989. Wood variation – Its causes and control. International Association of Wood Anatomists Journal Vol. 10: 1-363

Villagomez H Z, Peterson D M, Herrin L, Young R A. 2005. Antioxidant activity of different components of pine species. Holzforschung Vol. 59: 156-162

Zule J, Cufar Ca, Tisler V. 2015. Lipophilic extractives in heartwood of european larch (*Larix decidua* Mill). Drvna Industrija Vol. 66: 305 – 313

Zulkahfi. 2019. Kadar ekstraktif dan ketahanan terhadap jamur pada kayu Jati Plus Perhutani dengan perlakuan peneresan. Tesis Fakultas Kehutanan UGM. Yogyakarta.

Zulkahfi, Irawati D, Listyanto T, Rodiana D, Lukmandaru, G. 2020. Kadar ekstraktif dan sifat warna kayu jati plus Perhutani umur 11 tahun dari KPH Ngawi. Ilmu Kehutanan Vol. 14: 213-227