

REFERENSI

- Aldaqaal, S. M., Maqbul, A. A., Alhammad, A. A., Alghamdi, A. S., Alharbi, B. A., Alharbi, M. T., Alhazmi, O. M., & Zaylaee, Y. O. (2020). The Impact of Body Mass Index on the Clinicopathological and Prognostic Factors of Colorectal Cancer in Saudi Arabia. *Cureus*, *12*(11), e11789. <https://doi.org/10.7759/cureus.11789>
- American Joint Committee on Cancer. Colon and Rectum. Amin MB, Edge S, Greene F, Byrd DR, Brookland RK, et al, eds. AJCC Cancer Staging Manual. 8th ed. New York, NY: Springer; 2017.
- Aune *et al.* (2011) 'Dietary fibre, whole grains, and risk of colorectal cancer: Systematic review and dose-response meta-analysis of prospective studies', *BMJ (Online)*, p. 1082. doi: 10.1136/bmj.d6617.
- Board *et al.* (2021) 'Colorectal Cancer: Statistics', pp. 5–6. <https://www.cancer.net/cancer-types/colorectal-cancer/statistics>
- Boyd *et al.* (1976) 'Plasma CEA, tumor CEA, and tumor histology', *Journal of Surgical Oncology*, *8*(6), pp. 507–512. doi: 10.1002/jso.2930080610.
- Candrawati *et al.* (2018). 'Correlation of neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio, lymphocyte-to-monocyte ratio and carcinoembryonic antigen level in colorectal cancer', *Jurnal Kedokteran dan Kesehatan Indonesia* *9*(2): pp. 82–88.
- Canreg fkkmk. (2018). *RKBR Maret 2018*. Universitas Gadjah Mada. <https://canreg.fk.ugm.ac.id/laporan-data/registrasi-kanker-berbasis-rumah-sakit-dr-sardjito-fkkmk-ugm/rkbr-maret-2018/>
- Care, P., It, D. (2019) 'Clinics in Surgery The Role of Carcinoembryonic Antigen (CEA) Testing in', *4*, pp. 2–5.
- Chung *et al.* (2013). Prognostic significance of serum carcinoembryonic antigen normalization on survival in rectal cancer treated with preoperative chemoradiation. *Cancer research and treatment*, *45*(3), 186–192. <https://doi.org/10.4143/crt.2013.45.3.186>
- Coussens, L. M., & Werb, Z. (2002). Inflammation and cancer. *Nature*, *420*(6917), 860–867. <https://doi.org/10.1038/nature01322>
- Derwinger, K., Kodeda, K., Bexen-Lindskog, E., Taflin, H. (2009). *Tumour differentiation grade is associated with TNM staging and the risk of node metastasis in colorectal cancer*. *Acta Oncologica*, *49*(1), 57–62. doi:10.3109/02841860903334411
- Dolan, R. D., Lim, J., McSorley, S. T., Horgan, P. G., & McMillan, D. C. (2017). The role of the systemic inflammatory response in predicting outcomes in patients with operable cancer: Systematic review and meta-analysis. *Scientific reports*, *7*(1), 16717. <https://doi.org/10.1038/s41598-017-16955-5>
- Edge *et al.* (2009). *AJCC Cancer staging manual*. Springer Verlag.
- Ester, M. S. 2014. Perbedaan Kadar Cea Dan Ca 19-9 Dalam Serum Berdasarkan Differensiasi Sel Pada Kanker Kolorektal. Medan: Departemen Penyakit Dalam USU
- Fearon, E.R., Bommer, G.T. *Molecular Biology of Colorectal Cancer*. In V.T. DeVita, T.S. Lawrence & S.A. 2008. p. 152-167.

- Filiz *et al.* (2009) ‘Persistent high postoperative carcinoembryonic antigen in colorectal cancer patients- Is it important?’, *Clinics*, 64(4), pp. 287–294. doi: 10.1590/S1807-59322009000400004.
- Fleming *et al.* (2012) ‘Colorectal carcinoma: Pathologic aspects’, *Journal of Gastrointestinal Oncology*, 3(3), pp. 153–173. doi: 10.3978/j.issn.2078-6891.2012.030.
- Frontin, K. M. (2002) ‘Colon cancer’, *Ethnicity and Disease*, 12(1), pp. 1–9. <https://emedicine.medscape.com/article/277496-overview>
- Globocan (2020) ‘Colorectal Cancer Incidence in The World’, *The Global Cancer Observatory*, 419, pp. 1–2.
- Hamilton, S.R. (2000). Carcinoma of the colon and rectum. *World health organization classification of tumors. Pathology and genetics of tumors of the digestive system*, pp.105-119.
- Howlader N, Noone AM, Krapcho M, M. D. (2016) ‘Key Statistics for Colorectal Cancer’, *SEER Cancer Statistics Review*, p. Web. Available at: <https://www.cancer.org/cancer/colon-rectal-cancer/about/key-statistics.html>.
- Ilantzis *et al.* (2002) ‘Deregulated expression of the human tumor marker CEA and CEA family member CEACAM6 disrupts tissue architecture and blocks colonocyte differentiation’, *Neoplasia*, 4(2), pp. 151–163. doi: 10.1038/sj/neo/7900201.
- Jakubowska *et al.* (2020) ‘Pre- and postoperative neutrophil and lymphocyte count and neutrophil-to-lymphocyte ratio in patients with colorectal cancer’, *Molecular and Clinical Oncology*, 13(5), pp. 1–10. doi: 10.3892/mco.2020.2126.
- Josh, Farkas. 2019. *Pulmcrit: Neutrophil-lymphocyte ratio (NLR): Free upgrade to your WBC*. EMCrit Project. <https://emcrit.org/pulmcrit/nlr/>
- Kidoguchi, S., Sugano, N., Tokudome, G., Yokoo, T., Yano, Y., Hatake, K., & Nishiyama, A. (2020). *New Concept of Onco-Hypertension and Future Perspectives. Hypertension*. doi:10.1161/hypertensionaha.12
- Kuipers *et al.* (2015) ‘Colorectal cancer.’, *Nature reviews. Disease primers*, 1, p. 15065. doi: 10.1038/nrdp.2015.65.
- Li *et al.* (2014) ‘Prognostic role of neutrophil-to-lymphocyte ratio in colorectal cancer: A systematic review and meta-analysis’, *International Journal of Cancer*, 134(10), pp. 2403–2413. doi: 10.1002/ijc.28536.
- Li *et al.* (2020) ‘Predictive values of neutrophil-to-lymphocyte ratio on disease severity and mortality in COVID-19 patients: a systematic review and meta-analysis’, *Critical Care*, 24(1), pp. 1–10. doi: 10.1186/s13054-020-03374-8.
- Littlejohn *et al.* (2012). Systematic review and meta-analysis of the evidence for flexible sigmoidoscopy as a screening method for the prevention of colorectal cancer, *British Journal of Surgery*, Volume 99, Issue 11, November 2012, Pages 1488–1500, <https://doi.org/10.1002/bjs.8882>
- Locker *et al.* (2006) ‘ASCO 2006 update of recommendations for the use of tumor markers in gastrointestinal cancer’, *Journal of Clinical Oncology*, 24(33), pp. 5313–5327. doi: 10.1200/JCO.2006.08.2644.
- Lote, H., Spiteri, I., Ermini, L., Vatsiou, A., Roy, A., McDonald, A., ... Valeri, N. (2017). *Carbon dating cancer: defining the chronology of metastatic*

- progression in colorectal cancer. Annals of Oncology*, 28(6), 1243–1249. doi:10.1093/annonc/mdx074
- Mallappa *et al.* (2013) ‘Preoperative neutrophil to lymphocyte ratio >5 is a prognostic factor for recurrent colorectal cancer’, *Colorectal Disease*, 15(3), pp. 323–328. doi: 10.1111/codi.12008.
- Martins, E. C. *et al.* (2019) ‘Neutrophil-lymphocyte ratio in the early diagnosis of sepsis in an intensive care unit: A case-control study’, *Revista Brasileira de Terapia Intensiva*, 31(1), pp. 63–70. doi: 10.5935/0103-507X.20190010.
- Matsuda T, Saito Y, Fujii T, Uraoka T, Nakajima T, Kobayashi N, Emura F, Ono A, Shimoda T, Ikematsu H, Fu KI, Sano Y, Fujimori T. Size does not determine the grade of malignancy of early invasive colorectal cancer. *World J Gastroenterol.* 2009 Jun 14;15(22):2708-13. doi: 10.3748/wjg.15.2708. PMID: 19522020; PMCID: PMC2695885.
- Ozdemir *et al.* (2014) ‘Pretreatment neutrophil/lymphocyte ratio as a prognostic aid in colorectal cancer’, *Asian Pacific Journal of Cancer Prevention*, 15(6), pp. 2647–2650. doi: 10.7314/APJCP.2014.15.6.2647.
- Pine, *et al.* (2015) ‘Systemic neutrophil-to-lymphocyte ratio in colorectal cancer : the relationship to patient survival , tumour biology and local lymphocytic response to tumour’, *British Journal of Cancer*, (September 2014), pp. 204–211. doi: 10.1038/bjc.2015.87.
- Rifai, N., Horvath, A. R., Wittwer, C. T. (2017). *Tietz textbook of clinical chemistry and molecular diagnostics*. Saunders.
- Rogers AC, Handelman GS, Solon JG, McNamara DA, Deasy J, Burke JP. Meta-analysis of the clinicopathological characteristics and peri-operative outcomes of colorectal cancer in obese patients. *Cancer Epidemiol.* 2017 Dec;51:23-29. doi: 10.1016/j.canep.2017.09.004. Epub 2017 Oct 4. PMID: 28987964.
- Sepulveda *et al.* (2017) ‘Molecular Biomarkers for the Evaluation of Colorectal Cancer : Guideline From the American Society for Clinical Pathology , College of American Pathologists , Association for Molecular Pathology , and the American Society of Clinical Oncology’, 35(13). doi: 10.1200/JCO.2016.71.9807.
- Siregar, G. A., & Anshari, F. (2019). Absolute Neutrophil Count Levels among Degree of Differentiation and Tumor Location in Colorectal Cancer Patients in Medan. *Open access Macedonian journal of medical sciences*, 7(20), 3472–3474. <https://doi.org/10.3889/oamjms.2019.443>
- Society, A. C. (2020) ‘American Cancer Society. Cancer Facts & Figures 2020’, *American Cancer Society*, pp. 1–52. Available at: <http://www.cancer.org/acs/groups/content/@nho/documents/document/caff2007pwsecuredpdf.pdf>.
- Summers R. M. (2010). *r Radiology*, 255(3), 707–720. <https://doi.org/10.1148/radiol.10090877>
- Sutrisna, I. W. W. (2018) ‘Correlation between histopathologic grading and carcinoembryonic antigen levels in colon carcino’, *Medicina*, 49(1), pp. 22–28. doi: 10.15562/medicina.v49i1.256.

- Tan *et al.* (2018) ‘Prognostic Significance of lymphocyte to monocyte ratio in colorectal cancer: A meta-analysis’, *International Journal of Surgery*. doi: 10.1016/j.ijso.2018.05.030.
- Tan, E., Gouvas, N., Nicholls, R. J., Ziprin, P., Xynos, E., & Tekkis, P. P. (2009). *Diagnostic precision of carcinoembryonic antigen in the detection of recurrence of colorectal cancer. Surgical Oncology*, 18(1), 15–24. doi:10.1016/j.suronc.2008.05.00
- Tazzyman, S., Lewis, C. E., & Murdoch, C. (2009). Neutrophils: key mediators of tumour angiogenesis. *International journal of experimental pathology*, 90(3), 222–231. <https://doi.org/10.1111/j.1365-2613.2009.00641.x>
- Thirunavukarasu *et al.* (2011) ‘C-stage in Colon Cancer: Implications of Carcinoembryonic Antigen Biomarker in Staging , Prognosis , and Management’, pp. 689–697. doi: 10.1093/jnci/djr078.
- Tomislav, D. (2021). Colon Cancer Clinical Presentation: History, Physical Examination. <https://emedicine.medscape.com/article/277496-clinical>.
- Torre *et al.* (2015) ‘Global cancer statistics, 2012’, *CA: A Cancer Journal for Clinicians*, 65(2), pp. 87–108. doi: 10.3322/caac.21262.
- Ueno, H., Mochizuki, H., Hashiguchi, Y., Ishiguro, M., Kajiwara, Y., Sato, T., ... Talbot, I. C. (2008). *Histological Grading of Colorectal Cancer. Annals of Surgery*, 247(5), 811–818. doi:10.1097/sla.0b013e318167580f
- Umar *et al.* (2004). Revised Bethesda Guidelines for hereditary nonpolyposis colorectal cancer (Lynch syndrome) and microsatellite instability.” *Journal of the National Cancer Institute* vol. 96,4, 261-268. doi:10.1093/jnci/djh034
- Wardhana, A., Tomy, L. 2019. Neutrophil Lymphocyte Ratio (NLR) as a predictive factor radiological response of Neoadjuvant Chemotherapy (NAC) in Locally Advanced Rectal Cancer (LARC). Bali: Bali Medical Journal (Bali MedJ) 2021, Vol 10, Number 2: 717-723
- WCRF (2021) ‘Cancer Facts and Figures’, *World Cancer Research Fund International*, pp. 1–4. Available at: <http://www.wcrf.org/int/cancer-facts-figures/worldwide-data>.
- World Health Organization (2014) ‘Cancer Country Profiles: Indonesia’, *Cancer Country Profiles*, pp. 22–23.
- Yan *et al.* (2016) ‘The CEA-/lo colorectal cancer cell population harbors cancer stem cells and metastatic cells’, *Oncotarget*, 7(49), pp. 80700–80715. doi: 10.18632/oncotarget.13029
- Yarbro *et al.* (1999) “American Joint Committee on Cancer prognostic factors consensus conference.” *Cancer* vol. 86,11: 2436-46. doi:10.1002/(sici)1097-0142(19991201)86:11<2436::aid-cnrc35>3.0.co;2-#
- Zhang, S., Lin, M. and Zhang, H. (2015) ‘Diagnostic value of carcinoembryonic antigen and carcinoma antigen 19-9 for colorectal carcinoma’, 8(8), pp. 9404–9409.
- Zou, Z., Liu, H., Ning, N., Li, S., Du, X., & Li, R. (2016). Clinical significance of pre-operative neutrophil lymphocyte ratio and platelet lymphocyte ratio as prognostic factors for patients with colorectal cancer. *Oncology Letters*, 11, 2241-2248. <https://doi.org/10.3892/ol.2016.4216>