

Abhijeet V. Puri dan Amol D. Gholap. (2018). *Comparative Evaluation Of Papaya And Banana Starch As Disintegrant In Diclofenac Sodium Tablet Formulation*. International Journal of Pharmacy and Biological Sciences. www.ijpbs.com or www.ijpbsonline.com ISSN: 2230-7605 (Online); ISSN: 2321-3272 (Print)

Azhar Ahmed & Farukh Khan. (2013). *Extraction of Starch from Taro (Colocasia esculenta) and Evaluating it and further using Taro Starch as Disintegrating Agent in Tablet Formulation with Over All Evaluation*. Inveni Rapid: Novel Excipients Vol. 2013, Issue 2. ISSN 2278-408X. www.inveni.in

Anief, M. (2013). Ilmu Meracik Obat Teori dan Praktek, Penerbit University Press. Yogyakarta.

Akin-Ajani, O. D., Itiola, O. A., Odeku, O. A., (2014) *Effect of acid modification on the material and compaction properties of fonio and sweet potato starches*. Starch/Starke € 2014, 66, 749–759

Allen, L. V., Popovich, N. G., and Ansel, H. C. (2011) *Ansel's Pharmaceutical Dosage Forms and Drug Delivery System*, 9th Edition, Lippincott Williams and Wilkins, Philadelphia

Aulton's, M.E (2007) *Aulton's Pharmaceutics: The Design and Manufacture Medicine, 3 rd edition*, Churchill Livingstone Elsevier, London.

Dauda Usaini et all, 2019, *Binder and Disintegrant Performance of Native and Thermally Modified Dioscorea cayenensis Starches in Paracetamol Tablet*, Tropical Journal of Natural Product Research Trop J Nat Prod Res, <https://www.tjnpr.org/>

Ditjen POM RI (2020) *Farmakope Indonesia Edisi VI*, Depkes RI, Jakarta.

Gideon et all (2013). *Evaluation of disintegrating property of native and microcrystalline starches derived from digitaria iburua*. West African Journal of Pharmacy (2013) 24 (1) 64-71. Nigeria

Gunawan, D. dan Sri M, 2004, *Ilmu Obat Alam (Farmakognosi) Jilid 1*, Penebar Swadaya, Jakarta.

Gunsel W.C, Swart C. J.,and Kanig J. L., 1970, *Tablet in Lachman, L., Lieberman H.A., (Editor), The Theory Practise of Industrial Pharmacy*, Leo and Febiger, Philadelphia

L) sebagai bahan penghancur tablet. UIN Alauddin Makassar.

<http://journal.uui.ac.id/index.php/JIF>

Kandekar UY. (2020). *An Alternative Natural Disintegrant for Pharmaceutical Application Exploration of Elephant Foot Yam (Amorphophallus paeoniifolius) Starch.* Indian J of Pharmaceutical Education and Research. 2021;55(1s):s209-s219

Kementerian Perdagangan Republik Indonesia. (2020). *Analisis Komoditi Jagung Bulan Januari 2020.* Kementerian Perdagangan Republik Indonesia. Jakarta.

Kementerian Pertanian Republik Indonesia (2019). *Buletin Konsumsi Pangan.* Pusat Data dan Sistem Informasi Kementerian Pertanian. Jakarta

Kementerian Pertanian Republik Indonesia (2020). *Outlook Jagung Komoditas Pertanian Subsektor Tanaman Pangan.* Pusat Data dan Sistem Informasi Kementerian Pertanian. Jakarta.

Kusuma R et all. (2013). *Evaluation of Colocasia esculenta Starch as an Alternative Tablet Excipient to Maize Starch: Assessment by Preformulation and Formulation Studies.* International Journal of Pharma Sciences and Research (IJPSR). ISSN : 0975-9492. <http://www.ijpsr.info/>

Lachman L, Herbert A.L. And Joseph I.K. (1994). *Teori dan Praktek Farmasi Industri II,* Universitas Indonesia, Jakarta.

Lalduhsanga Puachu et all. (2016). *Physicochemical and disintegrant properties of glutinous rice starch of Mizoram India.* International Journal of Biological Macromolecules 95 (2017) 1298–1304. <http://dx.doi.org/10.1016/j.ijbiomac.2016.11.029> 0141-8130/

Lalduhsanga Puachu et all. (2018). *Taro starch (Colocasia esculenta) and citric acid modified taro starch as tablet disintegrating agents.* International Journal of Biological Macromolecules 118 (2018) 397–405. <https://doi.org/10.1016/j.ijbiomac.2018.06.086>

Madu SJ et all. (2011). *The Role Of Acid Treated Sweet Potato Starch (Microcrystalline Starch) On Disintegrant Property Of Paracetamol Tablet Formulation.* International Journal of Pharmaceutical Research and Innovation, Vol. 4: 32-39.

Mariam VL et all. (2015). *Effect Of Thermal And Chemical Modifications On The Mechanical And Release Properties Of Paracetamol Tablet Formulations Containing Corn,*

Cassava And Sweet Potato Starches As Filler-Binders. Asian Pacific Journal of Tropical Biomedicine 5(7): 585-590. <http://dx.doi.org/10.1016/j.apjtb.2015.05.003>

Mhana BY et all. (2017). *Evaluation of pearl millet starch as tablet disintegrant*. LIMUJ, Volume 2, PP 152-163, 2017. <http://doi.org/10.21502/limuj.019.02.2017>

Musa et all (2011) *Evaluaton Of Barley Hordeum Vulgare Starch As Tablet Disintegrant*, nt J Pharm Pharm Sci, Vol 3, Suppl 3, 2011, 7579, Nigeria

Nnabuike DN & Tochukuwu JN. (2017). *Optimization Of Metronidazole Tablet Formulation Using Manihot Utilissima Starch And A Combination Of Processing Techniques*. Future Journal of Pharmaceutical Sciences 3 (2017) 65 -70. <http://dx.doi.org/10.1016/j.fjps.2017.03.002>

Nwachukwu N & Ubieko EA (2020) *Disintegrant Properties of Native Starches obtained from Cassava, Sweet Potato and Corn in Ibuprofen Tablet Formulations*. Journal of Drug Delivery and Therapeutics. 2020; 10(5):264-273. <http://dx.doi.org/10.22270/jddt.v10i5.4324>

Odeniyi MA, Babalola AO, Ayorinde JO. (2013). *Evaluation of Cedrela gum as a binder and bioadhesive component in ibuprofen tablet formulations*. Braz J Pharm Sci 2013; 49(1): 95-105.

Olufunke D, et all. (2015). *Disintegrant properties of banana starch obtained from the unripe fruits of Musa sapientum L*. Journal of Applied Pharmaceutical Science Vol. 4 (09), pp. 083-088, <http://doi.org/10.7324/JAPS.2014.40915>

Olufunke D, et all. (2016). *Evaluation of the disintegrant properties of native and modified forms of fonio and sweet potato starches*. Starch/Stärke 2016, 68, 169–174. <http://doi.org/10.1002/star.201500188>

Olayemi OJ, et all. (2016). *Evaluation Of Disintegrant Properties Of Neorautanenia Mitis Starch*. Journal of Phytomedicine and Therapeutic Vol. 15(2) 52 – 63. www.jopat.gov.ng

Poedjiadi, A., dan Supriyanti, F.M.T. (2009) *Dasar-dasar Biokimia*, Universitas Indonesia, Jakarta.

Puri A. V et. al. (2013). *Formulation and Evaluation of Diclofenac Sodium Tablet using Isolated starch from Unripe Papaya Fruits as Disintegrant*. Indo American Journal of Pharm Research.2013;3(11). <http://www.iajpr.com/index.php/en/> ISSN NO: 2231-6876

Excipient. Journal of Chemical and Pharmaceutical Research 4(6):3134-3138. ISSN : 0975-7384 <http://www.jocpr.com/>

Rashid, I., Al Omari, M. M. H., & Badwan, A. A. (2013). From native to multifunctional starch- based excipients designed for direct compression formulation. *Starch/Staerke*, 65(7–8), 552–571

Rudnic, E. M., and Kottke, M. K., 2002, *Tablet Dosage Form* in Banker, G. S., and Rhodes, C. T., *Modern Pharmaceutics*, 4th Edition Revised and Expanded, Marcel Dekker Inc., New York. 287-309.

Sunitasonartiya and Dr. Shikhaagrawal. (2018). *Formulation And Evaluation Of Ibuprofen Tablet Using Isolated Starch From Unripe Papaya Fruits As A Disintegrant*. International Journal of Advanced Research 6(6), 792-804. <http://dx.doi.org/10.21474/IJAR01/7277>

Sylvester OE et all (2016) *A Comparative Investigation Of The Disintegrant Efficiency Of Musa Paradisiaca L. And Musa Sapientum L. Starches In Paracetamol Tablet Formulations*. Journal of Pharmacy and Bioresources. <http://dx.doi.org/10.434/jpb.v13i2.7>

Singh, N., K. S. Sandhu, and M. Kaur, 2005, *Physicochemical Properties Including Granular Morphology, Amylose Content, Swelling and Solubility, Thermal and Pasting Properties of Starches from Normal, Waxy, High Amylose and Sugary Corn*, Department of Food Science and Technology , Guru Nanak Dev University , Amritsar, India.

Siregar, C.J.P., dan Wikarsa, S., 2010, *Teknologi Farmasi Sediaan Tablet DasarDasar Praktis*, Penerbit Buku Kedokteran EGC, Jakarta.

Soebagio, B., Sriwododo, Adhika A. S, 2009, *Uji Sifat Fisikokimia Pati Biji Durian (Durio Zibethinus Murr) Alami Dan Modifikasi Secara Hidrolisis Asam*, Universitas Padjajaran, Bandung.

Syamsuni, H.A., 2007, *Ilmu Resep, Kedokteran EGC*, Jakarta.

Thoorens G, Krier F, Leclercq B, Evrad B, (2014). *Microcrystalline cellulose, a direct compression binder in a quality by design environment - A review*. Int. J. Pharmaceutics, 2014; 473:64-72