



DAFTAR PUSTKA

- Abang, A. F., Nanga, S. N., Kuate, A. F., Kouebou, C., Suh, C., Masso, C., Saethre, M. G., & Fiaboe, K. K. M. (2021). Natural enemies of fall armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in different agro-ecologies. *Insects*, 12(6), 1–23. <https://doi.org/10.3390/insects12060509>
- Abrahams, P., Bateman, M., Beale, T., Clottee, V., Cock, M., Colmenarez, Y., Corniani, N., Early, R., Godwin, J., Gomez, J., Moreno, P. G., Murphy, S. T., Oppong-Mensah, B., Phiri, N., Pratt, C., Richards, G., Silvestri, S., & Witt, A. (2017). Fall armyworm: Impacts and implications for Africa; evidence note (2). In CABI. CABI. https://doi.org/10.1564/v28_oct_02
- Agboyi, L. K., Goergen, G., Beseh, P., Mensah, S. A., Clottee, V. A., Glikpo, R., Buddie, A., Cafa, G., Offord, L., Day, R., Rwmushana, I., & Kenis, M. (2020). Parasitoid complex of fall armyworm, *Spodoptera frugiperda*, in Ghana and Benin. *Insects*, 11(2), 1–15. <https://doi.org/10.3390/insects11020068>
- Agboyi, L. K., Layode, B. F. R., Fening, K. O., Beseh, P., Clottee, V. A., Day, R., Kenis, M., & Babendreier, D. (2021). Assessing the potential of inoculative field releases of *Telenomus remus* to control *Spodoptera frugiperda* in Ghana. *Insects*, 12(8), 1–15. <https://doi.org/10.3390/insects12080665>
- Altieri, M. A. (1999). The ecological role of biodiversity in agroecosystems. *Agriculture, Ecosystems and Environment*, 74(1–3), 19–31. [https://doi.org/10.1016/S0167-8809\(99\)00028-6](https://doi.org/10.1016/S0167-8809(99)00028-6)
- Altieri, M. A., & Letourneau, D. K. (1984). Vegetation diversity and insect pest outbreaks. *Critical Reviews in Plant Sciences*, 2(2), 131–169. <https://doi.org/10.1080/07352688409382193>
- Amala, U., & Shivalingaswamy, T. M. (2018). Effect of intercrops and border crops on the diversity of parasitoids and predators in agroecosystem. *Egyptian Journal of Biological Pest Control*, 28(1), 1–4. <https://doi.org/10.1186/s41938-017-0015-y>
- Ashley, T. R., B.R. W., F.M. D., & K.L. A. (1989). The fall armyworm: a bibliography. *Florida Entomologist*, 72(1), 152–202.
- Assefa, F., & Ayalew, D. (2019). Status and control measures of fall armyworm (*Spodoptera frugiperda*) infestations in maize fields in Ethiopia: a review. *Cogent Food & Agriculture*, 5, 1–16. <https://doi.org/10.1080/23311932.2019.1641902>
- Audi, A. H., & Biliyaminu, A. (2018). Understanding insect parasitoids life history behaviour for optimal utilization in biological control of tropical injurious pest - a review. *Bayero Journal of Pure and Applied Sciences*, 11(2), 39–40. <https://doi.org/10.4314/bajopas.v11i2.6>
- Ayala, O. R., Navarro, F., & Virla, E. G. (2013). Evaluation of the attack rates and level of damages by the fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae), affecting corn-crops in the northeast of Argentina. *Revista de La Facultad de Ciencias Agrarias*, 45(2), 1–12.
- BBPOPT. (2020). *Update penyebaran hama baru pada jagung (FAW) dan strategi pengendaliannya*. <https://bbpopt.id/>



- Beckingham, C. (2007). Sweet corn growing. NSW Goverment. <https://www.dpi.nsw.gov.au/>
- Beserra, E. B., & Parra, J. R. P. (2005). Impact of the number of *Spodoptera frugiperda* egg layers on parasitism by *Trichogramma atopovirilia*. *Scientia Agricola*, 62(2), 190–193. <https://doi.org/10.1590/s0103-90162005000200016>
- Bianchi, F. J. J. A., Booij, C. J. H., & Tscharntke, T. (2006). Sustainable pest regulation in agricultural landscapes: A review on landscape composition, biodiversity and natural pest control. *Proceedings of the Royal Society B: Biological Sciences*, 273(1595), 1715–1727. <https://doi.org/10.1098/rspb.2006.3530>
- Borror, D. J., Triplehorn, C. A., & Jhonson, N. F. (2005). Borror and Delong's Introduction to the Study of insects (7th ed.). Thomson Brooks/Cole.
- Brotodjojo, R. R. R. (2009). Pengendalian hama dengan pengelolaan agroekosistem dalam kerangka pertanian berkelanjutan untuk mendukung ketahanan pangan. *Pangan*, 18(55), 17–24.
- Buchori, D., Bandungsahari, & Nurindah. (2008). Conservation of agroecosystem through utilization of parasitoid diversity: lesson for promoting sustainable agriculture and ecosystem health. *HAYATI Journal of Biosciences*, 15(4), 165–172. <https://doi.org/10.4308/hjb.15.4.165>
- Buchori, D., Hidayat, P., Kartosuwondo, U., Nurmansyah, A., & Meilin, A. (2002). Dinamika interaksi antara parasitoid Trichogrammatidae dan inangnya: faktor-faktor yang berpengaruh terhadap kualitas Trichogrammatidae sebagai agen pengendali hidup. *Laporan Penelitian Hibah Bersaing Perguruan Tinggi VII/3*.
- Buchori, D., Meilin, A., Hidayat, P., & Sahari, B. (2010). Species distribution of *Trichogramma* and *Trichogrammatoidea* Genus (Trichogrammatoidea: Hymenoptera) in Orang. *J. Issaas*, 16(1), 83–96.
- Bueno, R. C. O. F., Bueno, A. F., Moscardi, F., Parra, J. R. P., & Hoffmann-Campo, C. B. (2010). Lepidopteran larva consumption of soybean foliage: Basis for developing multiple-species economic thresholds for pest management decisions. *Pest Management Science*, 67(2), 170–174. <https://doi.org/10.1002/ps.2047>
- Bueno, R. C. O. F., Carneiro, T. R., Pratirossi, D., Bueno, A. F., & Fernandes, O. A. (2008). Biology and thermal requirements of *Telenomus remus* reared on fall armyworm. *Ciencia Rural Santa Maria*, 38(1), 1–6.
- CABI. (2020). *Spodoptera frugiperda* (fall armyworm). <https://www.cabi.org/isc/datasheet/29810>
- Camargo, L. F., Brito, R. A., & Penteado-Dias, A. M. (2015). Redescription of *Campoletis sonorensis* (Cameron, 1886) (Hymenoptera, Ichneumonidae, Campopleginae), parasitoid of *Spodoptera frugiperda* (J. E. Smith, 1797) (Lepidoptera, Noctuidae) in Brazil. *Brazilian Journal of Biology*, 75(4), 989–998. <https://doi.org/10.1590/1519-6984.04914>
- Canico, A., Mexia, A., & Santos, L. (2020). First report of native parasitoids of fall armyworm *Spodoptera frugiperda* Smith (Lepidoptera: Noctuidae) in Mozambique. *Insects*, 11(9), 1–12. <https://doi.org/10.3390/insects11090615>



- Capinera, J. L. (2017). Fall armyworm, *Spodoptera frugiperda* Smith (Insecta: Lepidoptera: Noctuidae). *IFAS Extension. University of Florida*, 1–6. <https://extension.entm.psu.edu/fieldcropsipm/insects/fall-armyworm.php>
- Casmuz, A., Juarez, M. L., Socias, M. G., Murua, M. G., Prieto, S., Medina, S., Willink, E., & Gastaminza, G. (2010). Review of the host plants of fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Revista de La Sociedad Entomológica Argentina*, 69(3–4), 209–231.
- Cave, R. D. (2000). Biology, ecology and use in pest management of *Telenomus remus*. *Biocontrol News and Information*, 21(1), 21–26.
- Chapman, D., Purse, B. V., Roy, H. E., & Bullock, J. M. (2017). Global trade networks determine the distribution of invasive non-native species. *Global Ecology and Biogeography*, 26(8), 907–917. <https://doi.org/10.1111/geb.12599>
- Chen, X., Tang, P., Zeng, J., van Achterberg, C., & He, J. (2014). Taxonomy of parasitoid wasps in China: an overview. *Biological Control*, 68(1), 57–72. <https://doi.org/10.1016/j.biocontrol.2013.06.007>
- Coll, M. (2009). Conservation biological control and the management of biological control services: are they the same? *Phytoparasitica*, 37(3), 205–208. <https://doi.org/10.1007/s12600-009-0028-5>
- Cottrell, T. E., & Yeargan, K. V. (1999). Factors influencing dispersal of larval *Coleomegilla maculata* from the weed *Acalypha ostryaefolia* to sweet corn. *Entomologia Experimentalis et Applicata*, 90(3), 313–322. <https://doi.org/10.1046/j.1570-7458.1999.00451.x>
- Dainese, M., Montecchiari, S., Sitzia, T., Sigura, M., & Marini, L. (2017). High cover of hedgerows in the landscape supports multiple ecosystem services in Mediterranean cereal fields. *Journal of Applied Ecology*, 54(2), 380–388. <https://doi.org/10.1111/1365-2664.12747>
- Dasilva, C. S. B., Vieira, J. M., Loiacono, M., Margaria, C., & Parra, J. R. P. (2015). Evidence of exploitative competition among egg parasitoids of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in maize. *Revista Colombiana de Entomología*, 41(2), 184–186.
- Day, R., Abrahams, P., Bateman, M., Beale, T., Clottee, V., Cock, M., Colmenarez, Y., Corniani, N., Early, R., Godwin, J., Gomez, J., Moreno, P. G., Murphy, S. T., Oppong-Mensah, B., Phiri, N., Pratt, C., Silvestri, S., & Witt, A. (2017). Fall armyworm: Impacts and implications for Africa. *Outlooks on Pest Management*, 28(5), 196–201. https://doi.org/10.1564/v28_oct_02
- Delgado, C. L. (2003). Rising consumption of meat and milk in developing countries has created a new food revolution. *Journal of Nutrition*, 133(11), 3907–3910. <https://doi.org/10.1093/jn/133.11.3907s>
- Direktorat Jenderal Prasarana dan Sarana Pertanian. (2019). Saran bahan aktif insektisida untuk pengendalian hama ulat grayak *Spodoptera frugiperda*. <https://psp.pertanian.go.id/>
- Early, R., Gonzalez-Moreno, P., Murphy, S. T., & Day, R. (2018). Forecasting the global extent of invasion of the cereal pest *Spodoptera frugiperda*, the fall armyworm. *NeoBiota*, 40, 25–50. <https://doi.org/10.3897/neobiota.40.28165>
- Edwards, R. L. (1954). The host-finding and oviposition behaviour of *Mormoniella*



- vitripenniis* (Walker) (Hym., Pteromalidae), parasite of Muscoid flies. *Behaviour*, 7(2), 88–112.
- EPPO. (2019). *Spodoptera frugiperda* continues to spread in Asia. <https://iapps2010.me/2019/03/25/eppo-2019-053-spodoptera-frugiperda-continues-to-spread-in-asia/>
- FAO. (2018). Briefing Note on FAO Actions on Fall Armyworm in Africa. <http://www.fao.org/food-chain-%0Ahttp://www.fao.org/fall-armyworm/en/>
- FAO. (2020). Fall armyworm map. <http://www.fao.org/fall-armyworm/monitoring-tools/faw-map/en/>
- FAOSTAT. (2021). *World crop production data*. <http://www.fao.org/faostat/en/>
- Ferrer, F. (2001). Biological control of agricultural insect pests in Venezuela; advances, achievements, and future perspectives. *Biocontrol News and Information*, 22(3), 67–74. <https://www.researchgate.net/publication/308881474>
- Gagic, V., Tscharntke, T., Dormann, C. F., Gruber, B., Wilstermann, A., & Thies, C. (2011). Food web structure and biocontrol in a four-trophic level system across a landscape complexity gradient. *Proceedings: Biological Sciences*, 278(1720), 2946–2953. <https://doi.org/10.1098/rspb.20>
- Ganiger, P. C., Yeshwanth, H. M., Muralimohan, K., Vinay, N., Kumar, A. R. V., & Chandrashekara, K. (2018). Occurrence of the new invasive pest, fall armyworm, *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae), in the maize fields of Karnataka, India. *Current Science*, 115(4), 621–623. <https://doi.org/10.18520/cs/v115/i4/621-623>
- Garbach, K., Milder, J. C., Montenegro, M., Karp, D. S., & DeClerck, F. A. J. (2014). Biodiversity and ecosystem services in agroecosystems. *Encyclopedia of Agriculture and Food Systems*, 2, 21–40. <https://doi.org/10.1016/B978-0-444-52512-3.00013-9>
- Ghahari, H., Hayat, R., Tabari, M., Ostovan, H., & Imani, S. (2008). A contribution to the predator and parasitoid fauna of rice pests in Iran, and a discussion on the biodiversity and IPM in rice fields. *Linzer Biologische Beitraege*, 40(1), 735–764.
- Goergen, G., Kumar, P. L., Sankung, S. B., Togola, A., & Tamo, M. (2016). First report of outbreaks of the fall armyworm *Spodoptera frugiperda* (J E Smith) (Lepidoptera, Noctuidae), a new alien invasive pest in West and Central Africa. *PLoS ONE*, 11(10), 1–9. <https://doi.org/10.1371/journal.pone.0165632>
- Gurr, G. M., Wratten, S. D., & Altieri, M. A. (2004). Ecological engineering for pest management: advances in habitat manipulation for arthropods. CSIRO Publishing. <https://doi.org/10.1079/9780851999036.0000>
- Hailu, G., Niassy, S., Zeyaur, K. R., Ochatum, N., & Subramanian, S. (2018). Maize-legume intercropping and push-pull for management of fall armyworm, stemborers, and striga in Uganda. *Agronomy Journal*, 110(6), 2513–2522. <https://doi.org/10.2134/agronj2018.02.0110>
- Hajek, A. E. (2004). Natural enemies an introduction to biological control. Cambridge University Press.
- Hardke, J. T., Lorenz, G. M., & Leonard, B. R. (2015). Fall armyworm (Lepidoptera: Noctuidae) ecology in Southeastern cotton. *Journal of Integrated Pest Management*, 6(1), 1–10. <https://doi.org/10.1080/21588095.2015.1008300>



Management, 6(1), 1–8. <https://doi.org/10.1093/jipm/pmv009>

- Hatt, S., Boeraeve, F., Artru, S., Dufrêne, M., & Francis, F. (2018). Spatial diversification of agroecosystems to enhance biological control and other regulating services: an agroecological perspective. *Science of the Total Environment*, 621, 600–611. <https://doi.org/10.1016/j.scitotenv.2017.11.296>
- Hawkins, B. A., Cornell, H. V., & Hochberg, M. E. (1997). Predators, parasitoids, and pathogens as mortality agents in phytophagous insect populations. *Ecology*, 78(7), 2145–2152. <https://doi.org/10.1140/epjp/i2012-12064-4>
- Herlinda, S., & Irsan, C. (2015). Pengendalian Hayati Hama Tumbuhan. Unsri Press.
- Hollingsworth, J. P., Hartstack, A. W., & Lingren, P. D. (1970). The spectral response of *Campoletis perdistinctus*. *Journal of Economic Entomology*, 63(6), 1758–1761. <https://doi.org/10.1093/jee/63.6.1758>
- Hunter, M. D. (2002). Landscape structure, habitat fragmentation, and the ecology of insects. *Agricultural and Forest Entomology*, 4(1), 159–166.
- IITA. (2018). Fall armyworm has reached the Indian subcontinent! <https://www.iita.org/news-item/fall-armyworm-has-reached-the-indian-subcontinent/>
- IPPC. (2016). Les degats causes par *Spodoptera frugiperda*. IPPC Official Pest Report.
- IPPC. (2018). First detection of fall armyworm on the border of Thailand. IPPC Official Pest Report. <https://www.ippc.int/en/>
- IPPC. (2019a). First detection report of the fall armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae) on maize in Myanmar. IPPC Official Pest Report. <https://www.ippc.int/en/>
- IPPC. (2019b). Report of first detection of fall armyworm (FAW) in Republic of Korea. IPPC Official Pest Report. <https://www.ippc.int/en/>
- IPPC. (2019c). Report of first detection of *Spodoptera frugiperda*-fall armyworm (FAW) in Egypt. IPPC Official Pest Report. <https://www.ippc.int/en/>
- IPPC. (2019d). Reporting fall armyworm (*Spodoptera frugiperda*) in Yemen. IPPC Official Pest Report. <https://www.ippc.int/en/>
- IPPC. (2019e). The occurrence of fall armyworm (*Spodoptera frugiperda*) in Orang. IPPC Official Pest Report. <https://www.ippc.int/en/>
- Jaraleno-Teniente, J., Lomeli-Flores, J. R., Rodriguez-Leyva, E., Bujanos-Muniz, R., & Rodríguez-Rodríguez, S. E. (2020). Egg parasitoids of orang of *Spodoptera frugiperda* (Smith) (lepidoptera: Noctuidae) in maize and sorghum in Central Mexico. *Insects*, 11(3), 1–10. <https://doi.org/10.3390/insects11030157>
- Johnson, N. F. (1984). Systematics of nearctic *Telenomus*: classification and revisions of the podisi and phymatae species groups (Hymenoptera: Scelionidae). *Bulletin of the Ohio Biological Orang*, 6(3), 1–118.
- Johnson, S. J. (1987). Migration and the life history strategy of the fall armyworm, *Spodoptera frugiperda* in the western hemisphere. *Orang Journal of Tropical Insect Science*, 8(4-5-6), 543–549. <https://doi.org/10.1017/s1742758400022591>



- Kementerian Pertanian. (2016). Outlook komoditas pertanian tanaman pangan jagung. *Pusat Data Dan Sistem Informasi. Orang*, 85.
- Kenis, Plessis, H., Van den Berg, J., Ba, M. N., Goergen, G., Kwadjo, K. E., Baoua, I., Tefera, T., Buddie, A., Cafa, G., Offord, L., Rwmushana, I., & Polaszek, A. (2019). *Telenomus remus*, a candidate parasitoid for the biological control of *Spodoptera frugiperda* in Africa, is already present on the continent. *Insects*, 10, 1–10. <https://doi.org/10.3390/insects10040092>
- Kremen, C. (2005). Managing ecosystem services: What do we need to know about their ecology? *Ecology Letters*, 8(5), 468–479. <https://doi.org/10.1111/j.1461-0248.2005.00751.x>
- Kremen, Claire, Williams, N. M., Aizen, M. A., Gemmill-Herren, B., LeBuhn, G., Minckley, R., Packer, L., Potts, S. G., Roulston, T., Steffan-Dewenter, I., Vázquez, D. P., Winfree, R., Adams, L., Crone, E. E., Greenleaf, S. S., Keitt, T. H., Klein, A. M., Regetz, J., & Ricketts, T. H. (2007). Pollination and other ecosystem services produced by mobile organisms: a conceptual framework for the effects of land-use change. *Ecology Letters*, 10(4), 299–314. <https://doi.org/10.1111/j.1461-0248.2007.01018.x>
- Laminou, S. A., Ba, M. N., Karimoune, L., Douumma, A., & Muniappan, R. (2020). Parasitism of locally recruited egg parasitoids of the fall armyworm in Africa. *Insects*, 11(7), 1–13. <https://doi.org/10.3390/insects11070430>
- Langer, A., & Hance, T. (2004). Enhancing parasitism of wheat aphids through apparent competition: A tool for biological control. *Agriculture, Ecosystems and Environment*, 102(2), 205–212. <https://doi.org/10.1016/j.agee.2003.07.005>
- Laouali, A., Ibrahim, B., Ba, M. N., Laouali, K., & Rangaswamy, M. (2018). Native parasitoids recruited by the invaded fall armyworm in Niger. *Indian Journal of Entomology*, 80(4), 1253–1254. <https://doi.org/10.5958/0974-8172.2018.00338.3>
- Liao, Y. L., Yang, B., Xu, M. F., Lin, W., Wang, D. S., Chen, K. W., & Chen, H. Y. (2019). First report of *Telenomus remus* parasitizing *Spodoptera frugiperda* and its field parasitism in southern China. *Journal of Hymenoptera Research*, 73, 95–102. <https://doi.org/10.3897/JHR.73.39136>
- Lichtenberg, E. M., Kennedy, C. M., Kremen, C., Batary, P., Berendse, F., Bommarco, R., Bosque-Perez, N. A., Carvalheiro, L. G., Snyder, W. E., Williams, N. M., Winfree, R., Klatt, B. K., Astrom, S., Benjamin, F., Brittain, C., Chaplin-Kramer, R., Clough, Y., Danforth, B., Diekötter, T., ... Crowder, D. W. (2017). A global synthesis of the effects of diversified farming systems on arthropod diversity within fields and across agricultural landscapes. *Global Change Biology*, 23(11), 4946–4957. <https://doi.org/10.1111/gcb.13714>
- Mangold, J. R. (1978). Attraction of *Euphasiopteryx ochracea*, *Corethrella* sp., and Gryllids to broadcast songs of the Southern Mole Cricket. *The Florida Entomologist*, 61(2), 57–61.
- Marino, P. C., & Landis, D. A. (1996). Effect of landscape structure on parasitoid diversity and parasitism in agroecosystems. *Ecological Applications*, 6(1), 276–284. <https://doi.org/10.2307/2269571>
- Marino, P. C., & Landis, D. A. (2000). Parasitoid community structure: implications for biological control in agricultural landscapes. In B. Ekbom, M. E. Irwin, & Y.



Robert (Eds.), *Interchanges of insects between agricultural and surrounding landscapes* (pp. 183–193). Springer, Dordrecht.

Matson, P. A., Parton, W. J., Power, A. G., & Swift, M. J. (1997). Agricultural intensification and ecosystem properties. *Science*, 277, 504–509. <https://doi.org/10.1126/science.277.5325.504>

MEA. (2005). Ecosystems and human well-being. In *Island Press*. <http://books.google.com/books?hl=en&lr=&orang=vNd3Po5ZURgC&oi=fnd&pg=PR13&dq=%22MA+used+to+assess+options+that+can+enhance+the+contribution%22+%22has+always+depended+on+the+services+provided+by%22+%22his+April+2000+Millennium+Report+to+the+United%22+&ots=g8k>

Midega, C. A. O., Pittchar, J. O., Pickett, J. A., Hailu, G. W., & Khan, Z. R. (2018). A climate-adapted push-pull system effectively controls fall armyworm, *Spodoptera frugiperda* (J E Smith), in maize in East Africa. *Crop Protection*, 105(August 2017), 10–15. <https://doi.org/10.1016/j.cropro.2017.11.003>

Molina-Ochoa, J., Carpenter, J. E., Heinrichs, E. A., & Foster, J. E. (2003). Parasitoids and parasites of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in the Americas and Caribbean Basin: An inventory. *Florida Entomologist*, 86(3), 254–289. [https://doi.org/10.1653/0015-4040\(2003\)086\[0254:PAPOSF\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2003)086[0254:PAPOSF]2.0.CO;2)

Murua, M. G., Molina-Ochoa, J., & Coviella, C. (2006). Population dynamics of the fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae) and its parasitoids in northwestern Argentina. *Florida Entomologist*, 89(2), 175–182. [https://doi.org/10.1653/0015-4040\(2006\)89\[175:PDOTFA\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2006)89[175:PDOTFA]2.0.CO;2)

Nettles, W. C., & Burks, M. L. (1975). A substance from *Heliothis virescens* larvae stimulating larviposition by females of the tachinid, *Archytas marmoratus*. *Journal of Insect Physiology*, 21(5), 965–978. [https://doi.org/10.1016/0022-1910\(75\)90108-0](https://doi.org/10.1016/0022-1910(75)90108-0)

Ogunfunmilayo, A. O., Kazeem, S. A., Idoko, J. E., Adebayo, R. A., Fayemi, E. Y., Adedibu, O. B., Oloyede-Kamiyo, Q. O., Nwogwu, J. O., Akinbode, O. A., Salihu, S., Offord, L. C., Buddie, A. G., & Ofuya, T. I. (2021). Occurrence of natural enemies of fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in Nigeria. *PLoS ONE*, 16(7), 1–13. <https://doi.org/10.1371/journal.pone.0254328>

Orphanides, G. M., & Gonzales, D. (1970). Effects of adhesive materials and host location on parasitization by a uniparental race of *Trichogramma pretiosum*. *Journal of Economic Entomology*, 63(6), 1891–1897.

Otim, M. H., Aropet, S. A., Opio, M., Kanyesigye, D., Opolot, H. N., & Tay, W. T. (2021). Parasitoid distribution and parasitism of the fall armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in different maize producing regions of Uganda. *Insects*, 12(2), 1–20. <https://doi.org/10.3390/insects12020121>

Pagiola, S., von Ritter, K., & Bishop, J. (2004). Assessing the economic value of ecosystem conservation. The World Bank Environment Department.

Philpott, S. M. (2013). Biodiversity and pest control services. *Encyclopedia of Biodiversity: Second Edition*, 1, 373–385. <https://doi.org/10.1016/B978-0-12-384719-5.00344-0>

Plecas, M., Gagic, V., Jankovic, M., Petrovic-Obradovic, O., Kavallieratos, N. G.,



- Tomanovic, Z., Thies, C., Tscharntke, T., & Cetkovic, A. (2013). Landscape composition and configuration influence cereal aphid-parasitoid-hyperparasitoid interactions and biological control differentially across years. *Agriculture, Ecosystems and Environment*, 183, 1–10. <https://doi.org/10.1016/j.agee.2013.10.016>
- Power, A. G. (2010). Ecosystem services and agriculture: Tradeoffs and synergies. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1554), 2959–2971. <https://doi.org/10.1098/rstb.2010.0143>
- Prasanna, B. M., Huesing, J. E., Eddy, R., & Peschke, V. M. (2018). Fall armyworm in Africa: a guide for integrated pest management (First Edit, pp. 1–120). CDMX: CIMMYT. www.maize.org.
- Querino, R. B., & Zucchi, R. A. (2005). An illustrated key to the species of *Trichogramma* (Hymenoptera: Trichogrammatidae) of Brazil. *Zootaxa*, 1073, 37–60. <https://doi.org/10.11646/zootaxa.1073.1.3>
- Quicke, D. L. J. (2012). We know too little about parasitoid wasp distributions to draw any conclusions about latitudinal trends in species richness, body size and biology. *PLoS ONE*, 7(2). <https://doi.org/10.1371/journal.pone.0032101>
- Rand, T. A., Tylianakis, J. M., & Tscharntke, T. (2006). Spillover edge effects: the dispersal of agriculturally subsidized insect natural enemies into adjacent natural habitats. *Ecology Letters*, 9(5), 603–614. <https://doi.org/10.1111/j.1461-0248.2006.00911.x>
- Raymond, C. M., Bryan, B. A., MacDonald, D. H., Cast, A., Strathearn, S., Grandgirard, A., & Kalivas, T. (2009). Mapping community values for natural capital and ecosystem services. *Ecological Economics*, 68, 1301–1315. <https://doi.org/10.1016/j.ecolecon.2008.12.006>
- Redlich, S., Martin, E. A., & Steffan-Dewenter, I. (2018). Landscape-level crop diversity benefits biological pest control. *Journal of Applied Ecology*, 55(5), 2419–2428. <https://doi.org/10.1111/1365-2664.13126>
- Rizali, A., Oktaviyani, Putri, S. D. P. S., Doananda, M., & Linggani, A. (2021). Invasion of fall armyworm *Spodoptera frugiperda*, a new invasive pest, alters native herbivore attack intensity and natural enemy diversity. *Biodiversitas*, 22(8), 3482–3488. <https://doi.org/10.13057/biodiv/d220847>
- Root, R. B. (1973). Organization of a plant-arthropod association in simple and diverse habitats: the fauna of collards (Brassica Oleracea). *Ecological Monographs*, 43(1), 95–124.
- Rosenheim, J. A. (1996). An evolutionary argument for egg limitation. *Evolution*, 50(5), 2089–2094. <https://doi.org/10.2307/2410766>
- Rusch, A., Chaplin-Kramer, R., Gardiner, M. M., Hawro, V., Holland, J., Landis, D., Thies, C., Tscharntke, T., Weisser, W. W., Winqvist, C., Woltz, M., & Bommarco, R. (2016). Agricultural landscape simplification reduces natural pest control: A quantitative synthesis. *Agriculture, Ecosystems and Environment*, 221(1), 198–204. <https://doi.org/10.1016/j.agee.2016.01.039>
- Rwomushana, I., Bateman, M., Beale, T., Beseh, P., Cameron, K., Chiluba, M., Clottee, V., Davis, T., Day, R., Early, R., Godwin, J., Gonzalez-Moreno, P., Kansiime, M., Kenis, M., Makale, F., Mugambi, I., Murphy, S., Nunda, W., Phiri, N., ... Tambo, J. (2018). Fall armyworm: impacts and implications for Africa; Evidence note update. In CABI. [https://www.invasive-](http://www.invasive-)



species.org/Uploads/InvasiveSpecies/FAW Evidence Note October 2018.pdf

- Saaksjarvi, I. E., Ruokolainen, K., Tuomisto, H., Haataja, S., Fine, P. V. A., Cárdenas, G., Mesones, I., & Vargas, V. (2006). Comparing composition and diversity of parasitoid wasps and plants in an Amazonian rain-forest mosaic. *Journal of Tropical Ecology*, 22(2), 167–176. <https://doi.org/10.1017/S0266467405002993>
- Salazar-Mendoza, P., Rodriguez-Saona, C., & Fernandes, O. A. (2020). Release density, dispersal capacity, and optimal rearing conditions for *Telenomus remus*, an egg parasitoid of *Spodoptera frugiperda*, in maize. *Biocontrol Science and Technology*, 30(10), 1–20. <https://doi.org/10.1080/09583157.2020.1776841>
- Sarmento, R. A., Aguiar, R. W. S., Aguiar, R. A. S. S., Vieira, S. M. J., Oliveira, H. G., & Holtz, A. M. (2002). Biology review, occurrence and control of *Spodoptera frugiperda* (Lepidoptera, Noctuidae) in corn in Brazil. *Bioscience Journal*, 18(2), 41–48.
- Sartiami, D., Dadang, Harahap, I. S., Kusumah, Y. M., & Anwar, R. (2020). First record of fall armyworm (*Spodoptera frugiperda*) in Orang and its occurrence in three provinces. *IOP Conference Series: Earth and Environmental Science*, 468(1), 1–9. <https://doi.org/10.1088/1755-1315/468/1/012021>
- Sharanabasappa, Kalleshwaraswamy, C. M., Asokan, R., Swamy, H. M. M., Maruthi, M. S., Pavithra, H. B., Hedge, K., Navi, S., Prabhu, S. T., & Goergen, G. (2018). First report of the fall armyworm, *Spodoptera frugiperda* (J E Smith) (Lepidoptera: Noctuidae), an alien invasive pest on maize in India. *Pest Management in Horticultural Ecosystems*, 24(1), 50–54. <http://www.indianjournals.com/ijor.aspx?target=ijor:pmhe&volume=20&issue=1&article=010>
- Sharanabasappa, Kalleshwaraswamy, C. M., Maruthi, M. S., & Pavithra, H. B. (2018). Biology of invasive fall army worm *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) on maize. *Indian Journal of Entomology*, 80(3), 540–543. <https://doi.org/10.5958/0974-8172.2018.00238.9>
- Sheehan, W. (1998). Response by specialist and generalist natural enemies to agroecosystem diversification: a selective review. *Environmental Entomology*, 15(3), 456–461. <https://doi.org/10.1093/ee/15.3.456>
- Shylesha, A. N., Jalali, S. K., Gupta, A., Varshney, R., Venkatesan, T., Shetty, P., Ojha, R., Ganiger, P. C., Navik, O., Subaharan, K., Bahktavatsalam, N., & Ballal, C. R. (2018). Studies on new invasive pest *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) and its natural enemies. *Journal of Biological Control*, 32(3), 145–151. <https://doi.org/10.18311/jbc/2018/21707>
- Silveira, L. C. P., Souza, I. L., Tomazella, V. B., & Mendez, H. A. G. (2019). Parasitoid insects. In B. Souza, L. L. Vazquez, & R. C. Marucci (Eds.), *Springer Nature Switzerland* (Natural En). https://doi.org/10.1007/978-3-030-24733-1_41
- Sirami, C., Gross, N., Baillod, A. B., Bertrand, C., Carrie, R., Hass, A., Henckel, L., Miguet, P., Vuillot, C., Alignier, A., Girard, J., Batary, P., Clough, Y., Violle, C., Giralt, D., Bota, G., Badenhausser, I., Lefebvre, G., Gauffre, B., ... Fahrig, L. (2019). Increasing crop heterogeneity enhances multitrophic diversity across agricultural regions. *Proceedings of the National Academy of Sciences of the United States of America*, 116(33), 1–6.



<https://doi.org/10.1073/pnas.1906419116>

- Sisay, B., Simiyu, J., Mendesil, E., Likhayo, P., Ayalew, G., Mohamed, S., Subramanian, S., & Tefera, T. (2019). Fall armyworm, *Spodoptera frugiperda* infestations in East Africa: Assessment of damage and parasitism. *Insects*, 10(7), 1–10. <https://doi.org/10.3390/insects10070195>
- Snyder, W. E., & Ives, A. R. (2003). Interactions between specialist and generalist natural enemies: parasitoids, predators, and pea aphid biocontrol. *Ecology*, 84(1), 91–107.
- Sperber, C. F., Nakayama, K., Valverde, M. J., & De Siqueira Neves, F. (2004). Tree species richness and density affect parasitoid diversity in cacao agroforestry. *Basic and Applied Ecology*, 5(3), 241–251. <https://doi.org/10.1016/j.baae.2004.04.001>
- Staver, C., Guharay, F., Monterroso, D., & Muschler, R. G. (2001). Designing pest-suppressive multistrata perennial crop systems: shade-grown coffee in Central America. *Agroforestry Systems*, 53, 151–170141. <https://doi.org/10.1023/A>
- Supeno, B., Tarmizi, Meidiwarman, & Haryanto, H. (2021). Keragaman parasitoid yang berasosiasi dengan telur hama baru *Spodoptera frugiperda* di pulau Lombok. *Prosiding SAINTEK*, 3, 418–423.
- Tawakkal, M. I., Buchori, D., Maryana, N., & Pudjianto. (2020). New association between *Spodoptera frugiperda* J.E. Smith (Lepidoptera: Noctuidae) and native natural enemies: Bioprospection of native natural enemies as biological control agents. *IOP Conference Series: Earth and Environmental Science*, 771(1), 1–8. <https://doi.org/10.1088/1755-1315/771/1/012030>
- Tooker, J. F., & Hanks, L. M. (2000). Influence of plant community structure on natural enemies of pine needle scale (Homoptera: Diaspididae) in urban landscapes. *Environmental Entomology*, 29(6), 1305–1311. <https://doi.org/10.1603/0046-225X-29.6.1305>
- Trisyono, Y. A., Suputa, Aryuwandari, V. E. F., Hartaman, M., & Jumari, J. (2019). Occurrence of heavy infestation by the fall Armyworm *Spodoptera frugiperda*, a new alien invasive pest, in corn Lampung Orang. *Jurnal Perlindungan Tanaman Orang*, 23(1), 156–160. <https://doi.org/10.22146/jpti.46455>
- Tscharntke, T., Klein, A. M., Kruess, A., Steffan-Dewenter, I., & Thies, C. (2005). Landscape perspectives on agricultural intensification and biodiversity - ecosystem service management. *Ecology Letters*, 8(8), 857–874. <https://doi.org/10.1111/j.1461-0248.2005.00782.x>
- Van Driesche, R. G. (1983). Meaning of “percent parasitism” in studies of insect parasitoids. *Environmental Entomology*, 12(6), 1611–1622. <https://doi.org/10.1093/ee/12.6.1611>
- Van Lenteren, J. C., & Bueno, V. H. P. (2003). Augmentative biological control of arthropods in Latin America. *BioControl*, 48(2), 123–139. <https://doi.org/10.1023/A:1022645210394>
- Vinson, S. B. (1984). Parasitoid-Host Relationship. *Chemical Ecology of Insects*, 205–233. https://doi.org/10.1007/978-1-4899-3368-3_8
- Wenxia, D., Chun, X., & Chengyun, L. (2016). Effect of diversified cropping on insect pests and natural enemies in agroecosystems. *Chinese Journal of Eco-*



Agriculture, 24(4), 435–442.

- Whitfield, J. B. (1998). Phylogeny and evolution of host-parasitoid interactions in hymenoptera. *Annual Review of Entomology*, 43, 129–151. <https://doi.org/10.1146/annurev.ento.43.1.129>
- Wyckhuys, K. A. , & O'Neil, R. J. (2006). Population dynamics of *Spodoptera frugiperda* Smith (Lepidoptera: Noctuidae) and associated arthropod natural enemies in Honduran subsistence maize. *Crop Protection*, 25(11), 1180–1190. <https://doi.org/10.1016/j.cropro.2006.03.003>
- Yu, S. J. (1991). Insecticide resistance in the fall armyworm, *Spodoptera frugiperda* (J. E. Smith). *Pesticide Biochemistry and Physiology*, 39(1), 84–91. [https://doi.org/10.1016/0048-3575\(91\)90216-9](https://doi.org/10.1016/0048-3575(91)90216-9)
- Zhang, W., Ricketts, T. H., Kremen, C., Carney, K., & Swinton, S. M. (2007). Ecosystem services and dis-services to agriculture. *Ecological Economics*, 64(2), 253–260. <https://doi.org/10.1016/j.ecolecon.2007.02.024>
- Zhao, H., Li, J., Guo, L., & Wang, K. (2020). Crop diversity at the landscape level affects the composition and structure of the vegetation-dwelling arthropod communities in naked oat (*Avena chinensis*) fields. *Orang Journal of Environmental Research and Public Health*, 18(1), 1–13. <https://doi.org/10.3390/ijerph18010030>