

## DAFTAR PUSTAKA

- Alamsyah, A. 2006. Taklukkan Penyakit dengan Teh hijau. Tangerang: Agro Media Pustaka.
- Belitz, H. D., Grosch, W., & Schieberle, P. (2009). Coffee, tea, cocoa. *Food chemistry*, 938-970.
- Bhumiratana, N., Adhikari, K. and Chambers, E. (2011) 'Evolution of sensory aroma attributes from coffee beans to brewed coffee', *LWT - Food Science and Technology*, 44(10), pp. 2185–2192. doi: 10.1016/j.lwt.2011.07.001.
- Cabrera, A., Collins, W. C., & Salgado, J. F. (2006). Determinants of individual engagement in knowledge sharing. *The International Journal of Human Resource Management*, 17(2), 245-264.
- de Castro, D. L. S. *et al.* (2020) 'Effect of high-velocity and traditional resistance exercise on serum antioxidants and inflammation biomarkers in older women: A randomized crossover trial', *Experimental Gerontology*, 139(June). doi: 10.1016/j.exger.2020.111026.
- Chacko, S. M., Thambi, P. T., Kuttan, R., & Nishigaki, I. (2010). Beneficial effects of green tea: a literature review. *Chinese medicine*, 5(1), 1-9.
- Chan, K. E. *et al.* (2011) 'Combined angiotensin-converting enzyme inhibition and receptor blockade associate with increased risk of cardiovascular death in hemodialysis patients', *Kidney International*, 80(9), pp. 978–985. doi: 10.1038/ki.2011.228.
- Engel, E. *et al.* (2002) 'Relevance of omission tests to determine flavour-active compounds in food: Application to cheese taste', *Food Quality and Preference*, 13(7–8), pp. 505–513. doi: 10.1016/S0950-3293(02)00136-2.
- Eveline, R. (1997). Kajian Produk Olahan Teh di Indonesia. Bogor: Institut Pertanian Bogor
- Farkas, O., Jakus, J. and Héberger, K. (2004) 'Quantitative structure - Antioxidant activity relationships of flavonoid compounds', *Molecules*, 9(12), pp. 1079–1088. doi: 10.3390/91201079.
- Feng, Z., Li, Y., Li, M., Wang, Y., Zhang, L., Wan, X., & Yang, X. (2019). Tea aroma formation from six model manufacturing processes. *Food chemistry*, 285, 347-354.
- Florentino, R. F., Tanchoco, C. C., Ramos, A. C., Mendoza, T. S., Natividad, E. P., Tangco, J. B. M., ... & Muhilal, S. A. (1991). Vitamin A Publications. *Science & Medicine*, 33, 1141-1154.
- Flores-Martínez, D., Uriás-Orona, V., Hernández-García, L., Rubio-Carrasco, W., Silva-Gutiérrez, K., Guevara-Zambrano, M., Prieto-Cadena, J., Serna-Mendez, T., Muy-Rangel, D., and 4 and Niño-Medina, G. (2018) 'Physicochemical Parameters , Mineral Composition , and Nutraceutical Properties of Ready-to-Drink Flavored-Colored', *Journal of Chemistry*,

2018, pp. 1–7.

- Gdoura, R. *et al.* (2007) ‘Published-Ahead-of-Print on December 12, 2007 by Journal of Andrology’, 039024(212), pp. 1–32.
- Goldstein, R. H. (2003) ‘Petrographic analysis of fluid inclusions’, *Fluid Inclusions-Analysis and Interpretation*, 32(April), pp. 9–54.
- Guo, X., Song, C., Ho, C. T., & Wan, X. (2018). Contribution of l-theanine to the formation of 2, 5-dimethylpyrazine, a key roasted peanutty flavor in Oolong tea during manufacturing processes. *Food Chemistry*, 263, 18-28.
- Han, Z. X. *et al.* (2016) ‘Green tea flavour determinants and their changes over manufacturing processes’, *Food Chemistry*, 212, pp. 739–748. doi: 10.1016/j.foodchem.2016.06.049.
- Hiramoto, S. *et al.* (2004) ‘Melanoidin, a food protein-derived advanced Maillard reaction product, suppresses Helicobacter pylori in vitro and in vivo’, *Helicobacter*, 9(5), pp. 429–435. doi: 10.1111/j.1083-4389.2004.00263.x.
- Ho, C. T., Zheng, X. and Li, S. (2015) *Tea aroma formation, Food Science and Human Wellness*. Beijing Academy of Food Sciences. doi: 10.1016/j.fshw.2015.04.001.
- Hofmann, T. and Schieberle, P. (1997) ‘Identification of Potent Aroma Compounds in Thermally Treated Mixtures of Glucose/Cysteine and Rhamnose/Cysteine Using Aroma Extract Dilution Techniques’, *Journal of Agricultural and Food Chemistry*, 45(3), pp. 898–906. doi: 10.1021/jf960456t.
- Joshi, R., & Gulati, A. (2015). Fractionation and identification of minor and aroma-active constituents in Kangra orthodox black tea. *Food chemistry*, 167, 290-298.
- Jumri, J., Yusmarini, Y., & Herawati, N. (2015). *Mutu Permen Jelli Buah Naga Merah (Hylocereus polyrhizus) dengan Penambahan Karagenan dan Gum Arab* (Doctoral dissertation, Riau University).
- Kaneko, M. K. *et al.* (2006) ‘Conservation of a platelet activating domain of Aggrus/podoplanin as a platelet aggregation-inducing factor’, *Gene*, 378(1–2), pp. 52–57. doi: 10.1016/j.gene.2006.04.023.
- Kara, D. (2009). Evaluation of trace metal concentrations in some herbs and herbal teas by principal component analysis. *Food Chemistry*, 114(1), 347-354.
- Karori, S. M. *et al.* (2007) ‘Antioxidant capacity of different types of tea products’, *African Journal of Biotechnology*, 6(19), pp. 2287–2296. doi: 10.5897/AJB2007.000-2358.
- Kimia, P. S. *et al.* (2013) ‘Analisis Kromatografi Cair Kinerja Tinggi Untuk Penetapan Kadar Asam Galat, Kafein Dan Epigalokatekin Galat Pada Beberapa Produk Teh Celup’, *Analisis Kromatografi Cair Kinerja Tinggi Untuk Penetapan Kadar Asam Galat, Kafein Dan Epigalokatekin Galat Pada Beberapa Produk Teh Celup*, 32(4), pp. 362–369. doi:

10.22146/agritech.9578.

- Kumalaningsih, S. (2007). Antioksidan, sumber dan manfaatnya. *Diambil dari: <http://antioxidancentre.com/html>*.
- Lau, H. *et al.* (2018) 'Characterising volatiles in tea (*Camellia sinensis*). Part I: Comparison of headspace-solid phase microextraction and solvent assisted flavour evaporation', *Lwt*, 94(October 2017), pp. 178–189. doi: 10.1016/j.lwt.2018.04.058.
- Lee, L. W., Cheong, M. W., Curran, P., Yu, B., & Liu, S. Q. (2016). Modulation of coffee aroma via the fermentation of green coffee beans with *Rhizopus oligosporus*: I. Green coffee. *Food Chemistry*, 211, 916-924.
- Liang, Y.R., Ye, Q., Jin, J., Liang, H. Lu, J.L. Du, Y.Y., and Dong, J. J. (2008) 'Chemical and Instrumental Assessment of Green Tea Sensory Preference', *International Journal of Food Properties*, 11, pp. 258–272. doi: 10.1080/10942910701299430.
- Liu, F., Yang, X., Wu, X., Xi, X., Gao, H., Zhang, S., ... & Lu, R. (2018). A dispersive magnetic solid phase microextraction based on ionic liquid-coated and cyclodextrin-functionalized magnetic core dendrimer nanocomposites for the determination of pyrethroids in juice samples. *Food chemistry*, 268, 485-491.
- Lucida, H. (2006). Determination of the ionization constants and the stability of catechin from gambir (*Uncaria gambir* (Hunter) Roxb). In *ASOPMS 12 International conference*.
- Martono, E., & Vohts, J. M. (2012). Support effects in cobalt-based ethanol steam reforming catalysts: Reaction of ethanol on Co/CeO<sub>2</sub>/YSZ (1 0 0) model catalysts. *Journal of catalysis*, 291, 79-86.
- Muawanah, A. *et al.* (2012) 'Penggunaan Bunga Kecombrang (*Etlingera Elatior*) Dalam Proses Formulasi Permen Jelly', *Jurnal Kimia VALENSI*, 2(4). doi: 10.15408/jkv.v2i4.270.
- Nemecz, George. "Green tea." *Journal of Modern Pharmacy* 9.1 (2002): 27.
- Ofmannt, T. H. (2006) 'AGRICULTURAL AND Molecular and Sensory Studies on the Umami Taste of', pp. 2688–2694.
- Palupi, N. S., Zakaria, F. R., & Prangdimurti, E. (2007). Pengaruh pengolahan terhadap nilai gizi pangan. *Modul e-Learning ENBP, Departemen Ilmu & Teknologi Pangan-Fateta-IPB*, 1-14.
- Pinnell, S. R. (2003) 'Cutaneous photodamage, oxidative stress, and topical antioxidant protection', *Journal of the American Academy of Dermatology*, 48(1), pp. 1–22. doi: 10.1067/mjd.2003.16.
- Robinson, T., 1995, Kandungan Organik Tumbuhan Tinggi, ITB Bandung, 78.
- Rufián-Henares, J. Á., Andrade, C. D. and Morales, F. J. (2009) *Non-Enzymatic Browning: The Case of the Maillard Reaction, Assessing the Generation and*

*Bioactivity of Neo-Formed Compounds in Thermally Treated Foods.*

- Sari, F., & Velioglu, Y. S. (2011). Effects of particle size, extraction time and temperature, and derivatization time on determination of theanine in tea. *Journal of Food Composition and Analysis*, 24(8), 1130-1135.
- Scharbert, S., Holzmann, N. and Hofmann, T. (2004) 'Identification of the astringent taste compounds in black tea infusions by combining instrumental analysis and human bioresponse', *Journal of Agricultural and Food Chemistry*, 52(11), pp. 3498–3508. doi: 10.1021/jf049802u.
- Schuh, C., & Schieberle, P. (2006). Characterization of the key aroma compounds in the beverage prepared from Darjeeling black tea: quantitative differences between tea leaves and infusion. *Journal of agricultural and food chemistry*, 54(3), 916-924.
- Selmar, D., Bytof, G., Knopp, S. E., & Breitenstein, B. (2006). Germination of coffee seeds and its significance for coffee quality. *Plant Biology*, 8(02), 260-264.
- Shukla, Y. (2007) 'Tea and cancer chemoprevention: A comprehensive review', *Asian Pacific Journal of Cancer Prevention*, 8(2), pp. 155–166.
- Su, Y. L., Leung, L. K., Huang, Y., & Chen, Z. Y. (2003). Stability of tea theaflavins and catechins. *Food Chemistry*, 83(2), 189-195.
- Sunarharum, W. B., Williams, D. J. and Smyth, H. E. (2014) 'Complexity of coffee flavor: A compositional and sensory perspective', *Food Research International*, 62, pp. 315–325. doi: 10.1016/j.foodres.2014.02.030.
- Sundari, D., Nuratmi, B., & Winarno, M. (2009). Toksisitas Akut (LD50) dan Uji Gelagat Ekstrak Daun Teh Hijau (*Camellia Sinensis* (Linn.) Kunze) Pada Mencit.
- Supartono, W., Sukartiko, A. C., Yuliando, H., & Kristanti, N. E. (2015). Possibility of some indigenous spices as flavor agent of green tea. *Agriculture and Agricultural Science Procedia*, 3, 62-66.
- Susanti, R. and Yuniastuti, A. (2012) 'Efektivitas Ekstrak Tanin Seledri (*Apium graveolens* L) Terhadap Profil Lipid Tikus Putih Hiperkolesterolemi.', *Life Science*, 1(2).
- Syah, Andi Nur Alam. *Taklukkan penyakit dengan teh hijau*. AgroMedia, 2006.
- Teranishi, Y., Mizutani, H., Murata, M., Shimizu, M., & Matsushima, K. (1995). Increased spontaneous production of IL-8 in peripheral blood monocytes from the psoriatic patient: relation to focal infection and response to treatments. *Journal of Dermatological science*, 10(1), 8-15.
- The, C. E. (1993) 'cn. 47', pp. 47–48.
- Tsuge, S., Ohtani, H., & Watanabe, C. (2011). *Pyrolysis-GC/MS data book of synthetic polymers: pyrograms, thermograms and MS of pyrolyzates*. Elsevier.

- Tsushida, T. and Takeo, T. (1984) 'Occurrence of Theanine in *Camellia Japonica* and *Camellia sasanqua* Seedlings', *Agricultural and Biological Chemistry*, 48(11), pp. 2861–2862. doi: 10.1271/bbb1961.48.2861.
- Tuminah, S., 2004., Teh Sebagai Salah Satu Sumber Antioksidan. Cermin Dunia Kedokteran. No 144 tahun 2004. pp: 52-54.
- Wahyono, S., Widiyastuti, Y., Irianti, S., Juniman, J., & Suharto, D. (2000). Laporan Penelitian Parameter Identifikasi Beberapa Ekstrak Tanaman Obat Secara Fisik dan Kimia.
- Wang, L.F., Park, S.C., Chung, J.O., Baik, J.H., and Park, S. K. (2004) 'to the Greenness of Green Tea', *Sensory and Nutritive Qualities of Food*, 69(8), pp. 301–305.
- Wang, L. *et al.* (2019) 'Protective effect of green tea catechin against urban fine dust particle-induced skin aging by regulation of NF-KB, AP-1, and MAPKs signaling pathways', *Environmental Pollution*, 252, pp. 1318–1324. doi: 10.1016/j.envpol.2019.06.029.
- Willis, S., Verghese, M., McCollum, M., Cheatom, K. Willis, Z., Seay, K., Sunkara, R. and Walker, L. (2017) 'A Comparison of Selected Phytochemical and Antioxidant Potential of Two Tea Beverages', *Food and Nutrition Sciences*, 8, pp. 1039–1049. doi: 10.4236/fns.2017.811076.
- Winarsi, H. (2007). Antioksidan alami & radikal bebas.
- Wong, K. H., Abdul Aziz, S. and Mohamed, S. (2008) 'Sensory aroma from Maillard reaction of individual and combinations of amino acids with glucose in acidic conditions', *International Journal of Food Science and Technology*, 43(9), pp. 1512–1519. doi: 10.1111/j.1365-2621.2006.01445.x.
- Wu, L. Y. *et al.* (2010) 'Analysis of chemical composition of *Chrysanthemum indicum* flowers by GC/MS and HPLC', *Journal of Medicinal Plants Research*, 4(5), pp. 421–426.
- Wu, Y. *et al.* (2016) 'Study of characteristic aroma components of baked Wujiatai green tea by HS-SPME/GC-MS combined with principal component analysis', *CYTA - Journal of Food*, 14(3), pp. 423–432. doi: 10.1080/19476337.2015.1123298.
- Yi, T., Zhu, L., Peng, W. L., He, X. C., Chen, H. L., Li, J., ... & Chen, H. B. (2015). Comparison of ten major constituents in seven types of processed tea using HPLC-DAD-MS followed by principal component and hierarchical cluster analysis. *LWT-Food Science and Technology*, 62(1), 194-201.
- Zhang, L. *et al.* (2019) 'Chemistry and Biological Activities of Processed *Camellia sinensis* Teas: A Comprehensive Review', *Comprehensive Reviews in Food Science and Food Safety*, 18(5), pp. 1474–1495. doi: 10.1111/1541-4337.12479.
- Zhang, M. *et al.* (2020) 'Contribution of addition theanine/sucrose on the formation of chestnut-like aroma of green tea', *Lwt*, 129(April), p. 109512. doi:

10.1016/j.lwt.2020.109512.