



## DAFTAR PUSTAKA

- Adams-Chapman and Stoll. (2006) ‘Neonatal infection and long-term neurodevelopmental outcome in the preterm infant.’, *Curr Opin Infect Dis*, 19 (3), pp. 290–7.
- Agrawal, Deorari, and Paul. (2014) ‘Sepsis in the Newborn’, *Division of Neonatology Department of Pediatrics All India Institute of Medical Sciences Ansari Nagar, New Delhi*.
- Astri Pinilih, H. ranto (2019) ‘Profil dan Faktor Risiko Kematian Sepsis Neonatus di RSUD Sukadana Lampung Timur’, 38(4), pp. 1–10.
- Bauer PR, Kashyap R, League SC, Park JG, Block DR, B. and NA, *et al.* (2016) ‘Diagnostic accuracy and clinical relevance of an inflammatory biomarker panel for sepsis in adult critically ill patients’, *Diagn Microbiol Infect Dis*, 84(2), pp. 175–80.
- Boeddhha, Navin P., Kerklaan, Dorian., Dunbar, Albert., van Puffelen, Esther., Nagtzaam, Nicole M.A., Vanhorebeek, Ilse, *et al.* (2018) ‘HLA-DR Expression on Monocyte Subsets in Critically Ill Children’, *The Pediatric infectious disease journal*, 37(10), pp. 1034–1040. doi: 10.1097/INF.0000000000001990.
- Brooks Z.C (2001) ‘Performance-Driven Quality Control’, *American Association for Clinical Chemistry*.
- Cheron, Floccard, Allaouchiche, Bernard, Caroline, Poitevin, Françoise, *et al.* (2010) ‘Lack of recovery in monocyte human leukocyte antigen-DR expression is independently associated with the development of sepsis after major trauma’, *Critical Care*, 14(6), p. 208. doi: 10.1186/cc9331.
- Craig and Mai. (2009) ‘Neutrophil Recruitment to the Lungs during bacterial Pneumonia’, *Infection and Immunity. Am Society for Microbiol*, 77 No 2, pp. 568–75.
- Dai, Jiang, Wenjie, Zhigang, Jian, Yongfei, Tieliang *et al.* (2017) ‘Neutrophil CD64 as a diagnostic marker for neonatal sepsis: Meta-analysis’, *Advances in Clinical and Experimental Medicine*, 26(2), pp. 327–332. doi: 10.17219/acem/58782.
- Davey, Morgan, Matt Liuzzi, Anna Tyler, Christopher Khan, Mohd Szakmany, Tamas Hall, Judith *et al.* (2014) ‘Microbe-Specific Unconventional T Cells Induce Human Neutrophil Differentiation into Antigen Cross-Presenting Cells’, *The Journal of Immunology*, 193(7), pp. 3704–3716. doi: 10.4049/jimmunol.1401018.
- Dhlamini, Suchard, Wiggill, Fadahun, Ballo. (2013) ‘Neutrophil CD64 has a high negative predictive value for exclusion of neonatal sepsis’, *SAJCH South African Journal of Child Health*, 7(1), pp. 25–29. doi: 10.7196/SAJCH.482.
- Dimoula, Pradier, Olivier, Zaina, Dyanne, Hulya Vincent, Jean Louis *et al.* (2014) ‘Serial determinations of neutrophil CD64 expression for the diagnosis and monitoring of sepsis in critically ill patients’, *Clinical Infectious Diseases*, 58(6), pp. 820–829. doi: 10.1093/cid/cit936.
- Djordjevic D, Pejovic J, Surbatovic M, Jevdjic J, Radakovic S. (2015) ‘Prognostic value and daily trend of interleukin-6, neutrophil CD64 expression, C-reactive protein and lipopolysaccharide-binding protein in critically ill patients: Reliable predictors of outcome or not?’, *J Med Biochem*, 34(4), pp. 431–9.
- Dokter, I. and Indonesia, A. (2016) ‘Diagnosis dan Tata Laksana Sepsis pada Anak’.



- Faix, J. (2011) 'Established and novel biomarkers of sepsis', *Biomarkers in Medicine*, 5(2), pp. 117–130.
- Fang, Fan, Cong Hai Li, Juan An, Qi Yao, Hong Ji, Qiang *et al.* (2015) 'Ratios of CD64 expressed on neutrophils, monocytes, and lymphocytes may be a novel method for diagnosis of neonatal sepsis', *Journal of Infection in Developing Countries*, 9(2), pp. 175–181. doi: 10.3855/jidc.4992.
- Genel F, Atlihan F, Ozsu E. (2010) 'Monocyte HLA-DR expression as predictor of poor outcome in neonates with late onset neonatal sepsis.', *J Infect [Internet]*, 60(3), pp. 224–8. Available at: <http://dx.doi.org/10.1016/j.jinf.2009.12.004>.
- Glaser, M.A., Hughes, L.M., Jnah, A., Newberry, D. & Harris-Haman, P. (2020) 'Neonatal Sepsis: A Review of Pathophysiology and Current Management Strategies', *Advances in Neonatal Care*, 00(0), pp. 1–12.
- Hotchkiss, Monneret, Guillaume, Didier Zhuang, Yugang Peng, Hu Chen, Yanqing Yuanzhuo *et al.* (2013) 'Sepsis-induced immunosuppression: from cellular dysfunctions to immunotherapy', *Angewandte Chemie International Edition*, 6(11), 951–952. Nature Publishing Group, 37(10)(8), pp. 2746–52. doi: 10.1007/s00134-003-1686-2.
- Ikatan Dokter Anak Indonesia (2009) 'Pedoman Pelayanan Medis Ikatan Dokter Anak Indonesia', in *Pedoman Pelayanan Medis Ikatan Dokter Anak Indonesia*. Jakarta: Badan Penerbit IDAI.
- James Wynn (2016) 'Defining neonatal sepsis HHS Public access', *Curr Opin Paediatr*, 28(2), pp. 135–140. doi: 10.1097/MOP.0000000000000315.
- Justin E. Juskewitch, R. S. A., , Stacy C. League2, Sarah M. Jenkins, C. and Y. Smith3, Felicity T. Enders, Stefan K. Grebe2, William A. Carey, and W. C. H. (2016) 'Monocyte HLA-DR Expression and Neutrophil CD64 Expression as Biomarkers of Infection in Critically Ill Neonates and Infants', *Pediatr Res*, 78(6), pp. 683–690. doi: 10.1038/pr.2015.164.MONOCYTE.
- Kan, B., Razzaghian, H.R. & Lavoie, P. (2016) 'An Immunological Perspective on Neonatal Sepsis', *Trends in Molecular Medicine*, 22(4), pp. 290–302.
- Kermorvant-Duchemin E, Laborie S, Rabilloud M, Lapillonne A. (2008) 'Outcome and prognostic factors in neonates with septic shock', *Pediatr Crit Care Med*, 9 (2), pp. 186–91.
- Khaertynov, Anokhin, Vladimir, Ilshat, Albert A, Sergey A, Asiya, Sofronova, Svetlana A. (2018) 'Changes in HLA-DR Expression on Monocytes and Lymphocytes in Neonatal Sepsis', *BioNanoScience*. BioNanoScience, 8(2), pp. 647–653. doi: 10.1007/s12668-018-0519-2.
- Lawn JE, Cousens S. (2005) 'Lancet neonatal survival steering team', *million neonatal deaths: When? Where? Why?*, 365, pp. 891–900.
- Leal, Álvarez-Nemegyei, José, Juan Rosado-Quiab, Ulises Diego-Rodríguez, Nidia, Paz-Baeza *et al.* (2012) 'Risk factors and prognosis for neonatal sepsis in southeastern Mexico: analysis of a four-year historic cohort follow-up', *BMC Pregnancy and Childbirth*, 12, pp. 1–9. doi: 10.1186/1471-2393-12-48.
- Leijte, Rimmelé, Thomas, Matthijs, Niklas, Céline, Gossez, Morgane *et al.* (2020) 'Monocytic HLA-DR expression kinetics in septic shock patients with different pathogens, sites of infection and adverse outcomes', *Critical Care. Critical Care*, 24(1), pp. 1–9. doi: 10.1186/s13054-020-2830-x.
- Liu, Johnson, Hope Cousens, Simon Perin, Jamie Scott, Susana Lawn, Joy E *et al.* (2012) 'Global, regional, and national causes of child mortality: An updated systematic analysis for 2010 with time trends since 2000', *The Lancet*. Elsevier



- Ltd, 379(9832), pp. 2151–2161. doi: 10.1016/S0140-6736(12)60560-1.
- Lukaszewicz AC, Grienay M, Resche-Rigon M, Pirracchio R, F. V and Boval B, *et al.* (2009) ‘Monocytic HLA-DR expression in intensive care patients: interest for prognosis and secondary infection prediction.’, *CritCare Med*, 37(10), pp. 2746–52.
- Mahmoodpoor, Paknezhad, Seyedpouya, Kamran, Hadi, Ali Akbar Sanaie, Sarvin *et al.* (2018) ‘Flow cytometry of CD64, HLA-DR, CD25, and TLRs for diagnosis and prognosis of sepsis in critically ill patients admitted to the intensive care unit: A review article’, *Anesthesiology and Pain Medicine*, 8(6), pp. 6–11. doi: 10.5812/aapm.83128.
- Marsh, S. G.; Albert, E. D.; Bodmer, W. F.; Bontrop, R. E.; Dupont, B.; Erlich, H. A.; Fernández-Viña, M.; Geraghty, D. E.; Holdsworth, R.; Hurley, C. K.; Lau, M.; Lee, K. W.; Mach, B.; Maiers, M.; Mayr, W. R.; Müller, C. R.; Parham, P.; Petersdorf, E. W., J. (2010) ‘Nomenclature for factors of the HLA system’, *Tissue Antigens*, 75(4), pp. 291–455.
- Martinez, F. O. and Gordon, S. (2014) ‘The M1 and M2 paradigm of macrophage activation: Time for reassessment’, *F1000Prime Reports*, 6(March), pp. 1–13. doi: 10.12703/P6-13.
- Miyake, F., Ishii, M., Hoshina, T., Ichikawa, S., Araki, S., Kinjo, T., Shibata, E., Hachisuga, T. & Kusuvara, K. (2016) ‘Analysis of the Physiological Variation in Neutrophil CD64 Expression during the Early Neonatal Period’, *American Journal of Perinatology*, 33 (14), pp. 1377–1381.
- Monneret G, Lepape A, Voirin N, Bohé J, Venet F, Debard AL, T. and H, Bienvenu J, Gueyffier F, V. P. (2006) ‘Persisting low monocyte human leukocyte antigen-DR expression predicts mortality in septic shock’, *Intensive Care Med*, 32, pp. 1175–1183.
- Motta M, Zini A, Regazzoli A, Garzoli E, Chirico G, Caimi L, C. and M. (2014) ‘Diagnostic accuracy and prognostic value of the cd64 index in very low birth weight neonates as a marker of early-onset sepsis’, *Scand J Infect Dis*, 46, pp. 433–439.
- Motta, Zini, Alice, Antonio, Elena, Gaetano, Luigi *et al.* (2014) ‘Diagnostic accuracy and prognostic value of the CD64 index in very low birth weight neonates as a marker of early-onset sepsis’, *Scandinavian Journal of Infectious Diseases*, 46(6), pp. 433–439. doi: 10.3109/00365548.2014.896028.
- Murthy, Godinho, Anthony, Vasudeva, Leslie, Sreekumaran. (2019) ‘Risk factors of neonatal sepsis in India: A systematic review and meta-analysis’, *PLoS ONE*, 14(4), pp. 1–26. doi: 10.1371/journal.pone.0215683.
- Odabasi, I. O. and Bulbul, A. (2020) ‘Review Neonatal Sepsis’, *Sisli Etfal Hastan Tip Bul.*, 54(2), pp. 142–158. doi: 10.14744/SEMB.2020.00236.
- Ten Oever, J., Netea, M.G. & Kullberg, B. J. (2016) ‘Utility of immune response-derived biomarkers in the differential diagnosis of inflammatory disorders’, *Journal of Infection*, 72(1), pp. 1–18.
- Oza, Lawn, Joy E Hogan, Daniel R Mathers, Colin Cousens, Simon N. (2015) ‘Neonatal cause-of-death estimates for the early and late neonatal periods for 194 countries: 2000–2013’, *Bulletin of the World Health Organization*, 93(1), pp. 19–28. doi: 10.2471/blt.14.139790.
- Palojärvi, Anniina Petäjä, Jari Siitonens, Sanna Janér, Cecilia Andersson, Sture. (2013) ‘Low monocyte HLA-DR expression as an indicator of immunodepression in very low birth weight infants’, *Pediatric Research*, 73(4),



- pp. 469–475. doi: 10.1038/pr.2012.199.
- Paolucci, M., Landini, M. P. and Sambri, V. (2012) ‘How Can the Microbiologist Help in Diagnosing Neonatal Sepsis ?’, 2012(di). doi: 10.1155/2012/120139.
- Pediatrics, A. A. (2018) ‘Group B streptococcal infections’, *Red Book: 2018 Report of The Committee on Infectious Diseases.*, American A(In D. Kimberkin, M. Brady, M. Jackson, S. Long, eds.).
- Perry, Sara E. Mostafa, Sobhy M. Wenstone, Richard Shenkin, Alan McLaughlin, Paul J.. (2003) ‘Is low monocyte HLA-DR expression helpful to predict outcome in severe sepsis?’, *Intensive Care Medicine*, 29(8), pp. 1245–1252. doi: 10.1007/s00134-003-1686-2.
- Polin R. (2012) ‘Management of neonates with suspected or proven early-onset bacterial sepsis.’, *Committee on Fetus and Newborn*, 129, pp. 1006–15.
- Pradhan, Richeek Jain, Paresh Paria, Anshuman Saha, Anindya Sahoo, Jagdish Sen, Anway et al. (2016) ‘Ratio of neutrophilic CD64 and monocytic HLA-DR: A novel parameter in diagnosis and prognostication of neonatal sepsis’, *Cytometry Part B - Clinical Cytometry*, 90(3), pp. 295–302. doi: 10.1002/cyto.b.21244.
- Puopolo KM, Draper D, Wi S, Newman TB, Zupancic J, Lieberman E, et al (2011) ‘Estimating the probability of neonatal early-on\_set infection on the basis of maternal risk factors’, *Pediatrics and Neonatology*, 128, pp. 1155–63.
- Pusponegoro, T. S. (2016) ‘Sepsis pada Neonatus (Sepsis Neonatal)’, *Sari Pediatri*, 2(2), p. 96. doi: 10.14238/sp2.2.2000.96-102.
- Rajkumar Motiram Meshram, Vishal S Gajimwar, S. (2019) ‘Predictors of mortality in outborns with neonatal sepsis: A prospective observational study’, *Niger Postgrad Med J*, 26(4), pp. 216–222. doi: 10.4103/npmj.npmj\_91\_19.
- Reis Machado, Juliana Soave, Danilo Figueiredo Da Silva, Marcos Vinícius De Menezes, Liliana Borges
- Etchebehere, Renata Margarida Monteiro, Maria Luiza Gonçalves Dos Reis et al. (2014) ‘Neonatal sepsis and inflammatory mediators’, *Mediators of Inflammation*, 2014. doi: 10.1155/2014/269681.
- Sabeeh Jumah, D., Kadhum Hassan, ad and Mea, A. (2007) ‘Predictors of Mortality Outcome in Neonatal Sepsis’, *The Medical Journal of Basrah University*, 25(1), pp. 11–18. doi: 10.33762/mjbu.2007.48118.
- Sarode, R., Ingole, N., Jasani, B., Nataraj, G., Nanavati, R. & Mehta, P. (2017) ‘Role of CD64 in the diagnosis of neonatal sepsis’, *International Journal of Contemporary Medical Research*, 4(9), pp. 1959–1963.
- Satar M, Arısoy AE, Ç. İ. (2018) *Turkish Neonatology Society Neonatal Guide, Diagnosis and Treatment of Infections*.
- Shane AL (2017) ‘Neonatal sepsis’, *Lancet*, 390, pp. 1770–80.
- Sheikh A, Javed T Afzal MF Sheikh CA. (2011) ‘Course and Complications of Early Onset Nonnatal Sepsis : A Descriptive Study’, *Annals*, 16(4).
- Simonsen KA, Anderson-Berry AL, Delair SF, D. H. (2014) ‘Early-onset neonatal sepsis’, *Clin Microbiol Rev*, 27, pp. 21–47.
- Skirecki T, Mikaszewska-Sokolewicz M, Hoser G, Z.-B. (2016) ‘The early expression of HLA-DR and CD64 myeloid markers is specifically compartmentalized in the blood and lungs of pa\_tients with septic shock’, *Mediators Inflamm*.
- Song SH, Kim HK, Park MH, C. H. (2008) ‘Neutrophil CD64 expression is associated with severity and prognosis of disseminated intravascular coagulation.’, *Thromb Res*, 121, pp. 499–507.



- Tam, P.-Y. I. dan Bendel, C. M. (2017) 'Diagnostics for neonatal sepsis: Current approaches and future directions', *Pediatric Research*, 82(4), pp. 574–583. doi: 10.1038/pr.2017.134.
- Tang, Z., Qin, D., Tao, M., Lv, K., Chen, S., Zhu, X., Li, X., Chen, T., Zhang, M., Zhong, M., Yang, H., Xu, Y. & Mao, S. (2018) 'Examining the utility of the CD64 index compared with other conventional indices for early diagnosis of neonatal infection.', *Nature*, 8(1), pp. 1–7.
- Vester, Helen Dargatz, P. Huber-Wagner, S. Biberthaler, P. Van Griensven, M. (2015) 'HLA-DR expression on monocytes is decreased in polytraumatized patients', *European Journal of Medical Research*. BioMed Central, 20(1), pp. 1–9. doi: 10.1186/s40001-015-0180-y.
- W.E. Benitz (2010) 'Adjunct laboratory test in the diagnosis of early-onset neonatal sepsis', *Clinics in Perinatology*, 37 no 2, pp. 421–438.
- Wang, Y. and Jönsson, F. (2019) 'Expression, role, and regulation of neutrophil Fc $\gamma$  receptors', *Frontiers in Immunology*, 10(AUG), pp. 1–13. doi: 10.3389/fimmu.2019.01958.
- Ward, N. S., Casserly, B. and Ayala, A. (2008) 'The Compensatory Anti-inflammatory Response Syndrome (CARS) in Critically Ill Patients', *Clinics in Chest Medicine*, 29(4), pp. 617–625. doi: 10.1016/j.ccm.2008.06.010.
- Wibowo, T., Haksari, E.L., Wandita, S., Anggraini, A. & Safrida, E. N. (2019) 'PPK Sepsis Neonatorum Awitan Dini', pp. 1–8.
- Winkler, Martin Sebastian Rissiek, Anne Priefler, Marion Schwedhelm, Edzard Robbe, Linda Bauer, Antonia et al. (2017) 'Human leucocyte antigen (HLA-DR) gene expression is reduced in sepsis and correlates with impaired TNF $\alpha$  response: A diagnostic tool for immunosuppression?', *PLoS ONE*, 12(8), pp. 1–14. doi: 10.1371/journal.pone.0182427.
- World Health Organization (WHO) (2015) 'Region Sepsis and Other Infectious Conditions of Newborn', *Global Health Observatory data repository*.
- Wynn, J., Wong, H., Shanley, T., Bizzarro, M., Saiman, L. & Polin, R. (2014) 'Time for a neonatal – specific consensus definition for sepsis', *Pediatric Critical care Medicine*, 15(6), pp. 523–528. doi: 10.1097/PCC.0000000000000157.Time.
- Wynn, J. L. dan Wong, H. R. (2020) 'Pathophysiology of Neonatal Sepsis', *Pathophysiology of Neonatal Diseases*, pp. 1536–1552.
- Yusef, D. et al. (2018) 'Clinical characteristics and epidemiology of sepsis in the neonatal intensive care unit in the era of multi-drug resistant organisms: A retrospective review', *Pediatrics and Neonatology*. Elsevier Taiwan LLC, 59(1), pp. 35–41. doi: 10.1016/j.pedneo.2017.06.001.
- Zea-Vera, A. and Ochoa, T. J. (2015) 'Challenges in the diagnosis and management of neonatal sepsis', *Journal of Tropical Pediatrics*, 61(1), pp. 1–13. doi: 10.1093/tropej/fmu079.
- Zhuang, Peng, Hu Chen, Yuanzhuo Zhou, Shuqin Chen, Yanqing (2017) 'Dynamic monitoring of monocyte HLA-DR expression for the diagnosis, prognosis, and prediction of sepsis', *Frontiers in Bioscience - Landmark*, 22(8), pp. 1344–1354. doi: 10.2741/4547.