

DAFTAR PUSTAKA

- Ansharullah., Ibrahim M.N. & Wiranty, E. 2018. Karakteristik fisikokimia dan organoleptik surimi berbasis ikan gabus – tepung sagu pada penyimpanan dingin. REKA PANGAN. Vol. 12, Nomor 1.
- Asikin, A.N. & Kusumaningrum, I. 2018. Karakteristik protein ikan gabus berdasarkan ukuran berat ikan asal das Mahakan Kalimantan Timur. Jurnal Pengolahan Hasil Perikanan Indonesia 21(1):137.
- Baehaki, A., I. Widiastuti, Nainggolan, C. & Gofar, N. 2020. Antioxidant activities of snakehead (*channa striata*) fish skin : peptides hydrolysis using protease TP2 isolate from swamp plant silage. Potravinarstvo Slovak Journal of Food Sciences.
- Barkia, A., Bougatef, A., Khaled, H.B & Nasri, M. 2009. Antioxidant activities of sardinella heads and or viscera protein hydrolysates prepared by enzymatic treatment. Journal of Food Biochemistry. 34; 3003-320.
- Benjakul, S., Yarnpakdee, S., Senphan, T., Halldorsdottir, S.M. & Kristinsson, H.G. 2013. Fish protein hydrolysates: production, bioactivities, and applications. Chapter 9.
- Botterweck, AAM, Vergaen H, GoldBohm RA, KleinJans J & van den Brant PA. 2007. Intake of butylated hydroxyanisole and butylated hydroxytoluene and stomach cancer risk: results from analyses in the Netherlands cohort study. Food and Chemical Toxicology. 38 (7): 599–605.
- Cahyaningrum, K., Amir, H., Siti Ari, B. 2016. Aktivitas antioksidan ekstrak rumput laut cokelat (*Sargassum polycystum*). AGRITECH, Vol. 36, No. 2.
- Chasanah, E., Mala Nurilmala, Ayu Ratih, P, Diini, F. 2015. Komposisi Kimia, Kadar Albumin Dan Bioaktivitas Ekstrak Protein Ikan Gabus (*Channa Striata*) Alam Dan Hasil Budidaya. JPB Kelautan Dan Perikanan Vol. 10 No. 2 Tahun 2015: 123±132.
- Choonpicharn, S., Jaturasitha, S., Rakariyatham, N., Suree, N & Niamsup, H. 2014. Antioxidant and antihypertensive activity of gelatin hydrolysate from Nile tilapia skin. Food Science and Technology, 52(5), 3134–3139. doi:10.1007/s13197-014-1581-6. Coligan J.E., Dunn, B.M, Speicher D.W. & Wingfield, P.T. 2002. Current Protocols in Protein Science. John Wiley & Sons, Inc.
- Dekkers, E., Raghavan, S., Kristinsson, H.G. & Marshall, M.R. 2011. Oxidative stability of mahi mahi red muscle dipped in tilapia protein hydrolysates. Food Chemistry. 124, 640–645.
- Elavarasan K., Naveen Kumar V and Shamasundar BA. 2013. Antioxidant and Functional Properties of Fish Protein Hydrolysates from Fresh Water Carp (*Catla catla*) as Influenced by the Nature of Enzyme. Journal of Food Processing and Preservation, 38(3), 1207–1214. doi:10.1111/jfpp.12081.
- Gajanan, P. G., Elavarasan, K., & Shamasundar, B. A. 2016. Bioactive and functional properties of protein hydrolysates from fish frame processing waste using plant proteases. Environmental Science and Pollution Research, 23(24), 24901–24911. doi:10.1007/s11356-016-7618-9.
- Gbogouri, G.A., Linder, M., Fanni, J. & Parmentier, M. 2004. Influence of hydrolysis degree on the functional properties of salmon byproduct hydrolysates. Journal of Food Science. 69, 615–622.
- Hariyati, M. Mahendradatta., A.B. Tawali. & J. Langkon. 2020. Enzymatic hydrolysis of proteins from snakehead-fish (*Channa striata*). Materials Science and Engineering. 885- 012015.
- Halliwell B. 2007. Biochemistry of Oxidative Stress Biochemical Society Transactions, 35: 1147-1150.
- Haslaniza, H., M.Y. Maskat, A.W.M. Wan & S. Mamot. 2010. The effects of enzyme concentration, temperature and incubation time on nitrogen content and degree of

- hydrolysis of protein precipitate from cockle (*Anadara granosa*) meat wash water. International Food Research Journal. 17: 147-152.
- Kahl, R. & Kappus, H., 1993. Toxicology of the synthetic antioxidants BHA and BHT in comparison with the natural antioxidant vitamin E. Zeitschrift fur Lebensmittel-untersuchung und -forschung 196 (4), 329–338.
- KKP. 2020. Produksi Perikanan. Diakses pada tanggal 25 Februari 2022 (https://statistik.kkp.go.id/home.php?m=prod_ikan_prov&i=2#panel-footer).
- Kristinsson H.G & Rasco, B.A .2000. Biochemical and functional properties of Atlantic salmon (*Salmo salar*) muscle proteins hydrolyzed with various alkaline proteases. J Agric Food Chem. 48:657–666.
- Le Vo, T.D., Nguyen, T.T.H., Phan, D.V., Nguyen, H.D.M. & Tran, H.Q. 2016. Investigation of antioxidant activity of the hydrolysate derived from Tra catfish byproducts using Alcalase 2.4 L FG for application as a natural antioxidant ingredient. Journal Science and Technology Development 19(6):109–121 DOI 10.32508/stdj.v19i3.577.
- Mutamimah, D., Ibrahim, B., & Trilaksani, W. 2018. Antioxidant activity of protein hydrolysate produced from tuna eye (*Thunnus* sp.) by enzymatic hydrolysis. Jurnal Pengolahan Hasil Perikanan Indonesia. 21(3): 522-531.
- Nićiforović, N., Mihailović, V., Mašković, P., Solujić, S., Stojković, A., & Muratspahić, D. P. (2010). Antioxidant activity of selected plant species; potential new sources of natural antioxidants. Food and Chemical Toxicology, 48(11), 3125–3130. doi:10.1016/j.fct.2010.08.007.
- Noman A, Xu Y, AL-Bukhaiti, WQ, Abed SM, Ali AH, Ramadhan AH and Xia W. 2018. Influence of Enzymatic Hydrolysis Conditions On The Degree of Hydrolysis and Functional Properties of Protein Hydrolysate Obtained from Chinese Sturgeon (*Acipenser sinensis*) by Using Papain Enzyme. Process Biochemistry 67, 19–28. doi.org/10.1016/j.procbio.2018.01.009.
- Puspawati, N.M., P.P. Dewi, N.W. Bogoriani & N.K. Ariati. Produksi hidrolisat protein antioksidan melalui hidrolisis enzimatis protein kulit ayam broiler dengan enzim papain.
- Rabiei, S., Rezaei, M., Asgharzade, S., Nikoo, M. & Kopai, M.R. 2019. Antioxidant and cytotoxic properties of protein hydrolysates obtained from enzymatic hydrolysis of Klunzinger's mullet (*Liza klunzingeri*) muscle. Braz. J. Pharm. Sci. 55:e18304. Jurnal Kimia 14 (2) p-ISSN 1907-9850.
- Ramakrishnan VV, Ghaly AE, Brooks, Budge SM. 2013. Extraction of proteins from mackerel fish processing waste using alcalase enzyme. Bioprocessing and Biotechniques. 3(2): 1–9.
- Sivaramakrishnan, T., Swain, S., Saravanan, K., Sankar. R, Kiruba, Roy, S. D., Biswas, L., & Shalini, B. 2017. Turkish Journal of Fisheries and Aquatic Sciences, 17(3). doi:10.4194/1303-2712-v17_3_20.
- Sobri, A., Herpandi, Lestari, S. 2017. Uji Pengaruh Suhu Pengeringan pada Karakteristik Kimia dan Sensori Kaldu Bubuk Kepala Ikan Gabus (*Channa striata*). Fishtech – Jurnal Teknologi Hasil Perikanan. Vol. 6, No.2: 97-106.
- Souissi, N., Bougatef, Ali., Ellouz, Y.T., & Nasri, M. 2007. Biochemical and Functional Properties of Sardinella (*Sardinella aurita*) By-Product Hydrolysates. Food Technology and. Biotechnology. 45 (2) 187–194.
- Ubadillah & Hersoelistyorini. 2010. Kadar Protein dan Sifat Organoleptik Nugget Rajungan dengan Substitusi Ikan Lele. Jurnal Pangan dan Gizi Vol 01 No. 02.
- Wasswa, J., Tang, J., Gu, X., & Yuan, X. 2007. Influence of the extent of enzymatic hydrolysis on the functional properties of protein hydrolysate from grass carp (*Ctenopharyngodon idella*) skin. Food Chemistry. 104, 1698–1704.



UNIVERSITAS
GADJAH MADA

Aktivitas Antioksidan Hidrolisat Protein dari Hasil Samping Ikan Gabus (*Channa striata*) Yang Diproses Secara Enzimatis

VENNY AGUSTIN, Mgs. M. Prima Putra, S.Pi., M.Sc., Ph.D.; Dr. Amir Husni, S.Pi., M.P

Universitas Gadjah Mada, 2022 | Diunduh dari <http://etd.repository.ugm.ac.id/>

- Wu, H.C., Chen, H.M., & Shiau, C.Y. 2003. Free amino acids and peptides as related to antioxidant properties in protein hydrolysates of mackerel (*Scomber austriasicus*). Food Res Int. 36:949–957.
- Yuliani, Marwati., Wardana, H., Emmawati, A. & Candra, K.P. 2018. Karakteristik kerupuk ikan dengan substitusi tepung tulang ikan gabus (*Channa striata*) sebagai fortifikasi kalsium. .Volume 21 Nomor 2.

