

DAFTAR PUSTAKA

- Amin, A.P., Bach, R.G., Caruso, M.L., Kennedy, K.F., Spertus, J.A. 2017. Association of Variation in Contrast Volume With Acute Kidney Injury in Patients Undergoing Percutaneous Coronary Intervention. *JAMA Cardiol.* 2: 1007–1012.
- Andreucci, M., Faga, T., Pisani, A., Sabbatini, M., *et al.* 2014. The Choice of the Iodinated Radiographic Contrast Media to Prevent Contrast-Induced Nephropathy. *Advances in Nephrology.* 11 pages.
- Azzalini, L., Laricchia, A., Regazzoli, D., Mitomo, S., *et al.* 2019. Ultra-Low Contrast Percutaneous Coronary Intervention to Minimize the Risk for Contrast-Induced Acute Kidney Injury in Patients With Severe Chronic Kidney Disease. *J. Invasive Cardiol.* 31: 176–182.
- Barbieri, L., Verdoia, M., Nardin, M., Marino, P., Suryapranata, H., De Luca, G. 2016. Gender Difference in the Risk of Contrast-Induced Nephropathy in Patients Undergoing Coronary Angiography or Percutaneous Coronary Intervention. *Angiology.* 68: 542–546.
- Barrett, B.J., Parfrey, P.S., 2006. Clinical practice. Preventing nephropathy induced by contrast medium. *N. Engl. J. Med.* 354: 379–386.
- Campos, A.M., Placido-Sposito, A., Freitas, W.M., Moura, F.A., *et al.* 2016. ST-elevation myocardial infarction risk in the very elderly. *BBA Clin.* 6: 108–112.
- Chagnac, A., Herman, M., Zingerman, B., Erman, A., Rozen-Zvi, B., *et al.* 2008. Obesity-induced glomerular hyperfiltration: its involvement in the pathogenesis of tubular sodium reabsorption. *Nephrol. Dial. Transplant. Off. Publ. Eur. Dial. Transpl. Assoc. - Eur. Ren. Assoc.* 23: 3946–3952.
- Das, S.R., Alexander, K.P., Chen, A.Y., Powell-Wiley, T.M., *et al.* 2011. Impact of body weight and extreme obesity on the presentation, treatment, and in-hospital outcomes of 50,149 patients with ST-Segment elevation myocardial infarction results from the NCDR (National Cardiovascular Data Registry). *J. Am. Coll. Cardiol.* 58: 2642–2650
- Duraes, A.R., Bitar, Y.S., Freitas, A.C.T., Filho, I.M., Freitas, B.C., Fernandez, A.M. 2017. Gender differences in ST-elevation myocardial infarction (STEMI) time delays: experience of a public health service in Salvador-Brazil. *Am. J. Cardiovasc. Dis.* 7: 102–107.

- Evola, S., Lunetta, M., Macaione, F., Fonte, G., Milana, G., *et al.* 2012. Risk factors for contrast induced nephropathy: a study among Italian patients. *Indian Heart J.* 64: 484–491.
- Freeman RV, O'Donnell M, Share D *et al.* 2002. Nephropathy requiring dialysis after percutaneous coronary intervention and the critical role of an adjusted contrast dose. *Am J Cardiol.* 90:1068-1073.
- Giancarlo M. 2009. Contrast volume during primary percutaneous coronary intervention and subsequent contrast induced nephropathy and mortality. *Ann Intern Med.* 150:170–177.
- Heyman, S.N., Rosenberger, C., Rosen, S., Khamaisi, M. 2013. Why is diabetes mellitus a risk factor for contrast-induced nephropathy? *Biomed Res. Int.* 2013: 123589.
- Ibanez, B., James, S., Agewall, S., Antunes, M.J., *et al.* 2018. 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: The Task Force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Soci. *Eur Heart J.* 39: 119–177.
- Jain, T., Shah, S., Shah, J., Jacobsen, G., *et al.* 2018. Contrast-Induced Nephropathy in STEMI Patients With and Without Chronic Kidney Disease. *Crit Pathways in Cardiol.* 17: 25–31.
- Jaipaul, N., Manalo, R., Sadjadi, S.-A., McMillan, J. 2010. Obesity is not associated with contrast nephropathy. *Ther. Clin. Risk Manag.* 6: 213–217.
- Jessani, S., Levey, A.S., Bux, R., Inker, L., Islam, M., Chaturvedi, N., *et al.* 2014. Estimation of GFR in South Asians: a study from the general population in Pakistan. *American journal of kidney diseases.* 63(1):49–58.
- Kabeer, M.A., Cross, J., Hamilton, G., Rashid, S.T. 2021. Obesity as a Risk Factor for Radiographic Contrast-Induced Nephropathy. 72: 274–278.
- KDIGO. 2012. Clinical Practice Guideline for Acute Kidney Injury. *Journal of the international society of nephrology.* 2: 124-138.
- Kochar, A., Chen, A.Y., Sharma, P.P., Pagidipati, N.J., *et al.* 2018. Long-Term Mortality of Older Patients With Acute Myocardial Infarction Treated in US Clinical Practice. *J. Am. Heart Assoc.* 7: e007230.
- Kooiman, J., Seth, M., Share, D., Dixon, S., Gurm, H.S. 2014. The Association between Contrast Dose and Renal Complications Post PCI across the Continuum of Procedural Estimated Risk. *PLoS One* 9, e90233.

- Kumar, D., Liaquat, H., Sial, J.A., Saghir, T., *et al.* 2020. Risk Factors Associated With Contrast-Induced Nephropathy after Primary Percutaneous Coronary Intervention. *Cureus* 12: e9721–e9721.
- Laskey, W. K., Jenkins, C., Selzer, F., Marroquin, O., Wilensky, R.L., Glaser R, Cohen HA, Holmes Jr, D.R. 2007. Volume-to-creatinine clearance ratio: a pharmacokinetically based risk factor for prediction of early creatinine increase after percutaneous coronary intervention. *J Am CollCardiol.* 50:584-90.
- Liu, Y., Tan, N., Zhou, Y.L., *et al.* 2012. The contrast medium volume to estimated glomerular filtration rate ratio as a predictor of contrast-induced nephropathy after primary percutaneous coronary intervention. *Int Urol Nephrol.* 44:221–9.
- Mager, A., Assa, H.V., Lev, E., Bental, T., *et al.* 2011. The Ratio of Contrast Volume to Glomerular Filtration Rate Predicts Outcomes After Percutaneous Coronary Intervention for ST-Segment Elevation Acute Myocardial Infarction. *Catheterization and Cardiovascular Interventions.* 78:198–201.
- Marenzi, G., Lauri, G., Assanelli, E., *et al.* 2004. Contrast-Induced Nephropathy in Patients Undergoing Primary Angioplasty for Acute Myocardial Infarction. *J Am CollCardiol.* 9:1780–5.
- McCullough, P. A., Bertrand, M. E., Brinker, J. A., *et al.* 2006. A meta-analysis of the renal safety of isoosmolar iodixanol compared with low-osmolar contrast media. *Journal of the American College of Cardiology*, 48(4): 692–9.
- McCullough, P. A. 2008. Contrast-induced acute kidney injury. *J Am CollCardiol.* 51:1419-28.
- McCullough, P., Choi, J., Feghali, G., Schussler, J., *et al.* 2016. Contrast-Induced Acute Kidney Injury. *Journal of the American College of Cardiology.* 68:1465–73.
- Mehran, R., Nikolsky, E. 2006. Contrast-induced nephropathy: Definition, epidemiology, and patients at risk. *Kidney International.* 69: S11–S15.
- Mehran, R., Nikolsky, E., Kirtane, A.J., Caixeta, A., Wong, S.C., *et al.* 2009. Ionic Low-Osmolar Versus Nonionic Iso-Osmolar Contrast Media to Obviate Worsening Nephropathy After Angioplasty in Chronic Renal Failure Patients: The ICON (Ionic versus non-ionic Contrast to Obviate worsening Nephropathy after angioplasty in chronic renal failure patients) Study. *JACC Cardiovasc. Interv.* 2: 415–421.

- Mehran, R., Dangas, G., Weisbord, S. 2019. Contrast-Associated Acute Kidney Injury. *N Engl J Med*. 380:2146-55.
- Michels, W.M., Grootendorst, D.C., Verduijn, M., Elliott, E.G., Dekker, F.W., Krediet, R.T. 2010. Performance of the cockcroft-gault, MDRD, and New CKD-EPI formulas in relation to GFR, age, and body size. *Clin J Am Soc Nephrol*. 5(6):1003–9.
- Morcos, S. 2005. Prevention of Contrast Media–induced Nephrotoxicity after Angiographic Procedures. *J Vasc Interv Radiol*; 16:13–23.
- Morcos, R., Kucharik, M., Bansal, P., Taii, H., *et al.* 2019. Contrast-Induced Acute Kidney Injury:Review and Practical Update. *Clinical Medicine Insights: Cardiology*.13: 1-9.
- Neeland, I.J., Das, S.R., Simon, D.N., Diercks, D.B., Alexander, K.P., *et al.* 2017. The obesity paradox, extreme obesity, and long-term outcomes in older adults with ST-segment elevation myocardial infarction: results from the NCDR. *Eur. Hear. journal. Qual. care Clin*. 3: 183–191.
- Nozue, T., Michishita, I., Iwaki, T., Mizuguchi, I., *et al.* 2009. Contrast medium volume to estimated glomerularfiltration rate ratio as a predictor ofcontrast-induced nephropathy developing afterelective percutaneous coronary intervention. *Journal of Cardiology*. 54:214-220.
- Nyman, U., Almen, T., Aspelin, P., Hellstrom, M., *et al.* 2005. Contrast-Medium-Induced Nephropathy Correlated to the Ratio Between Dose in Gram Iodine and Estimated GFR in ml/min. *Acta Radiologica*.8: 13 pages.
- Palli, E., Makris, D., Papanikolaou, J., Garoufalis, G., Zakyntinos, E. 2014. Contrast-induced nephropathy in aged critically ill patients. *Oxid. Med. Cell. Longev*. 756469.
- Park, H.S., Kim, C. J., Yi, J.E., *et al.* 2015. Contrast Volume/Raw eGFR Ratio for Predicting Contrast-Induced Acute Kidney Injury in Patients Undergoing Percutaneous Coronary Intervention for Myocardial Infarction. *Cardiorenal medicine*. 5(1): 61–8.
- Rihal, C., Textor, S., Grill, D., *et al.* 2002. Incidence and prognostic importance of acute renal failure after PCI. *Circulation*. 105:2259–2264.
- Reed, M., Meier, P., Tamhane, U., Welch, K., *et al.* 2009. The relative renal safety of iodixanol compared with low-osmolar contrast media. A meta-analysis of randomized controlled trials. *Journal of the American College of Cardiology*. 7: 645–654.

- Schweiger, M., Chambers, C., Davidson, C., Zhang, S., *et al.* 2007. Prevention of Contrast Induced Nephropathy: Recommendations for the High Risk Patient Undergoing Cardiovascular Procedures. *Catheterization and Cardiovascular Interventions*. 69:135–140.
- Sherwin PF, Cambron R, Johnson JA, Pierro JA. 2005. Contrast dose-to-creatinine clearance ratio as a potential indicator of risk for radiocontrast-induced nephropathy: correlation of D/CrCL with area under the contrast concentration time curve using iodixanol. *Invest Radiol*. 40:598-603.
- Shoukat, S., Gowani, S.A., Jafferani, A., *et al.* 2010. Contrast-Induced Nephropathy in Patients Undergoing Percutaneous Coronary Intervention. *Cardiology Research and Practice*. 2010: 649164.
- Stevens LA, Coresh J, Greene T, Levey AS. 2006. Assessing kidney function-measured and estimated glomerular filtration rate. *N Engl J Med*. 354: 2473-83.
- Tan, N., Liu, Y., Zhou, Y.L., *et al.* 2012. Contrast Medium Volume to Creatinine Clearance Ratio: A Predictor of Contrast-Induced Nephropathy in the First 72 Hours Following Percutaneous Coronary Intervention. *Catheterization and Cardiovascular Interventions*. 79:70–75.
- Wang, K., Li, H.-L., Bei, W.-J., Guo, X.-S., *et al.* 2017. Association of left ventricular ejection fraction with contrast-induced nephropathy and mortality following coronary angiography or intervention in patients with heart failure. *Ther. Clin. Risk Manag*. 13: 887–895.
- Wang, J., Zhang, C., Liu, Z., Bai, Y. 2021. Risk factors of contrast-induced nephropathy after percutaneous coronary intervention: a retrospective analysis. *J. Int. Med. Res*. 49: 3000605211005972–3000605211005972.
- Yang, H.-Y., Huang, J.-H., Hsu, C.-Y., Chen, Y.-J. 2012. Gender Differences and the Trend in the Acute Myocardial Infarction: A 10-Year Nationwide Population-Based Analysis. *Sci. World J*. 184075.
- Zachura, M., Wilczek, K., Kurzawski, J., Gierlotka, M., Gąsior, M., Sadowski, M. 2018. Gender-related differences in men and women with ST-segment elevation myocardial infarction and incomplete infarct-related artery flow restoration: a multicenter national registry. *Adv. Interv. Cardiol*. 14: 356–362.
- Zhao, F., Lei, R., Yang, S.-K., Luo, M., Cheng, W., *et al.* 2019. Comparative effect of iso-osmolar versus low-osmolar contrast media on the incidence of contrast-induced acute kidney injury in diabetic patients: a systematic review and meta-analysis. *Cancer Imaging* 19, 38.