

DAFTAR PUSTAKA

- Ahbap E., Kara E., Basturk T., Sakaci T., Koc Y., Sahutoglu T., Akgol C., Sevinc M., Ucar Z.A., Unsal A., 2015, Neutrophil to Lymphosit and Platelet to Lymphocyte Ratio: A Novel Markers of Inflammation in Chronic Haemodialysis Patients (Abstract), Sisli Etfal Educational and Research Hospital, Istanbul, Turkey.
- Al-Othman, A. M., Al-Naseeb, A. J. M., Almajwal, A. M., Al-Mummar, M. N., Alhamdan, A. A., Al-Saif, M. A., & Alorf, S. M. (2016). Association of malnutrition in peritoneal dialysis patients of Saudi Arabia. *Arabian Journal of Chemistry*, 9, S1059–S1062. <https://doi.org/10.1016/j.arabjc.2011.11.015>
- Amparo, F. C., Kamimura, M. A., Molnar, M. Z., Cuppari, L., Lindholm, B., Amodeo, C., Carrero, J. J., & Cordeiro, A. C. (2015). Diagnostic validation and prognostic significance of the Malnutrition-Inflammation Score in nondialyzed chronic kidney disease patients. *Nephrology Dialysis Transplantation*, 30(5), 821–828. <https://doi.org/10.1093/ndt/gfu380>.
- Angkananard T., Anothaisintawee T., McEvoy M., Attia J., Thakkestia A., 2018, Neutrophil Lymphocyte Ratio and Cardiovascular Disease Risk: A Systematic Review and Meta-Analysis, *Hindawi BioMed Research International*, Article ID 2703518, 11 pages <https://doi.org/10.1155/2018/2703518>.
- An X., Mao H. P., Wei X., C J., Yang X., Li Z., Yu X., Li Z., 2012, Elevated neutrophil to lymphocyte ratio predicts overall and cardiovascular mortality in maintenance peritoneal dialysis patients, *Oct;44(5):1521-8*. doi: 10.1007/s11255-012-0130-3.
- Balafa, O., Halbesma, N., Struijk, D. G., Dekker, F. W., & Krediet, R. T. (2011). Peritoneal albumin and protein losses do not predict outcome in peritoneal dialysis patients. *Clinical Journal of the American Society of Nephrology*, 6(3), 561–566. <https://doi.org/10.2215/CJN.05540610>
- Baroni, G., Schuinski, A., De Moraes, T. P., Meyer, F., & Pecoits-Filho, R. (2012). Inflammation and the peritoneal membrane: Causes and impact on structure and function during peritoneal dialysis. *Mediators of Inflammation*, 2012(June 2014). <https://doi.org/10.1155/2012/912595>
- Bauer, J., Capra, S., & Ferguson, M. (2002). Use of the scored Patient-Generated Subjective Global Assessment (PG-SGA) as a nutrition assessment tool in patients with cancer. *European Journal of Clinical Nutrition*, 56(8), 779–785. <https://doi.org/10.1038/sj.ejcn.1601412>
- Bebas, P., Berakhir, Y., & Tindakan, D. (2019). *Majalah Kedokteran. The Journal of Medical School*, 52(2), 84–88.
- Bennett K and Aditya B. An Overview of Diabetic Nephropath: Epidemiology, Pathophysiology, and Treatment. *Journal of Diabetic Nursing*. 2015; 18: 61-7.

- Bharadwaj, S., Ginoya, S., Tandon, P., Gohel, T. D., Guirguis, J., Vallabh, H., Jevenn, A., & Hanouneh, I. (2016). Malnutrition: Laboratory markers vs nutritional assessment. *Gastroenterology Report*, 4 (4), 272–280. <https://doi.org/10.1093/gastro/gow013>.
- Bross R., Zitterkoph J., Pithia J., Benner D., Rambod M., Kovesdy C. P., Kopple J. D., Zadeh K. K., 2009, Association of Serum Total Iron-Binding Capacity and Its Changes Over Time with Nutritional and Clinical Outcomes in Hemodialysis Patients *Am J Nephrol* 2009;29:571–581.
- Chan, J. Y. W., Che, K. I., Lam, K. M. C., Chow, K. M., Chung, K. Y., Li, P. K. T., & Szeto, C. C. (2007). Comprehensive malnutrition inflammation score as a marker of nutritional status in Chinese peritoneal dialysis patients. *Nephrology*, 12(2), 130–134. <https://doi.org/10.1111/j.1440-1797.2006.00693.x>.
- Chen C., Zhao H. Y., Zhang Y. H., 2019, Correlation between neutrophil-to-lymphocyte ratio and kidney dysfunction in undiagnosed hypertensive population from general health checkup, *J Clin Hypertens*; 22:47–56.
- Chen, T., & Yang, M. (2020). Platelet-to-lymphocyte ratio is associated with cardiovascular disease in continuous ambulatory peritoneal dialysis patients. *International Immunopharmacology*, 78(September 2019). <https://doi.org/10.1016/j.intimp.2019.106063>
- Cho, J. H., Hur, I. K., Kim, C. D., Park, S. H., Ryu, H. M., Yook, J. M., Choi, J. Y., Choi, H. J., Choi, H. J., Park, J. W., Do, J. Y., & Kim, Y. L. (2010). Impact of systemic and local peritoneal inflammation on peritoneal solute transport rate in new peritoneal dialysis patients: A 1-year prospective study. *Nephrology Dialysis Transplantation*, 25(6), 1964–1973. <https://doi.org/10.1093/ndt/gfp767>.
- Cho, Y., Hawley, C. M., & Johnson, D. W. (2014). Clinical Causes of Inflammation in Peritoneal Dialysis Patients. 2014. <https://doi.org/10.1155/2014/909373>.
- Chugh, S., Chaudhry, S., Ryan, T., & Margetts, P. J. (2014). Peritoneal Membrane Injury and Peritoneal Dialysis. *Advances in Nephrology*, 2014, 1–10. <https://doi.org/10.1155/2014/573685>.
- Cobo, G., Lindholm, B., & Stenvinkel, P. (2018). Chronic inflammation in end-stage renal disease and dialysis. *Nephrology Dialysis Transplantation*, 33, iii35–iii40. <https://doi.org/10.1093/ndt/gfy175>.
- Devuyst, O., & Rippe, B. (2014). Water transport across the peritoneal membrane. *Kidney International*, 85(4), 750–758. <https://doi.org/10.1038/ki.2013.250>.
- Ellulu M. S., Patimah I., Khaza'ai H., Rahmat A., Abed Y., 2017, Obesity and inflammation: the linking mechanism and the complications, *Arch Med Sci*; 13, 4: 851–863.
- Eloueyk, A. K., Alameddine, R. Y., Osta, B. A., & Awad, D. M. (2019). Correlations between serum inflammatory markers and comorbidities in patients with end-stage renal disease. *Journal of Taibah University Medical Sciences*, 14(6), 547–552. <https://doi.org/10.1016/j.jtumed.2019.10.003>.
- Flessner, M. F., Credit, K., Henderson, K., Vanpelt, H. M., Potter, R., He, Z., Henegar, J., & Robert, B. (2007). Peritoneal changes after exposure to

- sterile solutions by catheter. *Journal of the American Society of Nephrology*, 18(8), 2294–2302. <https://doi.org/10.1681/ASN.2006121417>.
- Forget, P., Khalifa, C., Defour, J. P., Latinne, D., Van Pel, M. C., & De Kock, M. (2017). What is the normal value of the neutrophil-to-lymphocyte ratio? *BMC Research Notes*, 10(1), 1–4. <https://doi.org/10.1186/s13104-016-2335-5>.
- Gracia-Iguacel, C., González-Parra, E., Barril-Cuadrado, G., Sánchez, R., Egido, J., Ortiz-Ardúan, A., & Carrero, J. J. (2014). Definiendo el síndrome de desgaste proteico energético en la enfermedad renal crónica: Prevalencia e implicaciones clínicas. *Nefrologia*, 34(4), 507–519.
- Grossniklaus D. A., O'Brien G. M. C., Clark P. C., Dunbar S. B., 2008, Nutrient Intake in Heart Failure Patients, *J Cardiovasc Nurs*. 23(4): 357–363.
- Gürel, A., Doğan, A., Çeliker, H., Ulu, R., Yilit, I. P., & Aygen, B. (2015). Neutrophil to lymphocyte and platelet to lymphocyte ratios in peritoneal dialysis-related peritonitis. *Turkish Nephrology, Dialysis and Transplantation Journal*, 24(2), 158–161. <https://doi.org/10.5262/tndt.2015.1002.03>.
- Han, S. H., & Han, D. S. (2012). Nutrition in patients on peritoneal dialysis. *Nature Reviews Nephrology*, 8(3), 163–175, <https://doi.org/10.1038/nrneph>.
- Haryanti E., Kandarini Y., Widiana I. G. R., Sudhana W., Loekman J., Suwitra K., 2010., Kejadian Peritonitis pada Pasien Continuous Ambulatory Peritoneal Dialysis: Identifikasi Mikroorganisme dan Sensitivitas Antibiotik, *Divisi Ginjal dan Hipertensi, Bagian/SMF Ilmu Penyakit Dalam, FK Unud/RSUP Sanglah, Denpasar*.
- Hidayati HK, Suhardi, Titiek. Hubungan antara Hipertensi, Merokok dan Minuman Suplemen Energi dan Kejadian Penyakit Ginjal Kronik. 2012; 24.
- Ho, L. C., Wang, H. H., Chiang, C. K., Hung, K. Y., & Wu, K. D. (2010). Malnutrition-inflammation score independently determined cardiovascular and infection risk in peritoneal dialysis patients. *Blood Purification*, 29(3), 308–316. <https://doi.org/10.1159/000280641>.
- Ikizler, T. A., Wingard, R. L., & Hakim, R. M. (1995). Malnutrition in peritoneal dialysis patients: Etiologic factors and treatment options. *Peritoneal Dialysis International*, 15(SUPPL. 5). <https://doi.org/10.1177/089686089501505S09>.
- Johansson, L. (2015). Nutrition in older adults on peritoneal dialysis. *Peritoneal Dialysis International*, 35(6), 655–658. <https://doi.org/10.3747/pdi.2014.00343>
- Kalantar-Zadeh, K., Kopple, J. D., Humphreys, M. H., & Block, G. (2004). Comparing outcome predictability of markers of malnutrition-inflammation complex syndrome in haemodialysis patients. *Nephrology Dialysis Transplantation*, 19(6), 1507–1519. <https://doi.org/10.1093/ndt/gfh143>.
- Kang EW, Goo Y.S., Han S.H., Yoon S. Y., Choi S. R., Factor affecting malnutrition in continuous ambulatory peritoneal dialysis patients: a cross-sectional study, *Korea J. Nephrol*. 2002; 21:943-55.
- Kato, S., Abe, T., Lindholm, B., & Maruyama, S. (2015). Neutrophil/lymphocyte ratio: A promising prognostic marker in patients with chronic kidney

- disease. Inflammation and Cell Signaling, 1–5. <https://doi.org/10.14800/ics.683>.
- Kaya, T., Açıkgoz, S. B., Yıldırım, M., Nalbant, A., Altaş, A. E., & Cinemre, H. (2019). Association between neutrophil-to-lymphocyte ratio and nutritional status in geriatric patients. *Journal of Clinical Laboratory Analysis*, 33(1), 1–6. <https://doi.org/10.1002/jcla.22636>
- Kazancıoğlu R., 2013, Risk factors for chronic kidney disease: an update, Division of Nephrology, Faculty of Medicine, Bezmialem Vakif University, Istanbul, Turkey *Kidney International Supplements*, 3, 368–371.
- Kennedy C. and Bargman J., 2020, Peritoneal Dialysis in the Obese Patient *CJASN* 15: ccc–ccc,. doi: <https://doi.org/10.2215/CJN.10300819>.
- Kiebalo, T., Holotka, J., Habura, I., & Pawlaczyk, K. (2020). Nutritional status in peritoneal dialysis: Nutritional guidelines, adequacy and the management of malnutrition. *Nutrients*, 12(6), 1–14. <https://doi.org/10.3390/nu12061715>.
- Kim S., Eliot M., Koestler D.K., Wen-Chih Wu, Kelsey K.T., 2018, Association of Neutrophil-to-Lymphocyte Ratio With Mortality and Cardiovascular Disease in the Jackson Heart Study and Modification by the Duffy Antigen Variant *JAMA Cardiol.*;3(6):455-462. doi:10.1001/jamacardio.
- Lai, K. N., & Leung, J. C. K. (2010). Inflammation in peritoneal dialysis. *Nephron - Clinical Practice*, 116(1). <https://doi.org/10.1159/000314544>
- Lambie, M., Chess, J., Donovan, K. L., Kim, Y. L., Do, J. Y., Lee, H. B., Noh, H., Williams, P. F., Williams, A. J., Davison, S., Dorval, M., Summers, A., Williams, J. D., Bankart, J., Davies, S. J., & Topley, N. (2013). Independent effects of systemic and peritoneal inflammation on peritoneal dialysis survival. *Journal of the American Society of Nephrology*, 24(12), 2071–2080. <https://doi.org/10.1681/ASN.2013030314>
- Leung J. C. K., Chan L. Y. Y., Lai K. N., Tang S. C. W., 2013, Inflammation in Peritoneal Dialysis, Chapter 4: 55-81.
- Liabsuetrakul T., 2011, Is International or Asian Criteria-based Body Mass Index Associated with Maternal Anaemia, Low Birthweight, and Preterm Births among Thai Population? An Observational Study, *J HEALTH POPUL NUTR Jun*;29(3):218-228.
- Liu, C. C., Ko, H. J., Liu, W. S., Hung, C. L., Hu, K. C., Yu, L. Y., & Shih, S. C. (2019). Neutrophil-to-lymphocyte ratio as a predictive marker of metabolic syndrome. *Medicine (United States)*, 98(43). <https://doi.org/10.1097/MD>.
- Locatelli, F., Fouque, D., Heimbürger, O., Drüeke, T. B., Cannata-Andía, J. B., Hörl, W. H., & Ritz, E. (2002). Nutritional status in dialysis patients: A European consensus. *Nephrology Dialysis Transplantation*, 17(4), 563–572. <https://doi.org/10.1093/ndt/17.4.563>
- Lu, X., Wang, S., Zhang, G., Xiong, R., & Li, H. (2018). High neutrophil-to-lymphocyte ratio is a significant predictor of cardiovascular and all-cause mortality in patients undergoing peritoneal dialysis. *Kidney and Blood Pressure Research*, 43(2), 490–499. <https://doi.org/10.1159/000488696>.
- Malhotra R., Marcelli D., Gersdorff G. V., Grassmann A., Schaller M., Bayh I., Scatizzi L., Etter M., Guinsburg A., Barth G., Marelli C., Carioni P., Sande F. V. D., Kooman J. P., Liu E, Toffelmire T., Wang Y., Thijssen S., Usvyat

- L. A., Kotanko P., 2015, Relationship of Neutrophil-to-Lymphocyte Ratio and Serum Albumin Levels with C-Reactive Protein in Hemodialysis Patients: Results from 2 International Cohort Studies, 130(4):263-70. doi: 10.1159/000437005. Epub
- Manani, S. M., Virzì, G. M., Clementi, A., Brocca, A., De Cal, M., Tantillo, I., Ferrando, L., Crepaldi, C., & Ronco, C. (2016). Pro-inflammatory cytokines: A possible relationship with dialytic adequacy and serum albumin in peritoneal dialysis patients. *Clinical Kidney Journal*, 9(1), 153–157. <https://doi.org/10.1093/ckj/sfv137>.
- Marques, R., Carias, E., Domingos, A., Guedes, A., Bernardo, I., & Neves, P. (2021). Prognostic value of lymphocyte cell ratios in peritoneal dialysis. *Portuguese Journal of Nephrology & Hypertension*, 35(1), 18–21. <https://doi.org/10.32932/pjnh.2021.04.112>.
- Marshall. M. R., 2021, A systematic review of peritoneal dialysis-related peritonitis rates over time from national or regional population-based registries and databases, *Peritoneal Dialysis International XX(X)*.
- Martinez J. D., 2019, The Relationship of a Novel Marker of Inflammation (Neutrophil to-Lymphocyte Ratio) to Nutritional Status, Diet and Clinical Outcomes in Hemodialysis Patients, Florida International University.
- Martins, E. C., Da Fe Silveira, L., Viegas, K., Beck, A. D., Júnior, G. F., Cremonese, R. V., & Lora, P. S. (2019). Neutrophil-lymphocyte ratio in the early diagnosis of sepsis in an intensive care unit: A case-control study. *Revista Brasileira de Terapia Intensiva*, 31(1), 63–70. <https://doi.org/10.5935/0103-507X.20190010>
- Mihai, S., Codrici, E., Popescu, I. D., Enciu, A. M., Albulescu, L., Necula, L. G., Mambet, C., Anton, G., & Tanase, C. (2018). Inflammation-related mechanisms in chronic kidney disease prediction, progression, and outcome. *Journal of Immunology Research*, 2018. <https://doi.org/10.1155/2018/2180373>
- Morelle, J., & Devuyst, O. (2015). Water and solute transport across the peritoneal membrane. *Current Opinion in Nephrology and Hypertension*, 24(5), 434–443. <https://doi.org/10.1097/MNH.0000000000000151>.
- Noh, H., Kim, J. S., Han, K. H., Lee, G. T., Song, J. S., Chung, S. H., Jeon, J. S., Ha, H., & Lee, H. B. (2006). Oxidative stress during peritoneal dialysis: Implications in functional and structural changes in the membrane. *Kidney International*, 69(11), 2022–2028. <https://doi.org/10.1038/sj.ki.5001506>
- Nowak, K. L., & Chonchol, M. (2018). Does inflammation affect outcomes in dialysis patients? *Seminars in Dialysis*, 31(4), 388–397. <https://doi.org/10.1111/sdi.12686>
- PERNEFRI. (2011). Perhimpunan Nefrologi Indonesia Perhimpunan Nefrologi Indonesia (PERNEFRI) 2011.
- Pineault, J., Lamarche, C., Bell, R., Lafrance, J. P., Ouellet, G., Leblanc, M., Pichette, V., Bezzaoucha, S., & Vallée, M. (2017). Association of Neutrophil-to-Lymphocyte Ratio With Inflammation and Erythropoietin Resistance in Chronic Dialysis Patients. *Canadian Journal of Kidney Health and Disease*, 4, 1–7. <https://doi.org/10.1177/2054358117735563>

- Preedy, V. R., & Patel, V. B. (2019). Handbook of famine, starvation, and nutrient deprivation: From biology to policy. In Handbook of Famine, Starvation, and Nutrient Deprivation: From Biology to Policy. <https://doi.org/10.1007/978-3-319-55387-0>.
- Purwanto B., Yuniasih D., Puspitasari M., 2018, Albumin Description of Continuous Ambulatory Peritoneal Dialysis Patients in Sardjito Hospital Yogyakarta ADMJ (Ahmad Dahlan Medical Journal), VOL 2, No. 1, 10-18.
- Radić, J., Bašić-Jukić, N., Vujičić, B., Klarić, D., Radulović, G., Jakić, M., Jurić, K., Altabas, K., Grđan, Ž., Kovačević-Vojtušek, I., Martinović, V. Č., Janković, N., Gulin, M., Ljutić, D., & Rački, S. (2017). Anemia is correlated with malnutrition and inflammation in croatian peritoneal dialysis patients: A multicenter nationwide study. *Peritoneal Dialysis International*, 37(4), 472–482. <https://doi.org/10.3747/pdi.2016.00013>.
- Rajendra W. U., Ramya N, Shankar S. P., 2020, A Study of Neutrophil/Lymphocyte Ratio in Chronic Kidney Disease, Department of General Medicine, Aarupadai Veedu Medical College, Pondicherry, India, J. Evid. Based Med. Healthc., pISSN- 2349-2562, eISSN- 2349-2570/ Vol. 7/Issue 24.
- Rambod, M., Kovesdy, C. P., & Kalantar-Zadeh, K. (2008). Malnutrition-Inflammation Score for risk stratification of patients with CKD: Is it the promised gold standard? Commentary. *Nature Clinical Practice Nephrology*, 4(7), 354–355, <https://doi.org/10.1038/ncpneph0834>.
- Ryota Yoshitomia,B, Masaru Nakayamab, Teppei Sakohb, Akiko Fukuib, Eisuke Katafuchib, Makiko Sekib,
- Rodríguez E. R., Ana M., Sobaler L.,Ortega R. M., Losada M. L. D., Parra L., Aparicio A., 2020, Association between Neutrophil-to-Lymphocyte Ratio with Abdominal Obesity and Healthy Eating Index in a Representative Older Spanish Population *Nutrients*, 12, 855; doi:10.3390/nu12030855.
- Sahathevan, S., Khor, B. H., Ng, H. M., Gafor, A. H. A., Daud, Z. A. M., Mafra, D., & Karupaiah, T. (2020). Understanding development of malnutrition in hemodialysis patients: A narrative review. *Nutrients*, 12(10), 1–30. <https://doi.org/10.3390/nu12103147>.
- Salzer W. L., 2018, Peritoneal dialysis-related peritonitis: challenges and solutions, *International Journal of Nephrology and Renovascular Disease*: 11 173–186.
- Satirapoj, B., Limwannata, P., Kleechaiyaphum, C., Prapakorn, J., Yatinan, U., Chotsriluecha, S., & Supasynndh, O. (2017). Nutritional status among peritoneal dialysis patients after oral supplement with ONCE dialyze formula. *International Journal of Nephrology and Renovascular Disease*, 10, 145–151. <https://doi.org/10.2147/IJNRD.S138047>.
- Saunders J., 2010, Malnutrition: causes and consequences, *Clinical Medicine*, Vol 10, No 6: 624–7.
- Shi, J., Yu, M., & Sheng, M. (2017). Angiogenesis and Inflammation in Peritoneal Dialysis: The Role of Adipocytes. *Kidney and Blood Pressure Research*, 42(2), 209–219. <https://doi.org/10.1159/000476017>.

- Soraya S., Ramayani O. R., Siregar R., Siregar B., Kelainan Gigi dan Mulut pada Penderita Penyakit Ginjal Kronik, *The Journal of Medical School (JMS)* Vol. 52, No. 2, 2019 | 89 – 94.
- Spatola, L., Finazzi, S., Calvetta, A., Angelini, C., & Badalamenti, S. (2019). Subjective Global Assessment–Dialysis Malnutrition Score and arteriovenous fistula outcome: A comparison with Charlson Comorbidity Index. *Journal of Vascular Access*, 20(1), 70–78. <https://doi.org/10.1177/1129729818779550>
- Spoto, B., Pisano, A., & Zoccali, C. (2016). Insulin resistance in chronic kidney disease: A systematic review. *American Journal of Physiology - Renal Physiology*, 311(6), F1087–F1108. <https://doi.org/10.1152/ajprenal.00340>.
- Stenvinkel, P., Lindholm, B., Lönnqvist, F., Katzarski, K., Heimbürger, O. 2000, Increases in serum leptin levels during peritoneal dialysis are associated with inflammation and a decrease in lean body mass. *Journal of the American Society of Nephrology*, 11(7), 1303–1309. <https://doi.org/10.1681/asn.v11i71303>.
- Stenvinkel P., Chung S. H., Heimbürger O., Lindholm B., 2001, Malnutrition, Inflammation, and Atherosclerosis in Peritoneal Dialysis Patients, Department of Nephrology and Baxter Novum, Huddinge University Hospital, Stockholm, Sweden, Proceedings of the ISPD—The IX th Congress of the ISPD Montréal, Canada, Peritoneal Dialysis International, Vol. 21, Supplement 3.
- Swartz, R., Messina, J., Reynolds, J., & Ranjit, U. (1991). Simultaneous catheter replacement and removal in refractory peritoneal dialysis infections. *Kidney International*, 40(6), 1160–1165. <https://doi.org/10.1038/ki.1991.329>
- Szeto C., C., and Li P., K., T., 2019, Peritoneal Dialysis–Associated Peritonitis, *CJASN* 14: 1100–1105, July, PD-Associated Peritonitis.
- Thaha dan Widiarta, 2019., The Role of Inflammation in Chronic Kidney Disease, *Indonesian Journal of Kidney and Hypertension*, InaKidney | Vol. II | Is. 3| Sept - Dec 2019.
- Tjekyan R. Prevalensi dan Faktor Risiko Penyakit Ginjal Kronik di RSUP Dr. Mohammad Hoesin Palembang Tahun 2012. *Majalah Kedokteran Sriwijaya*. 2012: 46.
- Tsudab S., Nakanoa T., Tsuruyac K., Kitazonoa T., 2019, High neutrophil/lymphocyte ratio is associated with poor renal outcomes in Japanese patients with chronic kidney disease, vol. 41, NO. 1, 238–243.
- Turkmen, K., Guney, I., Yerlikaya, F. H., Zeki, H., Turkmen, K., Guney, I., Yerlikaya, F. H., & Tonbul, H. Z. (2012). The Relationship Between Neutrophil-to- Lymphocyte Ratio and Inflammation in End-Stage Renal Disease Patients The Relationship Between Neutrophil-to-Lymphocyte Ratio and Inflammation in End-Stage Renal Disease Patients. 6049. <https://doi.org/10.3109/0886022X.2011.641514>
- Unal, A., Kavuncuoglu, F., Duran, M., Oguz, F., Kocyigit, I., Sipahioglu, M. H., Tokgoz, B., & Oymak, O. (2015). Inflammation is associated to volume

- status in peritoneal dialysis patients. *Renal Failure*, 37(6), 935–940. <https://doi.org/10.3109/0886022X.2015.1040337>
- Vanholder, R., Glorieux, G., & Lameire, N. (2002). The other side of the coin: Impact of toxin generation and nutrition on the uremic syndrome. *Seminars in Dialysis*, 15(5), 311–314. <https://doi.org/10.1046/j.1525-139X.2002.00076.x>
- Wanner, C., Zimmermann, J., Schwedler, S., & Metzger, T. (2002). Inflammation and cardiovascular risk in dialysis patients. *Kidney International, Supplement*, 61(80). <https://doi.org/10.1046/j.1523-1755.61.s80.18.x>
- Wu, C. L., Wu, H. M., Chiu, P. F., Liou, H. H., Chang, C. Bin, Tarng, D. C., & Chang, C. C. (2014). Associations between the duration of dialysis, endotoxemia, monocyte chemoattractant protein-1, and the effects of a short-dwell exchange in patients requiring continuous ambulatory peritoneal dialysis. *PLoS ONE*, 9(10). <https://doi.org/10.1371/journal.pone.0109558>
- Wu H., Ye H., Huang R., Yi C., Wu J., 2020, Xueqing Yu and Xiao Yang Incidence and risk factors of peritoneal dialysis-related peritonitis in elderly patients: A retrospective clinical study, *Peritoneal Dialysis International*, Vol. 40(1) 26–33.
- Xu, H., & Carrero, J. J. (2017). Insulin resistance in chronic kidney disease. *Nephrology*, 22, 31–34. <https://doi.org/10.1111/nep.13147>
- Yuan, Q., Wang, J., Peng, Z., Zhou, Q., Xiao, X., Xie, Y., Wang, W., Huang, L., Tang, W., Sun, D., Zhang, L., Wang, F., Zhao, M. H., Tao, L., He, K., & Xu, H. (2019). Neutrophil-to-lymphocyte ratio and incident end-stage renal disease in Chinese patients with chronic kidney disease: Results from the Chinese Cohort Study of Chronic Kidney Disease (C-STRIDE). *Journal of Translational Medicine*, 17(1), 1–8. <https://doi.org/10.1186/s12967-019-1808-4>
- Yulianti, M., Suhardjono, S., Kresnawan, T., & Harimurti, K. (2017). Faktor-faktor yang Berkorelasi dengan Status Nutrisi pada Pasien Continuous Ambulatory Peritoneal Dialysis (CAPD). *Jurnal Penyakit Dalam Indonesia*, 2(1), 2. <https://doi.org/10.7454/jpdi.v2i1.59>
- Zhang J., Lu X., Wang S., Li H., 2021, *BioMed Research International High Neutrophil-to-Lymphocyte Ratio and Platelet-to-Lymphocyte Ratio Are Associated with Poor Survival in Patients with Hemodialysis*, Volume 2021, Article ID 9958081, 6 pages.