



## REFERENCES

- Abderrahim, A., Belhamel, K., Chalard, P., and Figuéreido, G. (2019) “Chemotypes and radical scavenging activity of the essential oils from *Artemisia arborescens* L. growing in three areas of Bejaia (Algeria),” *Journal of Food Measurement and Characterization*. 13(3), 2491–2499.
- Ashokkumar, M. (2015) “Applications of ultrasound in food and bioprocessing,” *Ultrasonics Sonochemistry*, 25(1), 17–23.
- Aydar, A. Y., Bagdatlioglu, N. and Köseoglu, O. (2017) “Effect of ultrasound on olive oil extraction and optimization of ultrasound-assisted extraction of extra virgin olive oil by response surface methodology (RSM),” *Grasas y Aceites*, 68(2), 189–201.
- Ayim, I., Ma, H., Alenyorege, E.A., Ali, Z., and Donkor, P.O. (2018) “Influence of ultrasound pretreatment on enzymolysis kinetics and thermodynamics of sodium hydroxide extracted proteins from tea residue,” *Journal of Food Science and Technology*, 55(3), 1037–1046.
- Boukhatem, M. N., Boumaiza, A., Nada, G.H., Rajabi, M., and Mousa, S.A. (2020) “*Eucalyptus globulus* essential oil as a natural food preservative: Antioxidant, antibacterial and antifungal properties in vitro and in a real food matrix (orangina fruit juice),” *Applied Sciences (Switzerland)*, 10(16), 1–17.
- Cárcel, J., García-Pérez, J., Benedito, J., and Mulet, A. (2012) “Food process innovation through new technologies: Use of ultrasound,” *Journal of Food Engineering*, 110(2), 200–207.
- Carrera, C., Ruiz-Rodríguez, A., Palma, M., and Barroso, C.G. (2012) “Ultrasound assisted extraction of phenolic compounds from grapes,” *Analytica Chimica Acta*, 732, 100–104.
- Castro-Puyana, M., Marina, M. L. and Plaza, M. (2017) “Water as green extraction solvent: Principles and reasons for its use,” *Current Opinion in Green and Sustainable Chemistry*. 5, 31–36.
- Chemat, F., Rombaut, N., Sicaire, N., Meullemiestre, A., Fabiano-Tixier, A.-S., and Abert-Vian, M. (2017) “Ultrasound assisted extraction of food and natural products. Mechanisms, techniques, combinations, protocols and applications. A review,” *Ultrasonics Sonochemistry*, 34, 540–560.
- Chen, S., Zeng, Z., Hu, N., Bai, B., Wang, H, and Suo, Y. (2018) “Simultaneous optimization of the ultrasound-assisted extraction for phenolic



compounds content and antioxidant activity of *Lycium ruthenicum* Murr. fruit using response surface methodology," *Food Chemistry*, 242, 1–8.

Deus, V. L., Resende, L.M., Bispo, E.S., Franca, A.S., Beatriz, M., and Gloria, A. (2021) "FTIR and PLS-regression in the evaluation of bioactive amines, total phenolic compounds and antioxidant potential of dark chocolates," *Food Chemistry*. 357, 129754–129763.

Enny, H.L., Wijaya, H., and Lestari, N. (2012) "Mempelajari Ekstraksi dan Stabilitas Total Karotenoid,  $\alpha$  dan  $\beta$  *Cryptoxanthin* dalam Ekstrak Buah Merah," *Jurnal Riset Teknologi Industri*, 6(12), 11–19.

Esmaeilzadeh Kenari, R., and Dehghan, B. (2020) "Optimization of ultrasound-assisted solvent extraction of hemp (*Cannabis sativa* L.) seed oil using RSM: Evaluation of oxidative stability and physicochemical properties of oil," *Food Science and Nutrition*, 8(9), 4976–4986.

Fasakin, A. O. (2017) "Effect of sonication at different ultrasonic frequencies on the quality and quantity of fatty acids of the oil of *Chlorella vulgaris*," *Scientific Research and Essays*, 12(16), 155–162.

Ferreira, S. L. C., Bruns, R. E., Ferreira, H. S., Matos, G. D., David, J. M., Brandão, G. C., da Silva, E. G.P., Portugal, L. A., dos Reis, P. S., Souza, A. S., and dos Santos, W.N.L. (2007) "Box-Behnken design: An alternative for the optimization of analytical methods," *Analytica Chimica Acta*, 597(2), 179–186.

Ghadermazi, R., Keramat, J. and Goli, S. A. H. (2016) "Antioxidant activity and physical properties of hydroxypropylmethylcellulose films enriched with essential oils," *Journal of Food and Nutrition Research*, 55(1), 22–32.

Goula, A.M., Ververi, M., Adamopoulou, A., and Kaderides, K. (2017) "Green ultrasound-assisted extraction of carotenoids from pomegranate wastes using vegetable oils," *Ultrasonics Sonochemistry*, 34, 821–830.

Ilyasov, I. R., Beloborodov, V.L., Selivanova, I.A., Terekhov, R.P. (2020) "ABTS/PP Decolorization Assay of Antioxidant Capacity Reaction Pathways," *International Journal of Molecular Sciences*, 21(3), 1131–1158.

Jalili, F., Jafari, S.M., Emam-Djomeh, Z., Malekjani, N., and Farzaneh, Vahid. (2018) "Optimization of Ultrasound-Assisted Extraction of Oil from Canola Seeds with the Use of Response Surface Methodology," *Food Analytical Methods*, 11(2), 598–612.

Johnson, L. (2002) "Recovery, Refining, Converting, and Stabilizing Edible Fats



and Oils," in Akoh C.C., M. D. B. (ed.) *Food Lipids: Chemistry, Nutrition, and Biotechnology*. Marcel Dekker; New York, NY, USA, 205–244.

Kappahycus, A. and Kata, C. (2011) "Pengujian kandungan total fenol *Kappahycus alvarezii* dengan metode ekstraksi ultrasonik dengan variasi suhu dan waktu," *Prosiding SNST*, 3, 40–44.

Kumar, V., Mathela, C.S., Kumar, M., and Tewari, G. (2019) "Antioxidant potential of essential oils from some *Himalayan Asteraceae* and *Lamiaceae* species," *Medicine in Drug Discovery*. 1, 100004–100011.

Majid, H. and Silva, F. V. M. (2020) "Optimisation of ultrasound assisted extraction of antiacetylcholinesterase and antioxidant compounds from manuka (*Leptospermum scoparium*) for use as a phytomedicine against Alzheimer's disease," *New Zealand Journal of Forestry Science*, 50, 1–12.

Mayara, T. P., Del Castillo B.D., Serrão P.C.D., Lobato R.A.B., da Silva R., de Oliveira P.F., Ferreira S.P.S., Távora, T.P.L., and da Silva de A.S.S.M. (2016) "Antioxidant effect of plant extracts of the leaves of *Tithonia diversifolia* (Hemsl.) A. Gray on the free radical DPPH," *Journal of Chemical and Pharmaceutical Research*, 8(8), 1182–1189.

Medina-Torres, N., Ayora-Talavera, T., Espinosa-Andrews, H., Sánchez-Contreras, A., and Pacheco, N. (2017) "Ultrasound Assisted Extraction for the Recovery of Phenolic Compounds from Vegetable Sources," *Agronomy*, 7, 47–66.

Mentari, F. and Vifta, R. L. (2019) "Evaluasi Sifat Fisika-Kimia dan Karakteristik Antioksidan Ekstrak Etanol Biji Timun Suri (*Cucumis melo* L.var Terenkapsulasi Kitosan)," *Jurnal Riset Teknologi Industri*, 13(2), 125–135.

Mohammadpour, H., Sadrameli, S.M., Eslami, F., and Asoodeh, A. (2019) "Optimization of ultrasound-assisted extraction of *Moringa peregrina* oil with response surface methodology and comparison with Soxhlet method," *Industrial Crops and Products*, 131, 106–116.

Murtiningrum, Sarunggallo, Z.L., Roreng, M.K., Santoso, B., and Armiaty. (2019) "Chemical properties, carotenoid, tocopherol and fatty acid composition of three clones of red fruit (*Pandanus conoideus* Lam.) oil of different ripening stages," *International Food Research Journal*, 26(2), 649–655.

Murtiningrum, S. K. S. dan K. (2012) "Ekstraksi Minyak Dengan Metode Wet Rendering Dari Buah Pandan (*Pandanus Conoideus* L)," *Journal of*



*Agroindustrial Technology*, 15(1), 28–33.

Mwaurah, P. W., Kumar, S., Kumar, N., Attkan, A.K., Panghal, A., Singh, V.K., and Garg, M.K. (2020) “Novel oil extraction technologies: Process conditions, quality parameters, and optimization,” *Comprehensive Reviews in Food Science and Food Safety*, 19(1), 3–20.

Nde, D. B. and Foncha, A. C. (2020) “Optimization methods for the extraction of vegetable oils: A review,” *Processes*, 8(2), 1–21.

Nurmiah, S., Syarief, R., Sukarno., Peranginangin, R., and Nurtama, B. (2013) “Application of Response Surface Methodology in The Optimization of Process Conditions of Alkali Treated Cottonii (ATC) Processing,” *Jurnal Pascapanen dan Bioteknologi Kelautan dan Perikanan*, 8(1), 9–22.

Panadare, D. C., Gondaliya, A. and Rathod, V. K. (2020) “Comparative study of ultrasonic pretreatment and ultrasound assisted three phase partitioning for extraction of custard apple seed oil,” *Ultrasonics Sonochemistry*, 61, 104821–104829.

Perincek, O. and Colak, M. (2013) “Use of Experimental Box-Behnken Design for the Estimation of Interactions Between Harmonic Currents Produced by Single Phase Loads,” *International Journal of Engineering Research and Applications*, 3(2), 158–165.

Puertolas, E., Alvarez-Sabaté, S., and Cruz, Z. (2016) “Pulsed electric field: groundbreaking technology for improving olive oil extraction,” *Inform*, 3(27), 12–14.

Puspita, D., Kristen, U. and Wacana, S. (2020) “Aktivitas antioksidan ekstrak buah merah (*Pandanus conoideus*) pada minyak kelapa yang diinkubasi selama 15 Hari,” *Prosiding Seminar Nasional Kesehatan Universitas Kristen Satya Wacana*. July, 19–24.

Putra, S. (2018) “Prospek Buah Merah Menjanjikan,” *Direktorat Jenderal Holtikultura Kementerian Pertanian*, 1. <http://hortikultura.pertanian.go.id/?p=2860>. [15 Maret 2022]

Rohman, A., Riyanto, S., Yuniarti, N., Saputra, W.R., Utami, R., and Mulatsih, W. (2010) “Antioxidant activity, total phenolic, and total flavaonoid of extracts and fractions of red fruit (*Pandanus conoideus* Lam),” *International Food Research Journal*, 97–106.

Rohman, A. Riyanto, S., Mart Sasi, A., and Yusof, F.M. (2014) “The use of FTIR spectroscopy in combination with chemometrics for the authentication of red fruit (*Pandanus conoideus* Lam) oil from sunflower and palm oils,”



*Food Bioscience*, 7, 64–70.

Rohman, A., Sugeng, R. and Che Man, Y. B. (2012) “Characterizaton of red fruit (*Pandanus conoideus* Lam) oil,” *International Food Research Journal*, 19(2), 563–567.

Rosas-Mendoza, M. E., Coria-Hernández, J., Meléndez-Pérez, R., Arjona-Román, J.L. (2017) “Characteristics of chia (*Salvia hispanica* L.) seed oil extracted by ultrasound assistance,” *Journal of the Mexican Chemical Society*, 61(4), 326–335.

Sánchez-Vioque, R., Polissiou, M., Astraka, K., Mozos-Pascual, M. de los., Tarantilis, P., Herraiz-Peñaiver, D., and Santana-Méridas, O. (2013) “Polyphenol composition and antioxidant and metal chelating activities of the solid residues from the essential oil industry,” *Industrial Crops and Products*, 49, 150–159.

Sandhiutami, N., Ngatidjan, N. and Kristin, E. (2012) “Penetapan kadar tokoferol minyak buah merah (*Pandanus conoideus* Lam.) secara in vitro dan in vivo,” *Jurnal Tumbuhan Obat Indonesia*, 5(1), 45–52.

Sangkala, S., Jura, M. R. and Tangkas, I. M. (2014) “Uji Aktivitas Antioksidan Ekstrak Buah Merah (*Pandanus baccari* L.) di Daerah Poso Sulawesi Tengah,” *Jurnal Akademika Kimia*, 3(4), 198–205.

Sarungallo, Z. L. Murtiningrum, P. S. (2011) “Sifat Fisikokimia Minyak Kasar dan Hasil Degumming dari Buah Merah (*Pandanus conoideus* L.) yang diekstrak secara Tradisional Merdey,” *Jurnal Agrotek*, 6(1), 9–15.

Sarungallo, Z. L., Hariyadi, P., Andarwulan, N., Purnomo, E.H., and Wada, M. (2015<sup>a</sup>) “Analysis of  $\alpha$ -Cryptoxanthin,  $\beta$ -Cryptoxanthin,  $\alpha$  -Carotene, and  $\beta$ -Carotene of *Pandanus Conoideus* Oil by High-performance Liquid Chromatography (HPLC),” *Procedia Food Science*, 3, 231–243.

Sarungallo, Z.L., Hariyasi, P., Andarwulan, N., and Purnomo, E.H. (2015<sup>b</sup>) “Characterization of chemical properties, lipid profile, total phenol and tocopherol content of oils extracted from nine clones of red fruit (*Pandanus conoideus*),” *Kasetsart Journal - Natural Science*, 49(2), 237–250.

Sarungallo, Z. L., Murtiningrum, Santoso, B., Roreng, M.K., and Latumahina, R.M.M. (2016) “Nutrient content of three clones of red fruit (*Pandanus conoideus*) during the maturity development,” *International Food Research Journal*, 23, 1217–1225.

Sarungallo, Z. L., Hariyadi, P., Andarwulan, N., and Purnomo, E.H. (2019)



“Keragaman Karakteristik Fisik Buah, tanaman dan Rendemen Minyak dari 9 Klon Buah Merah (*Pandanus conoideus*),” *Agrikan: Jurnal Agribisnis Perikanan*, 12(1), 70–82.

Sarungallo, Z.L.; Hariyadi, P.; Andarwulan, A.; and Purnomo, E.H. (2020) “Effect of heat treatment prior to extraction on the yield and quality of red fruit (*Pandanus conoideus*) oil,” *Food Research*, 4(3), 659–665.

Sarungallo, Z. L. , Hariyadi, P., Andarwulan, N., and Purnomo, E. H. (2014) “The effect of extraction method on the chemical quality and fatty acid composition of red fruit (*Pandanus Conoideus*) oil,” *Jurnal Teknologi Industri Pertanian*, 24(3), 209–217.

Satriyanto, B., Widjanarko, S. B. and Yunianta (2012) “Heat Stability of Red Fruit Extract Color as Potential Source of Natural Pigments,” *Teknologi Pertanian*, 13(3), 157–168.

Sekarsari, S., Widarta, I. W. R. and Jambe, A. A. G. N. A. (2019) “The influence of time and temperature with ultrasonic waves on antioxidant activity of extracts guajava leaves (*Psidium guajava* L.),” *Jurnal Ilmu dan Teknologi Pangan (ITEPA)*, 8(3), 267–277.

Senrayan, J. and Venkatachalam, S. (2020) “Ultrasonic acoustic-cavitation as a novel and emerging energy efficient technique for oil extraction from kapok seeds,” *Innovative Food Science and Emerging Technologies*, 62, 102347–102358.

Setyaningrum, D. L., Riyanto, S. and Rohman, A. (2013) “Analysis of corn and soybean oils in red fruit oil using FTIR spectroscopy in combination with partial least square,” *International Food Research Journal*, 20(4), 1977–1981.

Sharma, M., Dadhwal, K., Gat, Y., Kumar, V., Panghal, A., Prasad, R., Kaur, S., and Gat, P. (2019) “A review on newer techniques in extraction of oleaginous flaxseed constituents,” *OC&L - Oilseeds and fats, Crops and Lipids*, 26, 1–7.

Sinaga, F. A. and Purba, P. (2018) “The influence of red fruit oil on creatin kinase level at maximum physical activity,” *Asian Journal of Pharmaceutical and Clinical Research*, 11(13), 104–109.

Soehendro, A. W., Manuhara, g., and Nurhartadi, E. (2015) “Effects of Temperatures on Antioxidant and Antimicrobia Activity of Melinjo Seed (*Gnetum gnemon* L.) with Ethanol and Water as Solvent,” *Teknosains Pangan*, IV(4), 15–24.



Stevanato, N. and da Silva, C. (2019) "Radish seed oil: Ultrasound-assisted extraction using ethanol as solvent and assessment of its potential for ester production," *Industrial Crops and Products*, 132(December 2018), 283–291.

Taha, A., Ahmed, E., Ismaiel, A., Ashokkumar, M., Xu, X., Pan, S., and Hu, H. (2020) "Ultrasonic emulsification: An overview on the preparation of different emulsifiers-stabilized emulsions," *Trends in Food Science and Technology*. 105, 363–377.

Tiwari, B. K. (2015) "Ultrasound: A clean, green extraction technology," *TrAC - Trends in Analytical Chemistry*, 71, 100–109.

Tungmannithum, D., Elamrani, A., Abid, M., Drouet, S., Kiani, R., Garros, L., Kabra, A., Addi, M., Hano, C. (2020) "A quick, green and simple ultrasound-assisted extraction for the valorization of antioxidant phenolic acids from moroccan almond cold-pressed oil residues," *Applied Sciences (Switzerland)*, 10(9), 3313–3333.

Valantina, Rubalya S.; Neelamegam, P. (2015) "Selective ABTS and DPPH-radical scavenging activity of peroxide from vegetable oils," *International Food Research Journal*, 22(1), 289–294.

Vatai, T., Škerget, M. and Knez, Ž. (2009) "Extraction of phenolic compounds from elder berry and different grape marc varieties using organic solvents and/or supercritical carbon dioxide," *Journal of Food Engineering*, 90(2), 246–254.

Vijayalakshmi, M. and Ruckmani, K. (2016) "Ferric reducing anti-oxidant power assay in plant extract," *Bangladesh Journal of Pharmacology*. Bangladesh: J Pharmacol, 11(3), 570–572.

Wada, M., Fujimoto, K., Nishigaki, T., Febriyanti, E., Ikeda, R., and Nakashima, K. (2013) "Determination of carotenoid in Buah Merah using HPLC.pdf," *Warta ITP Balai Besar Industri Agro*, 30(1), 1–8.

Wang, J., Lin, H., Gao, H., Zhao, J., Zhou, L., Han, J., Yu, Z., and Yung, F. (2011) "Antimicrobial and antioxidant activities of the flower essential oil of Halimodendron halodendron," *Natural Product Communications*, 6(11), 1749–1753.

Wazir, D., Ahmad, S., Muse, R., Mahmood, M., and Shukor, M.Y. (2011) "Antioxidant activities of different parts of Gnetum gnemon L.," *Journal of Plant Biochemistry and Biotechnology*, 20(2), 234–240.

Wen, C., Zhang, J., Zhang, H., Dzah, C.S., Zandile, M., Duan, Y., Ma, H., and



Luo, X. (2018) "Advances in ultrasound assisted extraction of bioactive compounds from cash crops – A review," *Ultrasonics Sonochemistry*, 48(July), 538–549.

Wijaya, H. and Pohan, H. G. (2009) "Kajian Teknis Standar Minyak Buah Merah", Prosiding PPI Standardisasi.

Wu, S.-Q., Li, R., Jing, Zi-Tao., Wang, Y., Tan, J., and Tang, Shu-Hua. (2020) "Evaluation of antioxidant active ingredients of spikenard essential oil by ultra-fast gas chromatography electronic nose and radical scavenging mechanism," *Industrial Crops and Products*, 151, 112489–112496.

Xia, N., Schirra, S., Hasselwander, S., Förstermann, U., and Li, H. (2018) "Red fruit (*Pandanus conoideus* Lam) oil stimulates nitric oxide production and reduces oxidative stress in endothelial cells," *Journal of Functional Foods*, 51, 65–74.

Yadav, L., Bhatnagar, V. and Mudgal, V. D. (2014) "High power ultrasound: An innovation in the food processing industry," *Food Science Research Journal*, 5(2), 190–196.

Yuhono, J. T. and Rini, P.E. (2007) "Analysis of marketing system and equilibriate point price at the red fruit oil (*Pandanus conoideus*) industry in Papua Province," *Bul.Littro*, XVIII(2), 188–202.

Zhang, W., Pan, Yong-Gui., Huang, W., Chen, H., and Yang, H. (2019) "Optimized ultrasonic-assisted extraction of papaya seed oil from Hainan/Eksotika variety," *Food Science and Nutrition*, 7(8), 2692–2701.

Zhou, J., Ma, Y., Jia, Y., Pang, M., Cheng, G., and Cai, S. (2019) "Phenolic profiles, antioxidant activities and cytoprotective effects of different phenolic fractions from oil palm (*Elaeis guineensis* Jacq.) fruits treated by ultra-high pressure," *Food Chemistry*, 288, 68–77.