

- Almajano, P., Carbo, R., Angel J., Lo'pez Jime'nez ,Michael H. dan Gordon,.(2018).Antioxidant and antimicrobial activities of tea infusions Chemistry 108 : 55–63
- Astill, C. M. R. Birch, C. Dacombe, P. G. Humphrey, dan P. T. Martin, (2001). *Factaors affecting the caffeine and polyphenol contents of black and green tea infusions*, J. Agric. Food Chem., vol. 49, no. 11, pp. 5340–5347.
- Baba, R., dan Kumazawa, K. (2014). Characterization of the potent odorants contributing to the characteristic aroma of chinese green tea infusions by aroma extract dilution analysis. Journal of Agricultural and Food Chemistry, 62(33), 8308–8313.
- Baptista, J. A. B., Tavares, J. F. d. P., dan Carvalho, R. C. B. (1998). Comparison of catechins and aromas among different green teas using HPLC/SPME-GC. Food Research International, 31(10), 729–736.
- Chen, C. N., Liang, C. M., Lai, J. R., Tsai, Y. J., Tsay, J. S., dan Lin, J. K. (2003). Capillary Electrophoretic Determination of Theanine, Caffeine, and Catechins in Fresh Tea Leaves and Oolong Tea and Their Effects on Rat Neurosphere Adhesion and Migration. Journal of Agricultural and Food Chemistry, 51(25), 7495–7503.
- Chen, C. -W, dan Ho, C. -T. (1995). Antioxidant Properties of Polyphenols Extracted From Green and Black Teas. Journal of Food Lipids, 2(1), 35–46.
- Deng, W.W., S. Ogita, dan H. Ashihara,(2008). *Biosynthesis of theanine (γ-ethylamino-l-glutamic acid) in seedlings of Camellia sinensis*, Phytochem. Lett., 1, 2, 115–119.
- Du, L., Li, J., Li, W., Li, Y., Li, T., dan Xiao, D. (2014). Characterization of volatile compounds of pu-erh tea using solid-phase microextraction and simultaneous distillationextraction coupled with gas chromatography-mass spectrometry. Food Research International, 57, 61–70.
- E.Z. Su, T. Xia, dan L.P. Gao., Immobilization of beta-glucosidase and its aroma-increasing effect on tea beverage, Food Bioprod. Process 88 (2010) 83–89.
- Erba, D., Riso, P., Bordoni, A., Foti, P., Biagi, P. L., dan Testolin, G. (2005). Effectiveness of moderate green tea consumption on antioxidative status and plasma lipid profile in humans. Journal of Nutritional Biochemistry, 16(3), 144–149.
- Friedman, M., Levin, C. E., Lee, S. U., dan Kozukue, N. (2009). Stability of green tea catechins in commercial tea leaves during storage for 6 months. Journal of Food Science, 74 (2).
- Han, Z. X., Rana, M. M., Liu, G. F., Gao, M. J., Li, D. X., Wu, F. G., dan Wei, S. (2016). *Green tea flavour determinants and their changes over manufacturing processes*. Food Chemistry, 212, 739–748.
- Ho, C.-T., Zheng, X., dan Li, S. (2015). Tea aroma formation. Food Science and Human Wellness, 4(1), 9–27.
- Kato, M., dan Shibamoto, T. (2001). Variation of major volatile constituents in various green teas from southeast Asia. Journal of Agricultural and Food Chemistry, 49(3), 1394–1396.

- Kawakami, M., dan Yamanishi, T. (1999). *Formation of aroma components in roasted or pan-fired green tea by roasting or pan-firing treatment*. Biosci Biotechnol Agrochem. 73, 893–906.
- Kuo, P.C., Y.Y. Lai, dan Y.J. Chen, (2011) Changes in volatile compounds upon aging and drying in oolong tea production, J. Sci. Food Agric. 91 (2011) 293–301.
- Lau, H., Liu, S. Q., Xu, Y. Q., Tan, L. P., Zhang, W. L., Lassabliere, B., dan Yu, B. (2018). Characterising volatiles in tea (*Camellia sinensis*). Part II: Untargeted and targeted approaches to multivariate analysis. Journal of Food Chemistry, 94(December 2017), 142–162.
- Liang, Y. R., Ye, Q., Jin, J., Liang, H., Lu, J. L., Du, Y. Y., dan Dong, J. J. (2008). Chemical and instrumental assessment of green tea sensory preference. International Journal of Food Properties, 11(2), 258–272.
- Liu, Y., Luo, L., Liao, C., Chen, L., Wang, J., dan Zeng, L. (2018). Effects of brewing conditions on the phytochemical composition, sensory qualities and antioxidant activity of green tea infusion: A study using response surface methodology. Journal of Food Chemistry, 269(2), 24–34.
- Lv, H.-P., Zhong, Q.-S., Lin, Z., Wang, L., Tan, J.-F., dan Guo, L. (2012). Aroma characterisation of Pu-erh tea using headspace-solid phase microextraction combined with GC/MS and GC-olfactometry. Food Chemistry, 130(4), 1074–1081.
- Ma, C., Li, J., Chen, W., Wang, W., Qi, D., Pang, S., dan Miao, A. (2018). Study of the aroma formation and transformation during the manufacturing process of oolong tea by solid-phase micro-extraction and gas chromatography mass spectrometry combined with chemometrics. Food Research International, 108(December 2017), 413–422.
- Mizukami, Y., Sawai, Y., dan Yamaguchi, Y. (2008). Changes in the concentrations of acrylamide, selected odorants, and catechins caused by roasting of green tea. Journal of Agricultural and Food Chemistry, 56(6), 2154–2159.
- Mu, B., Zhu, Y., Lv, H. P., Yan, H., Peng, Q. H., dan Lin, Z. (2018). The enantiomeric distributions of volatile constituents in different tea cultivars. Food Chemistry, 265(May), 329–336.
- Narukawa, M., Toda, Y., Nakagita, T., Hayashi, Y., dan Misaka, T. (2014). L Theanine elicits umami taste via the T1R1 + T1R3 umami taste receptor. Amino Acids, 46(6), 1583–1587.
- Ofmannt, T.H, (2006). Agricultural And Molecular and Sensory Studies on the Umami Taste of Tea, Journal of Food Science and Technology, 2688–2694.
- Rohdiana, Dadan dan W. Tantan. (2004). Aktifitas Antioksidan Beberapa Klon Teh Unggulan. Prosiding Seminar Nasional dan Kongres Perhimpunan Ahli Teknologi Pangan Indonesia (PATPI). 17-18 Desember. Jakarta.
- S. Tsuge, H. Ohtani, C. dan Waeanaabe,. (2011). Pyrolysis-GC/MS data book of synthetic polymers: pyrograms, thermograms and MS of pyrolyzates, 1st ed., Elsevier, Amsterdam.
- Sang, S. (2016). Tea : Chemistry and Processing. Encyclopedia of Food and Health Page 268 – 272.

- Sasaki, T., Koshi, E., Take, H., Michihata, T., Maruya, M., dan Enomoto, T. (2017). Characterisation of odorants in roasted stem tea using gas chromatography-mass spectrometry and gas chromatography olfactometry analysis. *Food Chemistry*, 220, 177–183.
- Sasaki, T., Koshi, E., Take, H., Michihata, T., Maruya, M., dan Enomoto, T. (2017). Characterisation of odorants in roasted stem tea using gas chromatography-mass spectrometry and gas chromatography olfactometry analysis. *Food Chemistry*, 220, 177–183.
- Schuh, C., dan Schieberle, P. (2006). Characterization of the key aroma compounds in the beverage prepared from Darjeeling black tea: Quantitative differences between tea leaves and infusion. *Journal of Agricultural and Food Chemistry*, 54(3), 916–924.
- Setyamidjaja, D. (2000). Teh, Budidaya dan Pengolahan Pasca Panen. *Kanisius*. Yogyakarta.
- Shukla, Y. (2007). Tea and Cancer Chemoprevention : A Comprehensive Riview. *Asian Pasific Journal of Cancer Preventiom*, 8, 155-166
- Siswoputranto, P.S. 1 978. Perkembangan Teh, Kopi, Cokelat Internasional. *Gramedia*. Jakarta.
- Supartono, W., Sukartiko, A. C., Yuliando, H., & Kristanti, N. E. (2015). Possibility of some indigenous spices as flavor agent of green tea. *Italian Oral Surgery*, 3, 62–66
- Türközü, D., dan Şanlıer, N. (2017). L-theanine, unique amino acid of tea, and its metabolism, health effects, and safety. *Critical Reviews in Food Science and Nutrition*, 57(8), 1681–1687
- Wang, C., Zhang, C., Kong, Y., Peng, X., Li, C., Liu, S., dan Xu, Y. (2017). A comparative study of volatile components in Dianhong teas from fresh leaves of four tea cultivars by using chromatography-mass spectrometry, multivariate data analysis, and descriptive sensory analysis. *Food Research International*, 100(June), 267–275.
- Wang, D., Kubota. K., Kobayashi. A., dan Juan. I. M. (2001). Analysis of Glycosidic Bound Aroma Precursor in Tea Leaves: Changes of in the Glycosidic Content of Tea Leaves During the Oolong Tea Manufacturing Process. *Food Chemistry*, 49: 5391-5396
- Wang, L. F., Lee, J. Y., Chung, J. O., Baik, J. H., So, S., dan Park, S. K. (2008). Discrimination of teas with different degrees of fermentation by SPME GC analysis of the characteristic volatile flavour compounds. *Food Chemistry*, 109(1), 196–206.
- Wang, L., Xu, R., Hu, B., Li, W., Sun, Y., Tu, Y., dan Zeng, X. (2010). Analysis of free amino acids in Chinese teas and flower of tea plant by high performance liquid combined with solid-phase extraction. *Food Chemistry*, 123(4), 1259–1266.
- Wang, T., Li, X., Yang, H., Wang, F., Kong, J., Qiu, D., dan Li, Z. (2018). Mass spectrometry-based metabolomics and chemometric analysis of Pu-erh teas of various origins. *Food Chemistry*, 268(June), 271–278
- Wang, Y., Zheng, P., Liu, P., Song, X., Guo, F., dan Li, Y. (2019). Novel insight into the role of withering process in characteristic flavor formation of teas using transcriptome analysis and metabolite profiling. *Food Chemistry*, 272(June 2018), 313–322.
- Yang, D., Liu, Y., Sun, M., Zhao, L., Wang, Y., Chen, X., dan Xia, T. (2012). Differential gene expression in tea (*Camellia sinensis* L.) call with different morphologies and catechin contents. *Journal of Plant Physiology*, 169(2), 163–175.



UNIVERSITAS
GADJAH MADA

Kontribusi Penambahan Putih Telur dalam Pembentukan Senyawa Aroma serta Potensi Antioksidan pada Teh Hijau

MUHAMMAD SYAFI'IE, Dr. Ir. Supriyadi, M.Sc. ; Prof. Dr. Ir. Sri Anggrahini, M.S.

Universitas Gadjah Mada, 2022 | Diunduh dari <http://etd.repository.ugm.ac.id/>

- Yang, Y., Zhang, M., Yin, H., Deng, Y., Jiang, Y., Yuan, H., dan Wang, J. (2018). LWT - Food Science and Technology Rapid profiling of volatile compounds in green teas using Micro-Chamber / Thermal Extractor combined with thermal desorption coupled to gas chromatography-mass spectrometry followed by multivariate statistical analysis. *Journal of Food Science and Technology*, 96(January), 42–50.
- Yashin, A. Y., Nemzer, B. V., Combet, E., dan Yashin, Y. I. (2015). Determination of the Chemical Composition of Tea by Chromatographic Methods: A Review. *Journal of Food Research*, 4(3), 56.
- Yin, J., Zhang, Y., Du, Q., Chen, J., Yuan, H., & Xu, Y. (2014). Effect of Ca²⁺ concentration on the tastes from the main chemicals in green tea infusions. *Food Chemistry*, 62, 941–946.