

DAFTAR PUSTAKA

1. Aneja, R.K., Carcillo, J.A. 2011. Differences between adult and pediatric septic shock. *Minerva Anesthesiol*, 77(10): 986–992.
2. AP, G., DLS, A., PSB, M., VH, I. 2016. Anti-Inflammatory Cytokines in Sepsis: Immunological Studies and In Silico Investigation. *J Pharm Care Heal Syst*, 3(3).
3. Anwary, A.Z., Hakim, D.D.L., Setiabudi, D. (2021) ‘Korelasi positif antara neutrophil lymphocyte count ratio dan C-reactive protein pada pasien sepsis anak’, *Sari Pediatri*, 23(1)
4. Anthony, D.A., Andrews, D.M., Watt, S.V., Trapani, J.A., & Smyth, M.J. (2010) ‘Functional dissection of the granzyme family: cell death and inflammation’, *Immunology*, Rev.235, pp. 73–92
5. Cabrera-Perez, J., Condotta, S.A., Badovinac, V.P., Griffith, T.S. 2014. Impact of sepsis on CD4 T cell immunity. *J Leukoc Biol*, 96(5): 767–777.
6. Celikbilek, M., Dogan, S., Ozbakir, O., Zararsiz, G., Küçük, H., Gürsoy, S., Yurci, A., Güven, K., Yücesoy, M. 2013. Neutrophil-Lymphocyte Ratio as a Predictor of Disease Severity in Ulcerative Colitis. *J Clin Lab Anal*, 27(1): 72–76.
7. Cruz, A.T., Lane, R.D., Balamuth, F., Aronson, P.L., Ashby, D.W., Neuman, M.I., Souganidis, E.S., Alpern, E.R., Schlapbach, L.J. 2020. Updates on pediatric sepsis. *J Am Coll Emerg Physicians Open*, 1(5): 981–993.
8. Dahlan, S. 2017. *Statistik Untuk Kedokteran Dan Kesehatan Edisi 6*. Jakarta.p258
9. Desy D, Antonius HP, Mulyadi M, Bambang S. Faktor risiko yang berperan pada mortalitas sepsis. *Sari Pediatri* 2014;15;281-8 De Souza, D. C., Barreira, E. R., & Faria, L. S. (2017) ‘The epidemiology of sepsis in childhood’, *Shock*, 47(:2-5).
10. Dellmann, H.D. & Eurell, J.A. (1998) *Textbook of Veterinary Histology*, Lippincott Williams & Wilkins, Maryland.
11. De Souza, D.C., Barreira, E.R., Faria, L.S. 2017. The epidemiology of sepsis in childhood. *Shock*, 47: 2–5.
12. Dellinger, R.P., Levy, M.M., Carlet, J.M., Bion, J., Parker, M.M., Jaeschke, R., Reinhart, K., Angus, D.C., Brun-Buisson, C., Beale, R., Calandra, T., Dhainaut, J.F., Gerlach, H., Harvey, M., Marini, J.J., Marshall, J., Ranieri, M., Ramsay, G., Sevransky, J., Thompson, B.T., Townsend, S., Vender, J.S., Zimmerman, J.L., Vincent, J.L. 2008. Surviving sepsis campaign: International guidelines for management of severe sepsis and septic shock: 2008. *Crit Care Med*, 36(1): 296–327.
13. Dursun, A., Ozsoylu, S., Akyildiz, B.N. 2018. Neutrophil-to-lymphocyte ratio and mean platelet volume can be useful markers to predict sepsis in children. *Pakistan J Med Sci*, 34(4): 918–922.
14. El-Nawawy, A., Mohsen, A.A., Abdel-Malik, M., Taman, S.O. 2017. Performance of the pediatric logistic organ dysfunction (PELOD) and (PELOD-2) scores in a pediatric intensive care unit of a developing country. *Eur J Pediatr*, 176(7): 849–855.
15. Elshinawy, M., Kamal, M., Nazir, H., Khater, D., Hassan, R., *et al.* (2020) ‘Sepsis-related anemia in a pediatric intensive care unit: transfusion-associated outcomes’, *Transfusion*, 60(Suppl 1), S4-S9.
16. Fialkow, L., Filho, L.F., Bozzetti, M.C., Milani, A.R., Rodrigues Filho, E.M., Ladniuk, R.M., Pierozan, P., de Moura, R.M., Prolla, J.C., Vachon, E., Downey, G.P. 2006. Neutrophil apoptosis: A marker of disease severity in sepsis and sepsis-induced acute respiratory distress syndrome. *Crit Care*, 10(6): 1–14.
17. Funk, D.J., Parrillo, J.E., Kumar, A. 2009. Sepsis and Septic Shock: A History. *Crit Care Clin*, 25(1): 83–101.

18. Gacouin, A., Tribut, O., Amiot, L., Tattevin, P. 2002. Shock Is Associated With Poor Outcome. , 18(6): 487–494.
19. Goldstein, B., Giroir, B., Randolph, A. 2005. International pediatric sepsis consensus conference: Definitions for sepsis and organ dysfunction in pediatrics. In *Pediatric Critical Care Medicine*.
20. Gomes, A.P., Alves, D.L.S., Miguel, P.S.B., Inoue, V.H., Correia Lopes, T.C., *et al.* (2016) ‘Anti-Inflammatory Cytokines in Sepsis: Immunological Studies and In Silico Investigation’, *Journal of Pharmaceutical Care & Health Systems*, 3(3).
21. Greg S Martin (2012) Sepsis, severe sepsis and septic shock: changes in incidence, pathogens and outcomes, *Expert Review of Anti-infective Therapy*, 10:6, 701-706.
22. Gurol G., Citci I.H., Terzi H.A., Atasoy A.R., Ozbek A., *et al.* (2015) ‘Are there standardized cutoff values for neutrophil-lymphocyte ratios in bacteremia or sepsis?’, *Journal of Microbiology biotechnology*, 25(4), pp. 521-525.
23. Guyton, A.C. (1997) *Buku Ajar Fisiologi Kedokteran*, Edisi 9, Irawati Setiawan, Penerjemah, ECG Penerbit Buku Kedokteran, Jakarta.
24. Hall, M.K., Kea, B., Wang, R. 2019. Recognising Bias in Studies of Diagnostic Tests Part 1: Patient Selection. *Emerg Med J*, 36(7): 431–434. <https://emj.bmj.com/content/36/7/431>.
25. Holub, M., Beran, O., Kaspríková, N., Chalupa, P. 2012. Neutrophil to lymphocyte count ratio as a biomarker of bacterial infections. *Cent Eur J Med*, 7(2): 258–261.
26. Ibrahim, M. K., Zambruni, M., Melby, C. L., & Melby, P. C. (2017). Impact of Childhood Malnutrition on Host Defense and Infection. *Clinical Microbiology Reviews*, 30(4), 919–971
27. Kusumo, P.D. 2012. Gangguan Imunodefisiensi Primer (PID). *Widya Kedokt*, 29(324): 14–22.
28. Maat M., Buysse C.M., Emonts M., Spanjaard L., Joosten K.F., *et al.* (2007) ‘Improved survival of children with sepsis and purpura: effects of age, gender, and era’, *Critical Care*, 11: pp. 1-10.
29. Martin G.S., Eaton S., Mealer M., & Moss M. (2012) ‘Extravascular lung water in patients with severe sepsis: a prospective cohort study’, *Critical Care*, 9(2): pp. 74-82.
30. Mathews, S., Rajan, A., & Soans, S.T. (2019) ‘Prognostic value of rise in neutrophil to lymphocyte ratio (NLR) and platelet to lymphocyte ratio (PLR) in predicting the mortality in paediatric intensive care’, *International Journal of Contemporary Pediatrics*, 6(3).
31. Maqbul PL, Leksana E, Harahap MS. Rasio Netrofil Limfosit dan Limfositopenia Sebagai Penanda Sepsis. *JAI (Jurnal Anestesiologi Indonesia)* [Online]. 2016 Jul;8(2):114-112.
32. Latief, A., Chairulfatah, A., Alam, A., Pudjiadi, A., Malisie, R., Hadinegoro, S. 2016. *Konsensus Diagnosa dan Tata Laksana Sepsis pada Anak*.
33. Mayr, F.B., Yende, S., Angus, D.C. 2014. Epidemiology of severe sepsis. *Virulence*, 5(1): 4–11.
34. Melda, M., Triasih, R., Nurnaningsih. 2021. Modifying the pelod-2 score to predict mortality in critically ill patients. *Paediatr Indones Indones*, 61(2): 61–68.
35. Midwinter, M.J. 2011. Management of sepsis. In *Fundamentals of Surgical Practice, Thrid Edition*. 248–275.
36. Oematan, Y., Manoppo, J.I.C., Runtunuwu, A.L. 2013. Peran Inflamasi Dalam Patofisiologi Sepsis Dan Syok Septik Pada Anak. *J Biomedik*, 1(3): 166–173.
37. Prasetyo, R.H. 2011. Clinical Pathology and Majalah Patologi Klinik Indonesia dan Laboratorium Medik. *J Indones*, 21(3): 261–265. <http://journal.unair.ac.id/download->

38. Rosales, C. 2018. Neutrophil: A cell with many roles in inflammation or several cell types, *Front Physiol*, 9(FEB): 1–17.
39. Rusmawatiningtyas, D., Nurnaningsih, N. 2017. Mortality rates in pediatric septic shock. *Paediatr Indones*, 56(5): 304.
40. Saputra, I.M.Y., Gustawan, W., Utama, M.D., Arhana, B. 2019. Rasio Neutrofil dan Limfosit (NLCR) Sebagai Faktor Risiko Terjadinya Infeksi Bakteri di Ruang Rawat Anak RSUP Sanglah Denpasar. *Sari Pediatr*, 20(6): 354.
41. Schroder J, Kahlke V, Staubach KH, Zabel P, Stuber F. Gender differences in human sepsis. *Arch Surg*. 1998;133:1200-5
42. Schulze-Koops, H., Davis, L.S., Haverty, T.P., Wacholtz, M.C., Lipsky, P.E. 1998. Reduction of Th1 cell activity in the peripheral circulation of patients with rheumatoid arthritis after treatment with a non-depleting humanized monoclonal antibody to CD4. *J Rheumatol*, 25(11): 2065–2076.
43. Silalahi, B., Ginting, F., Ginting, Y. (2015) *Hubungan Antara Neutrophil-Lymphocyte Count Ratio dengan Kadar Procalcitonin pada Pasien Sepsis*. Universitas Sumatera Utara, Medan.
44. Silva, E., Da Hora Passos, R., Ferri, M.B., De Figueiredo, L.F.P. 2008. Sepsis: From bench to bedside. *Clinics*, 63(1): 109–120.
45. Soedarmo, S.S.P., Garna, H., Hadinegoro, S.R.S., Satari, H.I. 2008. *Buku Ajar Infeksi & Pediatri Tropis*.
46. Suari, N.M.R., Latief, A., Pudjiadi, A.H. 2021. New pelod-2 cut-off score for predicting death in children with sepsis. *Paediatr Indones*, 61(1): 39
47. Sudiartha, I.P.G., Wiargitha, I.K., Mahadewa, T.G.B. (2020) ‘Perbedaan nilai neutrophil lymphocyte ratio (NLR) terhadap pemeriksaan kultur darah dalam mendiagnosis sepsis pada pasien peritonitis di RSUP Sanglah, Bali, Indonesia’, *Intisari Sains Medis*, 11(1).
48. Shorr, A., Micek, S., Welch, E., Doherty, J., Reichley, R., *et al.* (2010) ‘Inappropriate antibiotic therapy in gram-negative sepsis increases hospital length of stay’, *Critical Care Medicine*, 39(1), pp. 46-51.
49. Tan, B., Wong, J.J., Sultana, R., Koh, J.C.J.W., Jit, M., *et al.* (2019) ‘Global case-fatality rates in pediatric severe sepsis and septic shock: A systematic review and meta analysis’, *Jama Pediatric*, 173(4), pp. 352-62.
50. Wulandari, A., Martuti, S., Kaswadi, P. 2018. Perkembangan diagnosis sepsis pada anak. *Sari Pediatr*, 19(4): 237.
51. Yazici, M., Özkisacik, S., Onur Öztan, M., Gürsoy, H. 2010. Neutrophil/lymphocyte ratio in the diagnosis of childhood appendicitis. *Turk J Pediatr*, 52(4): 400–403.
52. Yuanita, Anum, Q. 2012. Peranan Toll-Like Receptor Terhadap Infeksi Bakteri Pada Kulit. *Mdvi*, 39(1): 42–48.
53. Zahorec, R. 2001. Ratio of neutrophil to lymphocyte counts--rapid and simple parameter of systemic inflammation and stress in critically ill. *Bratisl Lek Listy*, 102(1): 5–14.