

DAFTAR PUSTAKA

- Tiao MM, Tsai SS, Kuo HW, Chen CL, Yang CY. Epidemiological features of biliary atresia in Taiwan, a national study 1996-2003. *J Gastroenterol Hepatol*. 2008 Jan;23(1):62-6.
- Mathur, P., Gupta, R., Soni, V., Ahmed, R. and Goyal, R.B., 2014. Biliary atresia associated with polysplenia syndrome, dextrocardia, situs inversus totalis and malrotation of intestines. *Journal of neonatal surgery*, 3(1).
- Vij, M. and Rela, M., 2020. Biliary atresia: pathology, etiology and pathogenesis. *Future Science OA*, 6(5), p.FSO466. DOI : <https://dx.doi.org/10.2144%2Ffsoa-2019-0153>
- Kumar, R., Lal, B. B., Sood, V., Khanna, R., Kumar, S., Bharathy, K., & Alam, S. (2019). Predictors of Successful Kasai Portoenterostomy and Survival with Native Liver at 2 Years in Infants with Biliary Atresia. *Journal of clinical and experimental hepatology*, 9(4), 453–459. <https://doi.org/10.1016/j.jceh.2018.09.008>
- Baek, S. H., Kang, J. M., Ihn, K., Han, S. J., Koh, H., & Ahn, J. G. (2020). The Epidemiology and Etiology of Cholangitis After Kasai Portoenterostomy in Patients With Biliary Atresia. *Journal of pediatric gastroenterology and nutrition*, 70(2), 171–177. <https://doi.org/10.1097/MPG.0000000000002555>
- Liu, J., Dong, R., Chen, G., Dong, K., & Zheng, S. (2019). Risk factors and prognostic effects of cholangitis after Kasai procedure in biliary atresia patients: A retrospective clinical study. *Journal of pediatric surgery*, 54(12), 2559–2564. <https://doi.org/10.1016/j.jpedsurg.2019.08.026>
- Dewanti, A.Y., Makhmudi, A., Gunadi, dan Nuryastuti, T., 2015. Profil Pasien Atresia Bilier di RSUP dr. Sardjito Yogyakarta. Skripsi Sarjana Kedokteran pada Fakultas Kedokteran UGM Yogyakarta.
- Russo P, Magee JC, Boitnott J, et al. Design and validation of the biliary atresia research consortium histologic assessment system for cholestasis in infancy. *Clinical Gastroenterology and Hepatology : the Official Clinical Practice Journal of the American Gastroenterological Association*. 2011 Apr;9(4):357-362.e2. DOI: <https://doi.org/10.1016/j.cgh.2011.01.003>
- Russo, P., Magee, J.C., Anders, R.A., Bove, K.E., Chung, C., Cummings, O.W., Finegold, M.J., Finn, L.S., Kim, G.E., Lovell, M.A. and Magid, M.S., 2016. Key histopathological features of liver biopsies that distinguish biliary atresia from other causes of infantile cholestasis and their correlation with outcome:

A multicenter study. *The American journal of surgical pathology*, 40(12), p.1601. DOI: <https://dx.doi.org/10.1097%2FPA.0000000000000755>

Arjmand, A., Tsipouras, M.G., Tzallas, A.T., Forlano, R., Manousou, P. and Giannakeas, N., 2020. Quantification of liver fibrosis—A comparative study. *Applied Sciences*, 10(2), p.447. DOI: <https://doi.org/10.3390/app10020447>

Yamataka, A., Kato, Y., & Miyano, T. (2010). *BILIARY TRACT DISORDERS AND PORTAL HYPERTENSION. Ashcraft's Pediatric Surgery*, 557–577. DOI: <https://doi.org/10.1016/b978-1-4160-6127-4.00044-6>

Sinha, C. K., & Davenport, M. (2008). Biliary atresia. *Journal of Indian Association of Pediatric Surgeons*, 13(2), 49–56. <https://doi.org/10.4103/0971-9261.43015>

Fanna, M., Masson, G., Capito, C., Girard, M., Guerin, F., Hermeziu, B., Lachaux, A., Roquelaure, B., Gottrand, F., Broue, P., Dabadie, A., Lamireau, T., Jacquemin, E., & Chardot, C. (2019). Management of Biliary Atresia in France 1986 to 2015: Long-term Results. *Journal of pediatric gastroenterology and nutrition*, 69(4), 416–424. <https://doi.org/10.1097/MPG.0000000000002446>

Khalil, B. A., Perera, M. T., & Mirza, D. F. (2010). Clinical practice: management of biliary atresia. *European journal of pediatrics*, 169(4), 395–402. <https://doi.org/10.1007/s00431-009-1125-7>

Serinet, M. O., Wildhaber, B. E., Broué, P., Lachaux, A., Sarles, J., Jacquemin, E., Gauthier, F., & Chardot, C. (2009). Impact of age at Kasai operation on its results in late childhood and adolescence: a rational basis for biliary atresia screening. *Pediatrics*, 123(5), 1280–1286. <https://doi.org/10.1542/peds.2008-1949>

Wildhaber, B.E., 2012. Biliary atresia: 50 years after the first Kasai. *International Scholarly Research Notices*, 2012. <https://doi.org/10.5402/2012/132089>

Shneider, B. L., Magee, J. C., Karpen, S. J., Rand, E. B., Narkewicz, M. R., Bass, L. M., Schwarz, K., Whittington, P. F., Bezerra, J. A., Kerkar, N., Haber, B., Rosenthal, P., Turmelle, Y. P., Molleston, J. P., Murray, K. F., Ng, V. L., Wang, K. S., Romero, R., Squires, R. H., Arnon, R., ... Childhood Liver Disease Research Network (ChiLDReN) (2016). Total Serum Bilirubin within 3 Months of Hepatopertoenterostomy Predicts Short-Term Outcomes in Biliary Atresia. *The Journal of pediatrics*, 170, 211–7.e72. <https://doi.org/10.1016/j.jpeds.2015.11.058>

Yokoyama, H., Masaki, T., Inoue, I., Nakamura, M., Mezaki, Y., Saeki, C. Matsuura, T. (2019). *Histological and biochemical evaluation of transforming*

growth factor- β activation and its clinical significance in patients with chronic liver disease. *Heliyon*, 5(2), e01231. <https://doi.org/10.1016/j.heliyon.2019.e01231>

Gunadi, Gunawan, T. A., Widiyanto, G., Yuanita, A., Mulyani, N. S., & Makhmudi, A. (2018). Liver transplant score for prediction of biliary atresia patients' survival following Kasai procedure. *BMC research notes*, 11(1), 381. <https://doi.org/10.1186/s13104-018-3498-z>

Baek, S. H., Kang, J. M., Ihn, K., Han, S. J., Koh, H., & Ahn, J. G. (2020). The Epidemiology and Etiology of Cholangitis After Kasai Portoenterostomy in Patients With Biliary Atresia. *Journal of pediatric gastroenterology and nutrition*, 70(2), 171–177. <https://doi.org/10.1097/MPG.0000000000002555>

Nakajima, H., Koga, H., Okawada, M., Nakamura, H., Lane, G. J., & Yamataka, A. (2018). Does time taken to achieve jaundice-clearance influence survival of the native liver in post-Kasai biliary atresia?. *World journal of pediatrics : WJP*, 14(2), 191–196. <https://doi.org/10.1007/s12519-018-0139-5>

Namasemayam, D., & Nallusamy, M. (2017). Factors influencing outcome after hepatic portoenterostomy among extrahepatic bile duct atresia patients in Hospital Sultanah Bahiyah, Alor Setar. *The Medical journal of Malaysia*, 72(6), 329–332.

Buzzetti, E., Parikh, P. M., Gerussi, A., & Tsochatzis, E. (2017). Gender differences in liver disease and the drug-dose gender gap. *Pharmacological research*, 120, 97–108. <https://doi.org/10.1016/j.phrs.2017.03.014>

Guy, J., & Peters, M. G. (2013). Liver disease in women: the influence of gender on epidemiology, natural history, and patient outcomes. *Gastroenterology & hepatology*, 9(10), 633–639.

Shen, W. J., Chen, G., Wang, M., & Zheng, S. (2019). Liver fibrosis in biliary atresia. *World journal of pediatrics : WJP*, 15(2), 117–123. <https://doi.org/10.1007/s12519-018-0203-1>

Haafiz A. B. (2010). Liver fibrosis in biliary atresia. *Expert review of gastroenterology & hepatology*, 4(3), 335–343. <https://doi.org/10.1586/egh.10.29>

Almpanis, Z., Demonakou, M. and Tiniakos, D., 2016. Evaluation of liver fibrosis: “Something old, something new...”. *Annals of Gastroenterology: Quarterly Publication of the Hellenic Society of Gastroenterology*, 29(4), p.445.

- alla, I.A.S. and Wibawa, I.D.N., 2012. Pendekatan diagnosis dan terapi fibrosis hati. *Denpasar: Bagian/SMF Ilmu Penyakit Dalam FK Universitas Udayana/RS Sanglah*.
- Henwood, A., 2010. Microscopic quality control of haematoxylin and eosin—know your histology. *Connection*, 14, pp.115-120.
- Goodman, Z.D., 2007. Grading and staging systems for inflammation and fibrosis in chronic liver diseases. *Journal of hepatology*, 47(4), pp.598-607.
- Obayashi, J., Kawaguchi, K., Manabe, S., Nagae, H., Wakisaka, M., Koike, J., Takagi, M. and Kitagawa, H., 2017. Prognostic factors indicating survival with native liver after Kasai procedure for biliary atresia. *Pediatric surgery international*, 33(10), pp.1047-1052. <https://doi.org/10.1007/s00383-017-4135-y>
- Baruah, R.R., Bhatnagar, V., Agarwala, S. and Gupta, S.D., 2015. Correlation of pre-and post-operative liver function, duct diameter at porta hepatis, and portal fibrosis with surgical outcomes in biliary atresia. *Journal of Indian Association of Pediatric Surgeons*, 20(4), p.184. <https://doi.org/10.4103/0971-9261.161040>
- Goda, T., Kawahara, H., Kubota, A., Hirano, K., Umeda, S., Tani, G., Ishii, T., Tazuke, Y., Yoneda, A., Etani, Y. and Ida, S., 2013. The most reliable early predictors of outcome in patients with biliary atresia after Kasai's operation. *Journal of pediatric surgery*, 48(12), pp.2373-2377. <https://doi.org/10.1016/j.jpedsurg.2013.08.009>
- Robie, D.K., Overfelt, S.R. and Xie, L., 2014. Differentiating biliary atresia from other causes of cholestatic jaundice. *The American Surgeon*, 80(9), pp.827-831.
- Nguyen, A.H.P., Pham, Y.H.T., Vu, G.H., Nguyen, M.H., Hoang, T.N. and Holterman, A., 2021. Biliary atresia liver histopathological determinants of early post-Kasai outcome. *Journal of Pediatric Surgery*. <https://doi.org/10.1016/j.jpedsurg.2021.03.039>
- Gad, E.H., Kamel, Y., Salem, T.A.H., Ali, M.A.H. and Sallam, A.N., 2021. Short- and long-term outcomes after Kasai operation for type III biliary atresia: Twenty years of experience in a single tertiary Egyptian center-A retrospective cohort study. *Annals of Medicine and Surgery*, 62, pp.302-314. <https://doi.org/10.1016/j.amsu.2021.01.052>
- Czubkowski, P., Cielecka-Kuszyk, J., Rurarz, M., Kaminska, D., Markiewicz-Kijewska, M. and Pawlowska, J., 2016. The limited prognostic value of liver

histology in children with biliary atresia. *Annals of hepatology*, 14(6), pp.902-909. <https://doi.org/10.5604/16652681.1171781>

Shiha, G. and Zalata, K., 2011. Ishak versus METAVIR: terminology, convertibility and correlation with laboratory changes in chronic hepatitis C. *Liver biopsy*, 10, pp.155-170.

Abdalla AF, Zalata KR, Ismail AF, et al. Regression of fibrosis in pediatric autoimmune hepatitis: morphometric assessment of fibrosis versus semiquantitative methods. *Fibrogenesis Tissue Repair* 2009;2:2. <https://doi.org/10.1186/1755-1536-2-2>

Goodman ZD. Grading and staging systems for inflammation and fibrosis in chronic liver diseases. *J Hepatol* 2007;47:598-607. <https://doi.org/10.1016/j.jhep.2007.07.006>

Nallagangula, K.S., Nagaraj, S.K., Venkataswamy, L. and Chandrappa, M., 2017. Liver fibrosis: a compilation on the biomarkers status and their significance during disease progression. *Future science OA*, 4(1), p.FSO250. <https://doi.org/10.4155/fsoa-2017->

Shao, L.N., Zhang, S.T., Wang, N., Yu, W.J., Chen, M., Xiao, N., Duan, Y., Pan, L.Z., Song, W.Q., Xia, Y.X. and Zhang, L., 2020. Platelet indices significantly correlate with liver fibrosis in HCV-infected patients. *Plos one*, 15(1), p.e0227544. <https://doi.org/10.1371/journal.pone.0227544>

Yokoo, T., Wolfson, T., Iwaisako, K., Peterson, M.R., Mani, H., Goodman, Z., Changchien, C., Middleton, M.S., Gamst, A.C., Mazhar, S.M. and Kono, Y., 2015. Evaluation of liver fibrosis using texture analysis on combined-contrast-enhanced magnetic resonance images at 3.0 T. *BioMed research international*, 2015. <https://doi.org/10.1155/2015/387653>

Degtyareva, A., Razumovskiy, A., Kulikova, N., Ratnikov, S., Filippova, E., Gordeeva, E., Albegova, M., Rebrikov, D. and Puchkova, A., 2020. Long-Term Effects of Kasai Portoenterostomy for Biliary Atresia Treatment in Russia. *Diagnostics*, 10(9), p.686. <https://doi.org/10.3390/diagnostics10090686>