

## DAFTAR PUSTAKA

- Adrie, C., Azoulay, E., Francois, A., Clec'h, C., Darques, L., Schwebel, C., Nakache, D., Jamali, S., Goldgran-Toledano, D., Garrouste-Orgeas, M. and Timsit, J.F., 2007. Influence of gender on the outcome of severe sepsis: a reappraisal. *Chest*, 132(6), pp.1786-1793.
- Adrie, C., Francois, A., Alvarez-Gonzalez, A., Mounier, R., Azoulay, E., Zahar, J.R., Clec'h, C., Goldgran-Toledano, D., Hammer, L., Descorps-Declere, A. and Jamali, S., 2009. Model for predicting short-term mortality of severe sepsis. *Critical Care*, 13(3), pp.1-14.
- Angele, M.K., Pratschke, S., Hubbard, W.J. and Chaudry, I.H., 2014. Gender differences in sepsis: cardiovascular and immunological aspects. *Virulence*, 5(1), pp.12-19.
- Aune, D., Huang, W., Nie, J. and Wang, Y., 2021. Hypertension and the risk of all-cause and cause-specific mortality: an outcome-wide association study of 67 causes of death in the national health interview survey. *BioMed research international*, 2021.
- Berg, D. and Gerlach, H., 2018. Recent advances in understanding and managing sepsis. *F1000Research*, 7.
- Bouza, C. and López-Cuadrado, T., 2020. Epidemiology and trends of sepsis in young adults aged 20–44 years: a nationwide population-based study. *Journal of clinical medicine*, 9(1), p.77.
- Brady, J., Horie, S. and Laffey, J.G., 2020. Role of the adaptive immune response in sepsis. *Intensive Care Medicine Experimental*, 8(1), pp.1-19.
- Brown, R.L., Sequeira, R.P. and Clarke, T.B., 2017. The microbiota protects against respiratory infection via GM-CSF signaling. *Nature communications*, 8(1), pp.1-11.
- Chaudhry, H., Zhou, J., Zhong, Y.I.N., Ali, M.M., McGuire, F., Nagarkatti, P.S. and Nagarkatti, M., 2013. Role of cytokines as a double-edged sword in sepsis. *In vivo*, 27(6), pp.669-684.
- Chen, C.M., Cheng, K.C., Chan, K.S. and Yu, W.L., 2014. Age may not influence the outcome of patients with severe sepsis in intensive care units. *International Journal of Gerontology*, 8(1), pp.22-26.
- Chung, K.P., Chang, H.T., Lo, S.C., Chang, L.Y., Lin, S.Y., Cheng, A., Huang, Y.T., Chen, C.C., Lee, M.R., Chen, Y.J. and Hou, H.H., 2015. Severe lymphopenia is associated with elevated plasma interleukin-15 levels and increased mortality during severe sepsis. *Shock*, 43(6), pp.569-575.

- Condotta, S.A., Cabrera-Perez, J., Badovinac, V.P. and Griffith, T.S., 2013. T-cell-mediated immunity and the role of TRAIL in sepsis-induced immunosuppression. *Critical Reviews™ in Immunology*, 33(1).
- Darwis, I. and Probosuseno, P., 2019. Hubungan neutrophil lymphocyte ratio dengan outcome sepsis pada geriatri. *Jurnal Kedokteran Universitas Lampung*, 3(1), pp.147-153.
- de Pablo, R., Monserrat, J., Prieto, A. and Alvarez-Mon, M., 2014. Role of circulating lymphocytes in patients with sepsis. *BioMed research international*, 2014.
- Desruisseaux, M.S., Nagajyothi, Trujillo, M.E., Tanowitz, H.B. and Scherer, P.E., 2007. Adipocyte, adipose tissue, and infectious disease. *Infection and immunity*, 75(3), pp.1066-1078.
- Drewry, A.M., Samra, N., Skrupky, L.P., Fuller, B.M., Compton, S.M. and Hotchkiss, R.S., 2014. Persistent lymphopenia after diagnosis of sepsis predicts mortality. *Shock (Augusta, Ga.)*, 42(5), p.383.
- Evans, T., 2018. Diagnosis and management of sepsis. *Clinical Medicine*, 18(2), p.146.
- Forget, P., Khalifa, C., Defour, J.P., Latinne, D., Van Pel, M.C. and De Kock, M., 2017. What is the normal value of the neutrophil-to-lymphocyte ratio?. *BMC research notes*, 10(1), pp.1-4.
- Grimaldi, D., Louis, S., Pène, F., Sirgo, G., Rousseau, C., Claessens, Y.E., Vimeux, L., Cariou, A., Mira, J.P., Hosmalin, A. and Chiche, J.D., 2011. Profound and persistent decrease of circulating dendritic cells is associated with ICU-acquired infection in patients with septic shock. *Intensive care medicine*, 37(9), pp.1438-1446.
- Gyawali, B., Ramakrishna, K. and Dhamoon, A.S., 2019. Sepsis: The evolution in definition, pathophysiology, and management. *SAGE open medicine*, 7, p.2050312119835043.
- He, X.L., Liao, X.L., Xie, Z.C., Han, L., Yang, X.L. and Kang, Y., 2016. Pulmonary infection is an independent risk factor for long-term mortality and quality of life for sepsis patients. *BioMed research international*, 2016.
- Hotchkiss, R.S., Monneret, G. and Payen, D., 2013. Sepsis-induced immunosuppression: from cellular dysfunctions to immunotherapy. *Nature Reviews Immunology*, 13(12), pp.862-874.
- Hotchkiss, R.S., Osmon, S.B., Chang, K.C., Wagner, T.H., Coopersmith, C.M. and Karl, I.E., 2005. Accelerated lymphocyte death in sepsis occurs by both the death receptor and mitochondrial pathways. *The Journal of Immunology*, 174(8), pp.5110-5118.

- Hutchins, N.A., Unsinger, J., Hotchkiss, R.S. and Ayala, A., 2014. The new normal: immunomodulatory agents against sepsis immune suppression. *Trends in molecular medicine*, 20(4), pp.224-233.
- Jeon, K., Na, S.J., Oh, D.K., Park, S., Choi, E.Y., Kim, S.C., Seong, G.M., Heo, J., Chang, Y., Kwack, W.G. and Kang, B.J., 2019. Characteristics, management and clinical outcomes of patients with sepsis: a multicenter cohort study in Korea. *Acute and critical care*, 34(3), p.179.
- Jiang, J., Du, H., Su, Y., Li, X., Zhang, J., Chen, M., Ren, G., He, F. and Niu, B., 2019. Nonviral infection-related lymphocytopenia for the prediction of adult sepsis and its persistence indicates a higher mortality. *Medicine*, 98(29).
- Kalani, C., Venigalla, T., Bailey, J., Udeani, G. and Surani, S., 2020. Sepsis patients in critical care units with obesity: is obesity protective?. *Cureus*, 12(2).
- Kalil, A.C., Metersky, M.L., Klompas, M., Muscedere, J., Sweeney, D.A., Palmer, L.B., Napolitano, L.M., O'Grady, N.P., Bartlett, J.G., Carratalà, J. and El Solh, A.A., 2016. Executive summary: management of adults with hospital-acquired and ventilator-associated pneumonia: 2016 clinical practice guidelines by the Infectious Diseases Society of America and the American Thoracic Society. *Clinical Infectious Diseases*, 63(5), pp.575-582.
- Kartika, D. S., Kumala, S., Utami, H. and Subhan, A., 2020. Analisis faktor risiko yang mempengaruhi outcome pasien sepsis di Rumah Sakit Umum Pusat Fatmawati. *SCIENTIA: Jurnal Farmasi dan Kesehatan*, 10(1), pp.17-32..
- Kasten, K.R., Muenzer, J.T. and Caldwell, C.C., 2010. Neutrophils are significant producers of IL-10 during sepsis. *Biochemical and biophysical research communications*, 393(1), pp.28-31.
- Kaushik, R., Gupta, M., Sharma, M., Jash, D., Jain, N., Sinha, N., Chaudhry, A. and Chaudhry, D., 2018. Diagnostic and prognostic role of neutrophil-to-lymphocyte ratio in early and late phase of sepsis. *Indian journal of critical care medicine: peer-reviewed, official publication of Indian Society of Critical Care Medicine*, 22(9), p.660.
- Khan, M.S., Gutch, M., Kumar, S. and Kumar, M., 2020. Insulin resistance as a prognostic indicator in severe sepsis, septic shock and multiorgan dysfunction syndrome. *International Journal of Medicine and Public Health*, 10(1).
- Kruger, P., Saffarzadeh, M., Weber, A.N., Rieber, N., Radsak, M., von Bernuth, H., Benarafa, C., Roos, D., Skokowa, J., and Hartl, D., 2015. Neutrophils: between host defence, immune modulation, and tissue injury. *PLoS pathogens*, 11(3), e1004651.
- Kurniawan, J., 2020. Persistent lymphopenia as a predictor of in-hospitality in septic patients at dr. sardjito hospital. *Acta Interna: The Journal of Internal Medicine*, 10(1), pp.37-49.

- Lambden, S., Laterre, P.F., Levy, M.M. and Francois, B., 2019. The SOFA score—development, utility and challenges of accurate assessment in clinical trials. *Critical Care*, 23(1), pp.1-9.
- Li, M., Li, Z., Yao, Y., Jin, W.N., Wood, K., Liu, Q., Shi, F.D. and Hao, J., 2017. Astrocyte-derived interleukin-15 exacerbates ischemic brain injury via propagation of cellular immunity. *Proceedings of the National Academy of Sciences*, 114(3), pp.E396-E405.
- Li, S., Hu, X., Xu, J., Huang, F., Guo, Z., Tong, L., Lui, K.Y., Cao, L., Zhu, Y., Yao, J. and Lin, X., 2019. Increased body mass index linked to greater short- and long-term survival in sepsis patients: A retrospective analysis of a large clinical database. *International Journal of Infectious Diseases*, 87, pp.109-116.
- Liu, S., Li, Y., She, F., Zhao, X. and Yao, Y., 2021. Predictive value of immune cell counts and neutrophil-to-lymphocyte ratio for 28-day mortality in patients with sepsis caused by intra-abdominal infection. *Burns & trauma*, 9.
- Liu, S., Wang, X., She, F., Zhang, W., Liu, H. and Zhao, X., 2021. Effects of neutrophil-to-lymphocyte ratio combined with interleukin-6 in predicting 28-day mortality in patients with sepsis. *Frontiers in immunology*, 12, p.757.
- Liu, Q., Jin, W.N., Liu, Y., Shi, K., Sun, H., Zhang, F., Zhang, C., Gonzales, R.J., Sheth, K.N., La Cava, A. and Shi, F.D., 2017. Brain ischemia suppresses immunity in the periphery and brain via different neurogenic innervations. *Immunity*, 46(3), pp.474-487.
- Liu, X., Shen, Y., Wang, H., Ge, Q., Fei, A. and Pan, S., 2016. Prognostic significance of neutrophil-to-lymphocyte ratio in patients with sepsis: a prospective observational study. *Mediators of inflammation*, 2016.
- Ljungström, L., Andersson, R. and Jacobsson, G., 2019. Incidences of community onset severe sepsis, Sepsis-3 sepsis, and bacteremia in Sweden—A prospective population-based study. *PloS one*, 14(12), p.e0225700.
- Llitjos, J.F., Gassama, A., Charpentier, J., Lambert, J., de Roquetaillade, C., Cariou, A., Chiche, J.D., Mira, J.P., Jamme, M. and Pène, F., 2019. Pulmonary infections prime the development of subsequent ICU-acquired pneumonia in septic shock. *Annals of intensive care*, 9(1), pp.1-9.
- Martin, M.D., Badovinac, V.P. and Griffith, T.S., 2020. CD4 T cell responses and the sepsis-induced immunoparalysis state. *Frontiers in immunology*, 11, p.1364.
- Martín, S., Pérez, A. and Aldecoa, C., 2017. Sepsis and immunosenescence in the elderly patient: a review. *Frontiers in medicine*, 4, p.20.

- May, A.K., Dossett, L.A., Norris, P.R., Hansen, E.N., Dorsett, R.C., Popovsky, K.A. and Sawyer, R.G., 2008. Estradiol is associated with mortality in critically ill trauma and surgical patients. *Critical care medicine*, 36(1), p.62.
- Meisel, C., Schwab, J.M., Prass, K., Meisel, A. and Dirnagl, U., 2005. Central nervous system injury-induced immune deficiency syndrome. *Nature Reviews Neuroscience*, 6(10), pp.775-786.
- Minasyan, H., 2019. Sepsis: mechanisms of bacterial injury to the patient. *Scandinavian journal of trauma, resuscitation and emergency medicine*, 27(1), pp.1-22.
- Nasir, N., Jamil, B., Siddiqui, S., Talat, N., Khan, F.A. and Hussain, R., 2015. Mortality in sepsis and its relationship with gender. *Pakistan journal of medical sciences*, 31(5), p.1201.
- Ni, J., Wang, H., Li, Y., Shu, Y. and Liu, Y., 2019. Neutrophil to lymphocyte ratio (NLR) as a prognostic marker for in-hospital mortality of patients with sepsis: A secondary analysis based on a single-center, retrospective, cohort study. *Medicine*, 98(46).
- Oami, T., Karasawa, S., Shimada, T., Nakada, T.A., Abe, T., Ogura, H., Shiraishi, A., Kushimoto, S., Saitoh, D., Fujishima, S. and Mayumi, T., 2021. Association between low body mass index and increased 28-day mortality of severe sepsis in Japanese cohorts. *Scientific reports*, 11(1), pp.1-7.
- Otto, G.P., Sossdorf, M., Claus, R.A., Rödel, J., Menge, K., Reinhart, K., Bauer, M. and Riedemann, N.C., 2011. The late phase of sepsis is characterized by an increased microbiological burden and death rate. *Critical care*, 15(4), pp.1-8.
- Pietropaoli, A.P., Glance, L.G., Oakes, D. and Fisher, S.G., 2010. Gender differences in mortality in patients with severe sepsis or septic shock. *Gender medicine*, 7(5), pp.422-437.
- Purnamasari, D., 2019. The emergence of non-communicable disease in Indonesia. *Acta Medica Indonesiana*, 50(4), p.273.
- Resende, C.B., Borges, I., Gonçalves, W.A., Carneiro, R., Rezende, B.M., Pinho, V., Nobre, V. and Teixeira, M.M., 2020. Neutrophil activity in sepsis: a systematic review. *Brazilian Journal of Medical and Biological Research*, 53.
- Rhee, C., Jones, T.M., Hamad, Y., Pande, A., Varon, J., O'Brien, C., Anderson, D.J., Warren, D.K., Dantes, R.B., Epstein, L. and Klompas, M., 2019. Prevalence, underlying causes, and preventability of sepsis-associated mortality in US acute care hospitals. *JAMA network open*, 2(2), pp.e187571-e187571.
- Roquilly, A., McWilliam, H.E., Jacqueline, C., Tian, Z., Cinotti, R., Rimbart, M., Wakim, L., Caminschi, I., Lahoud, M.H., Belz, G.T. and Kallies, A., 2017.



Local modulation of antigen-presenting cell development after resolution of pneumonia induces long-term susceptibility to secondary infections. *Immunity*, 47(1), pp.135-147.

Rudd, K.E., Johnson, S.C., Agesa, K.M., Shackelford, K.A., Tsoi, D., Kievlan, D.R., Colombara, D.V., Ikuta, K.S., Kissoon, N., Finfer, S. and Fleischmann-Struzek, C., 2020. Global, regional, and national sepsis incidence and mortality, 1990–2017: analysis for the Global Burden of Disease Study. *The Lancet*, 395(10219), pp.200-211.

Sari, R., KARAKURT, Z., Ay, M., Çelik, M.E., Tekan, Ü.Y., Çiyiltepe, F., Kargin, F., SALTÜRK, C., MOÇİN, Ö.Y., Güngör, G. and ADIGÜZEL, N., 2019. Neutrophil to lymphocyte ratio as a predictor of treatment response and mortality in septic shock patients in the intensive care unit. *Turkish journal of medical sciences*, 49(5), pp.1336-1349.

Shen, X.F., Cao, K., Jiang, J.P., Guan, W.X. and Du, J.F., 2017. Neutrophil dysregulation during sepsis: an overview and update. *Journal of cellular and molecular medicine*, 21(9), pp.1687-1697.

Shi, K., Wood, K., Shi, F.D., Wang, X. and Liu, Q., 2018. Stroke-induced immunosuppression and poststroke infection. *Stroke and vascular neurology*, 3(1).

Singer, M., Deutschman, C.S., Seymour, C.W., Shankar-Hari, M., Annane, D., Bauer, M., Bellomo, R., Bernard, G.R., Chiche, J.D., Coopersmith, C.M. and Hotchkiss, R.S., 2016. The third international consensus definitions for sepsis and septic shock (Sepsis-3). *Jama*, 315(8), pp.801-810.

Sônego, F., Ferreira, R.G., Kanashiro, A., Leite, C.A.V.G., Nascimento, D.C., Colón, D.F., Borges, V.D.F., Alves-Filho, J.C. and Cunha, F.Q., 2016. Paradoxical roles of the neutrophil in sepsis: protective and deleterious. *Frontiers in immunology*, 7, p.155.

Starr, M.E. and Saito, H., 2014. Sepsis in old age: review of human and animal studies. *Aging and disease*, 5(2), p.126.

Szabo, C., Thiemermann, C., Wu, C.C., Perretti, M. and Vane, J.R., 1994. Attenuation of the induction of nitric oxide synthase by endogenous glucocorticoids accounts for endotoxin tolerance in vivo. *Proceedings of the National Academy of Sciences*, 91(1), pp.271-275.

Taylor, B.C., Wilt, T.J. and Welch, H.G., 2011. Impact of diastolic and systolic blood pressure on mortality: implications for the definition of “normal”. *Journal of general internal medicine*, 26(7), pp.685-690.

Vivianni, A., and Farhanah, N., 2016. Faktor–faktor prediktor mortalitas sepsis dan syok sepsis di ICU RSUP Dr Kariadi. *Jurnal Kedokteran Diponegoro*, 5(4), pp. 504-517.

- Wacharasint, P., Boyd, J.H., Russell, J.A. and Walley, K.R., 2013. One size does not fit all in severe infection: obesity alters outcome, susceptibility, treatment, and inflammatory response. *Critical care*, 17(3), pp.1-10.
- Wang, H.E., Griffin, R., Judd, S., Shapiro, N.I. and Safford, M.M., 2013. Obesity and risk of sepsis: A population-based cohort study. *Obesity*, 21(12), pp.E762-E769.
- Wang, H.E., Jones, A.R. and Donnelly, J.P., 2017. Revised national estimates of emergency department visits for sepsis in the United States. *Critical care medicine*, 45(9), p.1443.
- Wang, H.E., Shapiro, N.I., Griffin, R., Safford, M.M., Judd, S. and Howard, G., 2013. Inflammatory and endothelial activation biomarkers and risk of sepsis: a nested case-control study. *Journal of critical care*, 28(5), pp.549-555.
- Wang, J., 2018. Neutrophils in tissue injury and repair. *Cell and tissue research*, 371(3), pp.531-539.
- Wang, S., Liu, X., Chen, Q., Liu, C., Huang, C. and Fang, X., 2017. The role of increased body mass index in outcomes of sepsis: a systematic review and meta-analysis. *BMC anesthesiology*, 17(1), pp.1-11.
- Wesche, D.E., Lomas-Neira, J.L., Perl, M., Chung, C.S. and Ayala, A., 2005. Leukocyte apoptosis and its significance in sepsis and shock. *Journal of leukocyte biology*, 78(2), pp.325-337.
- Whiles, B., Deis, A., Miller, P. and Simpson, S., 2016. Comorbid conditions predict outcomes in patients with severe sepsis. *Chest*, 149(4), p.A170.
- Wilgus, T.A., Roy, S., and McDaniel, J.C., 2013. Neutrophils and wound repair: positive actions and negative reactions. *Advances in wound care*, 2(7), pp.379-380.
- Ye, W., Chen, X., Huang, Y., Li, Y., Xu, Y., Liang, Z., Wu, D., Liu, X. and Li, Y., 2020. The association between neutrophil-to-lymphocyte count ratio and mortality in septic patients: a retrospective analysis of the MIMIC-III database. *Journal of Thoracic Disease*, 12(5), p.1843.
- Yilmaz, H., Cakmak, M., Inan, O., Darcin, T. and Akcay, A., 2015. Can neutrophil-lymphocyte ratio be independent risk factor for predicting acute kidney injury in patients with severe sepsis?. *Renal failure*, 37(2), pp.225-229.
- Yousef, A.A.A.M., Amr, Y.M. and Suliman, G.A., 2010. The diagnostic value of serum leptin monitoring and its correlation with tumor necrosis factor- $\alpha$  in critically ill patients: a prospective observational study. *Critical Care*, 14(2), pp.1-9.

Zhang, J., Shi, K., Li, Z., Li, M., Han, Y., Wang, L., Zhang, Z., Yu, C., Zhang, F., Song, L. and Dong, J.F., 2018. Organ-and cell-specific immune responses are associated with the outcomes of intracerebral hemorrhage. *The FASEB Journal*, 32(1), pp.220-229.

Zhou, Q., Wang, M., Li, S., Zhang, J., Ma, Q., Ding, Y., Ge, H., Shen, N., Zheng, Y. and Sun, Y., 2018. Impact of body mass index on survival of medical patients with sepsis: a prospective cohort study in a university hospital in China. *BMJ open*, 8(9), p.e021979.