

- Abbas, A., Lichtman, A. and Pillai, S. (2014) *Cellular and Molecular Immunology*. Philadelphia: Saunderson Elsevier.
- Alqahtani, A., Alamer, E., Mir, M., Alasmari, A., Alshahrani, M.M., Asiri, M., *et al.* (2022) 'Bacterial Coinfections Increase Mortality of Severely Ill COVID-19 Patients in Saudi Arabia', *International Journal of Environmental Research and Public Health*, 19(4). doi:10.3390/ijerph19042424.
- Anesi, G.L. (2021) *COVID-19: Epidemiology, clinical features, and prognosis of the critically ill adult*, *Up*.
- Baihaqi, F.A. and Rumaropen, H. (2022) 'Faktor-Faktor yang Berhubungan dengan Lama Rawat Inap Pasien COVID-19 di RSUD Serui Provinsi Papua: Studi Potong Lintang', *Jurnal Penyakit Dalam Indonesia*, 8(4), p. 187. doi:10.7454/jpdi.v8i4.627.
- Bhatt, K., Agolli, A., H. Patel, M., Garimella, R., Devi, M., Garcia, E., *et al.* (2021) 'High mortality co-infections of COVID-19 patients: mucormycosis and other fungal infections', *Discoveries*, 9(1), p. e126. doi:10.15190/d.2021.5.
- Castañeda-Sánchez, J.I., Duarte, A.R.M., Domínguez-López, M.L., Cruz-López, J.J. de la and Luna-Herrera, J. (2017) 'B Lymphocyte as a Target of Bacterial Infections', *Lymphocyte Updates - Cancer, Autoimmunity and Infection* [Preprint]. doi:10.5772/intechopen.69346.
- Chan, L., Karimi, N., Morovati, S., Alizadeh, Kasra, Kakish, J.E., Vanderkamp, S., *et al.* (2021) 'The roles of neutrophils in cytokine storms', *Viruses*, 13(11), pp. 1–20. doi:10.3390/v13112318.
- Citu, C., Gorun, F., Motoc, A., Sas, I., Gorun, O.M., Burlea, B., *et al.* (2022) 'The Predictive Role of NLR, d-NLR, MLR, and SIRI in COVID-19 Mortality', *Diagnostics*, 12(1), pp. 2–11. doi:10.3390/diagnostics12010122.
- Contou, D., Claudinon, A., Pajot, O., Micaëlo, M., Longuet Flandre, P., Dubert, M., *et al.* (2020) 'Bacterial and viral co-infections in patients with severe SARS-CoV-2 pneumonia admitted to a French ICU', *Annals of Intensive Care*, 10(1). doi:10.1186/s13613-020-00736-x.
- Corona, G., Pizzocaro, A., Vena, W., Rastrelli, G., Semeraro, F., Isidori, A.M., *et al.* (2021) 'Diabetes is most important cause for mortality in COVID-19 hospitalized patients: Systematic review and meta-analysis', *Reviews in Endocrine and Metabolic Disorders*, 22(2), pp. 275–296. doi:10.1007/s11154-021-09630-8.
- Drăgoescu, A.N., Pădureanu, V., Stănculescu, A.D., Chiuțu, L.C., Tomescu, P., Geormăneanu, C., *et al.* (2022) 'Neutrophil to Lymphocyte Ratio (NLR)—A Useful Tool for the Prognosis of Sepsis in the ICU', *Biomedicines*, 10(1). doi:10.3390/biomedicines10010075.
- Gelzo, M., Cacciapuoti, S., Pinchera, B., De Rosa, A., Cerneria, G., Scialò, F., *et al.* (2021) 'Prognostic Role of Neutrophil to Lymphocyte Ratio in COVID-19 Patients: Still Valid in Patients That Had Started Therapy?', *Frontiers in Public Health*, 9(June), pp. 1–5. doi:10.3389/fpubh.2021.664108.
- Gens, R., Ourtani, A., Vos, A. De and Keyser, J. De (2021) 'Usefulness of the Neutrophil-to-Lymphocyte Ratio as a Predictor of Pneumonia and Urinary Tract Infection Within the First Week After Acute Ischemic Stroke', *Frontiers in Neurology*, 12(May), pp. 1–8. doi:10.3389/fneur.2021.671739.



- Gu, X., Sha, L., Zhang, S., Shen, D., Zhao, W. and Yi, Y. (2021) 'Neutrophils and Lymphocytes Can Help Distinguish Asymptomatic COVID-19 From Moderate COVID-19', *Frontiers in Cellular and Infection Microbiology*, 11(October), pp. 1–13. doi:10.3389/fcimb.2021.654272.
- Hashemi Moghanjoughi, P., Neshat, S., Rezaei, A. and Heshmat-Ghahdarjani, K. (2022) 'Is the Neutrophil-to-Lymphocyte Ratio an Exceptional Indicator for Metabolic Syndrome Disease and Outcomes?', *Endocrine Practice*, 28(3), pp. 1–13. doi:10.1016/j.eprac.2021.11.083.
- Huang, W.J., Huang, G.T., Zhan, Q.M., Chen, J.L., Luo, W.T., Wu, L.H., *et al.* (2020) 'The neutrophil to lymphocyte ratio as a novel predictor of asthma and its exacerbation: A systematic review and meta-analysis', *European Review for Medical and Pharmacological Sciences*, 24(22), pp. 11719–11728. doi:10.26355/eurrev_202011_23824.
- Hwang, S.Y., Shin, T.G., Jo, I.J., Jeon, K., Suh, G.Y., Lee, T.R., *et al.* (2017) 'Neutrophil-to-lymphocyte ratio as a prognostic marker in critically-ill septic patients', *American Journal of Emergency Medicine*, 35(2), pp. 234–239. doi:10.1016/j.ajem.2016.10.055.
- de Jager, C.P.C., Wever, P.C., Gemen, E.F.A., Kusters, R., van Gageldonk-Lafeber, A.B., van der Poll, T. and Laheij, R.J.F. (2012) 'The Neutrophil-Lymphocyte Count Ratio in Patients with Community-Acquired Pneumonia', *PLoS ONE*, 7(10), pp. 4–11. doi:10.1371/journal.pone.0046561.
- Jin, J.M., Bai, P., He, W., Wu, F., Liu, X.F., Han, D.M., *et al.* (2020) 'Gender Differences in Patients With COVID-19: Focus on Severity and Mortality', *Frontiers in Public Health*, 8(April), pp. 1–6. doi:10.3389/fpubh.2020.00152.
- Jose, R.J. and Manuel, A. (2020) 'COVID-19 cytokine storm: the interplay between inflammation and coagulation', *The Lancet Respiratory Medicine*, 8(6), pp. e46–e47. doi:10.1016/S2213-2600(20)30216-2.
- Kesehatan, K. (2020) 'Pedoman Pencegahan dan Pengendalian Corona Virus deases (Covid-19)', *Kementrian Kesehatan*, 5, p. 178. Available at: https://covid19.go.id/storage/app/media/Protokol/REV-05_Pedoman_P2_COVID-19_13_Juli_2020.pdf.
- Kobayashi, S.D., Malachowa, N. and DeLeo, F.R. (2018) 'Neutrophils and Bacterial Immune Evasion', *Journal of Innate Immunity*, 10(5–6), pp. 432–441. doi:10.1159/000487756.
- Kwok, W.C., Tam, A.R., Ho, J.C.M., Lam, D.C.L., Tam, T.C.C., Chan, K.P.F., *et al.* (2022) 'Asthma, from mild to severe, is an independent prognostic factor for mild to severe Coronavirus disease 2019 (COVID-19)', *Clinical Respiratory Journal*, (November 2021), pp. 293–300. doi:10.1111/crj.13480.
- Lagunas-Rangel, F.A. (2020) 'PX Neutrophil-to-lymphocyte ratio and lymphocyte-to-C-reactive protein ratio in patients with severe coronavirus disease 2019 (COVID-19) A meta-analysis.pdf'.
- Lee, J.S., Kim, N.Y., Na, S.H., Youn, Y.H. and Shin, C.S. (2018) 'Reference values of neutrophil-lymphocyte ratio, lymphocyte-monocyte ratio, platelet-lymphocyte ratio, and mean platelet volume in healthy adults in South Korea', *Medicine (United States)*, 97(26), pp. 1–5. doi:10.1097/MD.00000000000011138.
- Li, J., Liu, Z., Wu, G., Yi, M., Chen, Y., Li, K., *et al.* (2020) 'D-Dimer as a Prognostic Indicator in Critically Ill Patients Hospitalized With COVID-19 in Leishenshan Hospital, Wuhan, China', *Frontiers in Pharmacology*, 11(December), pp. 1–9. doi:10.3389/fphar.2020.600592.



Liu, F., Li, L., Xu, M., Wu, J., Luo, D., Zhu, Y., *et al.* (2020a) ‘Prognostic value of IL-6, CRP, and PCT in patients with COVID-19’, *J Clin Virology* [Preprint], (January).

Liu, F., Li, L., Xu, M., Wu, J., Luo, D., Zhu, Y., *et al.* (2020b) ‘Prognostic value of interleukin-6, C-reactive protein, and procalcitonin in patients with COVID-19’, *Journal of Clinical Virology*, 127(January), p. 104370.

Liu, T., Zhang, Jieying, Yang, Y., Ma, H., Li, Z., Zhang, Jiaoyue, *et al.* (2020) ‘The role of interleukin-6 in monitoring severe case of coronavirus disease 2019’, *EMBO Molecular Medicine*, 12(7), pp. 1–12. doi:10.15252/emmm.202012421.

Liu, Y., Du, X., Chen, J., Jin, Y., Peng, L., Wang, H.H.X., *et al.* (2020a) ‘Neutrophil-to-lymphocyte ratio as an independent risk factor for mortality in hospitalized patients with COVID-19’, *Journal of Infection*, 81(1), pp. e6–e12. doi:10.1016/j.jinf.2020.04.002.

Liu, Y., Du, X., Chen, J., Jin, Y., Peng, L., Wang, H.H.X., *et al.* (2020b) ‘Neutrophil-to-lymphocyte ratio as an independent risk factor for mortality in hospitalized patients with COVID-19’, *Journal of Infection*, 81(1), pp. e6–e12. doi:10.1016/j.jinf.2020.04.002.

Ma, Y., Zhang, Y. and Zhu, L. (2021) ‘Role of neutrophils in acute viral infection’, *Immunity, Inflammation and Disease*, 9(4), pp. 1186–1196. doi:10.1002/iid3.500.

Mason, R.J. (2020) ‘Pathogenesis of COVID-19 from a cell biology perspective’, *European Respiratory Journal*, 55(4), pp. 9–11. doi:10.1183/13993003.00607-2020.

Mojtabavi, H., Saghadzadeh, A. and Rezaei, N. (2020) ‘Interleukin-6 and severe COVID-19: a systematic review and meta-analysis’, *European Cytokine Network*, 31(2), pp. 44–49. doi:10.1684/ecn.2020.0448.

Nalbant, A., Kaya, T., Varim, C., Yaylaci, S., Tamer, A. and Cinemre, H. (2020) ‘Can the neutrophil/lymphocyte ratio (NLR) have a role in the diagnosis of coronavirus 2019 disease (COVID-19)?’, *Revista da Associacao Medica Brasileira*, 66(6), pp. 746–751. doi:10.1590/1806-9282.66.6.746.

Ni, J., Wang, H., Li, Y., Shu, Y. and Liu, Y. (2019) ‘Neutrophil to lymphocyte ratio (NLR) as a prognostic marker for in-hospital mortality of patients with sepsis’, *Medicine (United States)*, 98(46), pp. 9–13. doi:10.5061/dryad.k1jj600.a.

Ozma, M.A., Maroufi, P., Khodadadi, E., Köse, Ş., Esposito, I., Ganbarov, K., *et al.* (2020) ‘Clinical manifestation, diagnosis, prevention and control of SARS-CoV-2 (COVID-19) during the outbreak period’, *Infezioni in Medicina*, 28(2), pp. 153–165.

De Pablo, R., Monserrat, J., Prieto, A. and Alvarez-Mon, M. (2014) ‘Role of Circulating Lymphocytes in Patients with Sepsis’, *BioMed Research International*, 2014(671087), pp. 1–11. doi:10.1155/2014/671087.

Peñaloza, H.F., Lee, J.S. and Ray, P. (2021) ‘Neutrophils and lymphopenia, an unknown axis in severe COVID-19 disease’, *PLoS Pathogens*, 17(9), pp. 1–9. doi:10.1371/journal.ppat.1009850.

Ponti, G., Maccaferri, M., Ruini, C., Tomasi, A. and Ozben, T. (2020) ‘Biomarkers associated with COVID-19 disease progression’, *Critical Reviews in Clinical Laboratory Sciences*, 0(0), pp. 1–11. doi:10.1080/10408363.2020.1770685.

Rezaei, M., Marjani, M., Mahmoudi, S., Mortaz, E. and Mansouri, D. (2021) ‘Dynamic Changes of Lymphocyte Subsets in the Course of COVID-19’, *International Archives of Allergy and Immunology*, 182(3), pp. 254–262. doi:10.1159/000514202.



Rosales, C. (2018) 'Neutrophil: A cell with many roles in inflammation or several cell types?', *Frontiers in Physiology*, 9(FEB), pp. 1–17. doi:10.3389/fphys.2018.00113.

Sevencan, N.O. and Ozkan, A.E. (2019) 'Associations between neutrophil/lymphocyte ratio, platelet/lymphocyte ratio, albuminuria and uric acid and the estimated glomerular filtration rate in hypertensive patients with chronic kidney disease stages 1–3', *Archives of Medical Science*, 15(5), pp. 1232–1239. doi:10.5114/aoms.2018.76262.

Shi, N., Ma, Y., Fan, Y., Wang, J., Zhao, C., Li, G., *et al.* (2020) 'Predictive Value of the Neutrophil-to-Lymphocyte Ratio(NLR) for Diagnosis and Worse Clinical Course of the COVID-19: Findings from Ten Provinces in China', *SSRN Electronic Journal* [Preprint]. doi:10.2139/ssrn.3569838.

Sônego, F., Castanheira, F.V. e. S., Ferreira, R.G., Kanashiro, A., Leite, C.A.V.G., Nascimento, D.C., *et al.* (2016) 'Paradoxical roles of the neutrophil in sepsis: Protective and deleterious', *Frontiers in Immunology*, 7(APR), pp. 1–7. doi:10.3389/fimmu.2016.00155.

Song, X., Ji, J., Reva, B., Joshi, H., Calinawan, A.P., Mazumdar, M., *et al.* (2021) 'Post-anticoagulant d-dimer is a highly prognostic biomarker of covid-19 mortality', *ERJ Open Research*, 7(3). doi:10.1183/23120541.00018-2021.

Susilo, A., Rumende, C.M., Pitoyo, C.W., Santoso, W.D., Yulianti, M., Herikurniawan, H., *et al.* (2020) 'Coronavirus Disease 2019: Tinjauan Literatur Terkini', *Jurnal Penyakit Dalam Indonesia*, 7(1), p. 45. doi:10.7454/jpdi.v7i1.415.

Tavakolpour, S., Rakhshandehroo, T., Wei, E.X. and Rashidian, M. (2020) 'Lymphopenia during the COVID-19 infection: What it shows and what can be learned', *Immunology Letter*, (January), pp. 19–21.

Taylor, R.S.S.C.M.B.N. (2021) 'Histology, T-Cell Lymphocyte', in *StatPearls*. Treasure Island (FL): StatPearls Publishing, pp. 1–5.

Ulloque-Badaracco, J.R., Ivan Salas-Tello, W., Al-kassab-Córdova, A., Alarcón-Braga, E.A., Benites-Zapata, V.A., Maguiña, J.L. and Hernandez, A. V. (2021) 'Prognostic value of neutrophil-to-lymphocyte ratio in COVID-19 patients: A systematic review and meta-analysis', *International Journal of Clinical Practice*, 75(11), pp. 1–16. doi:10.1111/ijcp.14596.

Vafadar, E., Teimouri, A., Rezaee, R., Morovatdar, N. and Foroughian, M. (2020) 'Increased age, neutrophil-to-lymphocyte ratio (NLR) and white blood cells count are associated with higher COVID-19 mortality', *American Journal of Emergency Medicine*, 40(January), pp. 11–14. doi:<https://doi.org/10.1016/j.ajem.2020.12.003>.

Wagner, C., Kotsougiani, D., Pioch, M., Prior, B., Wentzensen, A. and Hänsch, G.M. (2008) 'T lymphocytes in acute bacterial infection: Increased prevalence of CD11b+ cells in the peripheral blood and recruitment to the infected site', *Immunology*, 125(4), pp. 503–509. doi:10.1111/j.1365-2567.2008.02863.x.

Wan, Y., Shang, J., Graham, R., Baric, R.S. and Li, F. (2020) 'Reconocimiento de receptores por el nuevo coronavirus de Wuhan: un análisis basado en estudios estructurales de una década del coronavirus del SARS', *Journal of Virology*, 94(7), pp. 1–9.

Wardika, I.K. and Sikesa, I.G.P.H. (2021) 'Pengukuran Interleukin-6 (IL-6), C-Reactive Protein (CRP) dan D-Dimer sebagai prediktor prognosis pada pasien COVID-19 gejala berat: sebuah tinjauan pustaka', *Intisari Sains Medis*, 12(3), p. 901. doi:10.15562/ism.v12i3.1158.

WHO (2022) *COVID-19 weekly epidemiological update*, World Health Organization.



Wu, Z. hong, Tang, Y. and Cheng, Q. (2021) 'Diabetes increases the mortality of patients with COVID-19: a meta-analysis', *Acta Diabetologica*, 58(2), pp. 139–144. doi:10.1007/s00592-020-01546-0.

Xiang, P., Liu, J., Liu, Y., Pu, L., Xiong, H., Li, C., *et al.* (2020) 'Neutrophil-to-lymphocyte ratio predicts critical illness patients with 2019 coronavirus disease in the early stage', *Journal of Translational Medicine*, 18(1), pp. 1–12. doi:10.1186/s12967-020-02374-0.

Yan, X., Li, F., Wang, X., Yan, J., Zhu, F., Tang, S., *et al.* (2020) 'Neutrophil to lymphocyte ratio as prognostic and predictive factor in patients with coronavirus disease 2019: A retrospective cross - sectional study', (April). doi:10.1002/jmv.26061.

Yang, A., Liu, J., Tao, W. and Li, H. (2020) 'The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients', *ELSEVIER* [Preprint]. doi:<https://doi.org/10.1016/j.intimp.2020.106504>.

Zhou, F., Yu, T., Du, R., Fan, G., Liu, Y., Liu, Z., *et al.* (2020) 'Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study', *The Lancet*, 395(10229), pp. 1054–1062. doi:10.1016/S0140-6736(20)30566-3.