

Daftar Pustaka

- Arif, S.K., Rukka, A.S. and Wahyuni, S., 2017. Comparison of neutrophils-lymphocytes ratio and procalcitonin parameters in sepsis patient treated in intensive care unit Dr. Wahidin Hospital, Makassar, *Indonesia. Journal of Medical Science*, 17(1), pp.17-21
- Durlu, N., BatiiSlam, Y. and Özatamer, O., 2002. The Effects Of Isoflurane And Sevofluran On Immune System In Minor Surgical Interventions, *Journal of Ankara Medical School*, pp. 001–008.
- Firman B., 2007. Comparison of the effect of Sevoflurane and Isoflurane on the number of peripheral polimorphonuclear. Doctoral dissertation, program Pascasarjana Universitas Diponegoro.
- Frohlich D, Schwall B, Taeger K, Hobbhahn J, Rothe G, Schmitz G *et al.* Effect of Volatile Anaesthetics on Human Neutrophyl Oxydative Response to the Bacterial Peptide FMLP. *Br J Anaesth* 1997; 78: 718-23.
- Gibson PH, Croal BL, Cuthbertson BH, Small GR, Ifezulike AI, Gibson G, *et al.* Preoperative neutrophil-lymphocyte ratio and outcome from coronary artery bypass grafting. *Am Heart J* 2007;154:995-1002
- Gonda, K., Shibata, M., Sato, Y., Washio, M., Takeshita, H., *et al.* Elevated neutrophil-to-lymphocyte ratio is associated with nutritional impairment, immune suppression, resistance to S-1 plus cisplatin, and poor prognosis in patients with stage IV gastric cancer. 2017. *Molecular and Clinical Oncology*, 7(6), 1073-1078.
- Hall, J. E., 2016. Resistance of the Body to Infection: Leukocytes, Granulocytes, the Monocyte-Macrophage System, and Inflammation *Guyton and Hall textbook of medical physiology*. 13th edition. Philadelphia, PA: Elsevier
- Holub, M., Beran, O., Kaspříková, N. and Chalupa, P., 2012. Neutrophil to Lymphocyte Count Ratio as a Biomarker of Bacterial Infections, *Open Medicine*, 7(2).

Jager, C.P., van Wijk P.T., Mathoera R.B and Wever P.C., 2010. Lymphocytopenia and neutrophil-lymphocyte count ratio predict bacteremia better than conventional infection markers in an emergency care unit. *Critical care*, 14(5), p.R192

Karadasli, H., Cocelli, L.P., and Ugur, M.G., 2012. Comparison of Effects of Low Flow Sevoflurane and Desflurane Anesthesia on Neutrophil and T-Cell Populations, *Current Therapeutic Research*, 73(1–2), pp. 41–51..

Kim, W. H., Seung Jin H., Sangwook Ko, Hahm T.S. and, Lee S.M., 2011. The effect of anesthetic techniques on neutrophil-to-lymphocyte ratio After Laparoscopy-Assisted Vaginal Hysterectomy, *Acta Anaesthesiologica Taiwanica*, 49(3), pp. 83–87

Koutsogiannaki, S., Bernier, R., Tazawa, K. and Yuki, K., 2019. Volatile Anesthetic Attenuates Phagocyte Function and Worsens Bacterial Loads in Wounds, *Journal of Surgical Research*, 233, pp. 323–330.

Kurosawa, S, & Kato, M. 2008. Anesthetics, immune cells, and immune responses. *Journal of anesthesia*, 22(3), 263-277.

Leng S, Xue QL, Huang Y, Semba R, Chaves P, Bandeen-Roche K, et al. Total and differential white blood cell counts and their associations with circulating interleukin-6 levels in community-dwelling older women. *J Gerontol A Biol Sci Med Sci* 2005;60:195-9

Li, J., Chen, Q., Luo, X., Hong, J., Pan, K., et al. Neutrophil-to-lymphocyte ratio positively correlates to age in healthy population. 2015. *Journal of clinical laboratory analysis*, 29(6), 437-443.

Luhulima, D., Marwito, M. and Eva, O., 2018. Neutrophil - lymphocyte count ratio in bacterial sepsis. *Indonesia journal of clinical pathology and medical laboratory*, 23(3), pp.257-262

Matsuoka, H., Kurosawa, S., Horinouchi, T., Kato, M. and Hashimoto, Y., 2001. Inhalation anesthetics induce apoptosis in normal peripheral lymphocytes in vitro. *Anesthesiology: The Journal of the American Society of Anesthesiologists*, 95(6), pp.1467-1472.

Merriman, H. Reinhold, H. A., and West, M. P., 2014. Infectious Diseases, in *Acute Care Handbook for Physical Therapists*. Elsevier, pp. 313–334

Morgan G.E., Baldini G., Butterworth J.F., Carli, F., Frölich M.A., Giesecke M., Hosur S., Ilfeld B.M., Madison S.J., Mariano, E.R., and McGlinch B.P., 2013. Inhalational Anesthetic

Morisaki H, Aoyoma Y, Shimada M, Ochiai R, and Takeda J., 1997 Leucocyte Distribution During Sevoflurane Anaesthesia. *Br J Anaesth*. 1997;80:502–3.

Neal CP, Mann CD, Sutton CD, Garcea G, Ong SL, Steward WP, et al. Evaluation of the prognostic value of systemic inflammation and socioeconomic deprivation in patients with resectable colorectal liver metastases. *Eur J Cancer* 2009;45:56-64

Ni Eochagain, A., Burns, D., Riedel, B., Sessler, D.I. and Buggy, D.J., 2018. The effect of anaesthetic technique during primary breast cancer surgery on neutrophil–lymphocyte ratio, platelet–lymphocyte ratio and return to intended oncological therapy. *Anaesthesia*, 73(5), pp.603-611.

O’Brien J, Wang J, Redmond HP, and Shorten G., 2010 Effect of sevoflurane on human neutrophil apoptosis. *Eur J Anaesthesiol*. 2010;20:111–5.

Sastroasmoro S., 2008. Inferensi dari sampel ke populasi. Dalam *Dasar-Dasar Metodologi Penelitian Klinis*. Edisi ke-3,. Sagung Seto. Jakarta. p. 12-25

Sedghi, S., Kutscher, H. L., Davidson, B. A., and Knight, P. R. 2017. Volatile anesthetics and immunity. *Immunological investigations*, 46(8), 793-804

Stoelting, R.K. Anast, N., Hillier, S.C., Elsharkawy H., Kachulis, H., Cummings K., Flood P., Maheshwar K., Slinger P., 2015. Inhaled anesthetics.in *Pharmacology and physiology in anesthetic practice*. 5th ed. Lippincott Williams & Wilkins. p>35-72

Stollings, L.M., Li-Jie Jia, Pei Tang, Dou H., Binfeng Lu..and Yan Xu, 2016. Immune modulation by volatile anesthetics. *Anesthesiology: The Journal of the American Society of Anesthesiologists*, 125(2), pp.399-41Tyther R, Halligant M, Wang J, Redmond HP, and Shorten G., 2009, Effects of Chronic Occupational Exposure to Anaesthetics Gases on the Rate of Neutrophil Apoptosis Among Anaesthetists. *Eur J Anaesthesiol*.;19:604–8.

Surhonne, N., Hebri C., Kannan S., Duggappa D.R., Raghavendra Rao R.S, and Mapari C.G., 2019. The effect of anesthetic techniques on neutrophil to lymphocyte ratio in patients undergoing infraumbilical surgeries, *Korean Journal of Anesthesiology*, 72(5), pp. 458–465

Takahashi J, Shono Y, Hirabayashi H, Kamimura M, Nakagawa H, Ebara S, et al. Usefulness of white blood cell differential for early diagnosis of surgical wound infection following spinal instrumentation surgery. *Spine* 2006;31:1020-5

Venkatraghavan, L., Tan, T. P., Mehta, J., Arekapudi, A., Govindarajulu, A., and Siu, E., 2015. Neutrophil Lymphocyte Ratio as a predictor of systemic inflammation-A cross-sectional study in a pre-admission setting. *F1000 Research* 2015, 4:123

Walsh, S.R., Cook, E.J., Goulder, F., Justin, T.A., and Keeling, N.J., 2005. Neutrophil-lymphocyte ratio as a prognostic factor in colorectal cancer. *J Surg Oncol*, 91:181-184

Yamanaka T, Matsumoto S, Teramukai S, Ishiwata R, Nagai Y, Fukushima M. The baseline ratio of neutrophils to lymphocytes is associated with patient prognosis in advanced gastric cancer. *Oncology* 2007;73:215-20.

Yuki, K. and Eckenhooff, R.G. 2016. Mechanisms of the Immunological Effects of Volatile Anesthetics: A Review, *Anesthesia & Analgesia*, 123(2), pp. 326–335

Zahorec, R., 2001. Ratio of neutrophil to lymphocyte counts-rapid and simple parameter of systemic inflammation and stress in critically ill. *Bratislavske lekarske listy*, 102(1), pp.5-14

.

.