

## DAFTAR PUSTAKA

- Aiello, G., Barra, D., Parisi, R., Arienzo, M., Donadio, C., Ferrara, L., Toscanesi, M., & Trifuoggi, M. (2021). Infralittoral ostracoda and benthic foraminifera of the Gulf of Pozzuoli (Tyrrhenian Sea, Italy). *Aquatic Ecology*, 55(3), 955–998. <https://doi.org/10.1007/s10452-021-09874-1>
- Amalisa, Mahasri, G., & Kismiyati. (2021). The correlation between ectoparasite infestation and total *Vibrio parahaemolyticus* bacteria in Pacific white shrimp (*Litopenaeus vannamei*) in Super Intensive Ponds. *IOP Conference Series: Earth and Environmental Science*, 888(1). <https://doi.org/10.1088/1755-1315/888/1/012003>
- Arifin, N. B., Fakhri, M., Fakhri, M., Yuniarti, A., Yuniarti, A., Hariati, A. M., & Hariati, A. M. (2018). Komunitas Fitoplankton Pada Sistem Budidaya Intensif Udang Vaname, *Litopenaeus vannamei* di Probolinggo, Jawa Timur. *Jurnal Ilmiah Perikanan Dan Kelautan*, 10(1), 46. <https://doi.org/10.20473/jipk.v10i1.8542>
- Asaduzzaman, M., Rahman, M. M., Azim, M. E., Islam, M. A., Wahab, M. A., Verdegem, M. C. J., & Verreth, J. A. J. (2010). Effects of C/N ratio and substrate addition on natural food communities in freshwater prawn monoculture ponds. *Aquaculture*, 306(1–4), 127–136. <https://doi.org/10.1016/j.aquaculture.2010.05.035>
- Avnimelech, Y. 1999. Carbon/nitrogen ratio as a control element in aquaculture systems. *Aquaculture*, 176: 227–235.
- Azim, M. E., & Little, D. C. (2008). The biofloc technology (BFT) in indoor tanks: Water quality, biofloc composition, and growth and welfare of Nile tilapia (*Oreochromis niloticus*). *Aquaculture*, 283(1–4), 29–35. <https://doi.org/10.1016/j.aquaculture.2008.06.036>
- Ballester, E. L. C., Marzarotto, S. A., Silva de Castro, C., Frozza, A., Pastore, I., & Abreu, P. C. (2017). Productive performance of juvenile freshwater prawns *Macrobrachium rosenbergii* in biofloc system. *Aquaculture Research*, 48(9), 4748–4755. <https://doi.org/10.1111/are.13296>
- Battauz, Y. S., de Paggi, S. B. J., & Paggi, J. C. (2017). Macrophytes as dispersal vectors of zooplankton resting stages in a subtropical riverine floodplain. *Aquatic Ecology*, 51(2), 191–201. <https://doi.org/10.1007/s10452-016-9610-3>
- Belfiore, A. P., Buley, R. P., Fernandez-Figueroa, E. G., Gladfelter, M. F., & Wilson, A. E. (2021). Zooplankton as an alternative method for controlling phytoplankton in catfish pond aquaculture. *Aquaculture Reports*, 21. <https://doi.org/10.1016/j.aqrep.2021.100897>

- Berger, R., Henriksson, E., Kautsky, L., & Malm, T. (2003). Effects of filamentous algae and deposited matter on the survival of *Fucus vesiculosus* L. germlings in the Baltic Sea. *Aquatic Ecology*, 37, 1-11.
- Bougis, P. (1976). *Marine Plankton Ecology*. North-Holland Publishing Company. pp. 7-174.
- Boyd, C. E. (2000). *Effluent Composition & Water Quality Standards -- Implementing GAA's Responsible Aquaculture Program*.  
<https://www.researchgate.net/publication/285843894>
- Bradley, P. B., Sanderson, M. P., Frischer, M. E., Brofft, J., Booth, M. G., Kerkhof, L. J., & Bronk, D. A. (2010). Inorganic and organic nitrogen uptake by phytoplankton and heterotrophic bacteria in the stratified Mid-Atlantic Bight. *Estuarine, Coastal and Shelf Science*, 88(4), 429-441.  
<https://doi.org/10.1016/j.ecss.2010.02.001>
- Brandão, S. N., Hoppema, M., Kamenev, G. M., Karanovic, I., Riehl, T., Tanaka, H., Vital, H., Yoo, H., & Brandt, A. (2019). Review of Ostracoda (Crustacea) living below the Carbonate Compensation Depth and the deepest record of a calcified ostracod. In *Progress in Oceanography* (Vol. 178). Elsevier Ltd.  
<https://doi.org/10.1016/j.pocean.2019.102144>
- Brito, R., Chimal, M. E., Gelabert, R., Gaxiola, G., & Rosas, C. (2004). Effect of artificial and natural diets on energy allocation in *Litopenaeus setiferus* (Linnaeus, 1767) and *Litopenaeus vannamei* (Boone, 1931) early postlarvae. *Aquaculture*, 237(1-4), 517-531. <https://doi.org/10.1016/j.aquaculture.2004.05.012>
- Brito, L. O., dos Santos, I. G. S., de Abreu, J. L., de Araújo, M. T., Severi, W., & Gálvez, A. O. 2016. Effect of the addition of diatoms (*Navicula* spp.) and rotifers (*Brachionus plicatilis*) on water quality and growth of the *Litopenaeus vannamei* postlarvae reared in a biofloc system. *Aquaculture Research*, 47(12): 3990-3997.
- BSN. 2006. Bening udang vaname (*Litopenaeus vannamei*) kelas benih sebar. Diakses pada 18 Januari 2021 dari <http://kkp.go.id/an-component/media/upload-gambar-pendukung/DIT%20PERBENIHAN/SNI%20Perbenihan/>
- Cardozo, A. P., Oliveira Britto, V., & Odebrecht, C. (2011). Temporal variability of plankton and nutrients in shrimp culture ponds vs. adjacent estuarine water. In *Pan-American Journal of Aquatic Sciences* (Vol. 6, Issue 1).
- Casé, M., Leça, E. E., Leitão, S. N., SantAnna, E. E., Schwamborn, R., & de Moraes Junior, A. T. (2008). Plankton community as an indicator of water quality in tropical shrimp culture ponds. *Marine Pollution Bulletin*, 56(7), 1343-1352.  
<https://doi.org/10.1016/j.marpolbul.2008.02.008>
- Chung, M. H., & Lee, K.-S. (2008). Species Composition of the Epiphytic Diatoms on the Leaf Tissues of Three *Zostera* Species Distributed on the Southern Coast of Korea. *Algae*, 23(1), 75-81.

- Ciji, A., & Akhtar, M. S. (2020). Nitrite implications and its management strategies in aquaculture: a review. *Reviews in Aquaculture* 12(2), 878–908.  
<https://doi.org/10.1111/raq.12354>
- Dinesh Kumar, S., Santhanam, P., Krishnaveni, N., Raju, P., Begum, A., Ahmed, S. U., Perumal, P., Pragnya, M., Dhanalakshmi, B., & Kim, M. K. (2020). Baseline assessment of water quality and ecological indicators in *Penaeus vannamei* farm wastewater along the Southeast coast of India. *Marine Pollution Bulletin*, 160.  
<https://doi.org/10.1016/j.marpolbul.2020.111579>
- El-Zeiny, A. M., & Abd El-Hamid, H. T. (2022). Environmental and human risk assessment of heavy metals at northern Nile Delta region using geostatistical analyses. *Egyptian Journal of Remote Sensing and Space Science*, 25(1), 21–35.  
<https://doi.org/10.1016/j.ejrs.2021.12.005>
- Epa, U. P. K. (2017). Effect of blind feeding during the first month of the culture cycle on growth of *Penaeus monodon*, Fabricius cultured in semi-intensively managed ponds in the North West of Sri Lanka. In *Journal of the University of Kelaniya: Vol. XXXII*.
- Evans, C. D., Monteith, D. T., & Cooper, D. M. (2005). Long-term increases in surface water dissolved organic carbon: Observations, possible causes and environmental impacts. *Environmental Pollution*, 137(1), 55–71.  
<https://doi.org/10.1016/j.envpol.2004.12.031>
- Falkowski, P. (2012). The Power of Phytoplankton. *Nature*.
- Fielder, D. S., Purser, G. J., & Battaglene, S. C. (2000). Effect of rapid changes in temperature and salinity on availability of the rotifers *Brachionus rotundiformis* and *Brachionus plicatilis*. *Aquaculture*, 189. [www.elsevier.nl/locate/aqua-online](http://www.elsevier.nl/locate/aqua-online)
- Firdaus, & Wijayanti. (2019). *Fitoplankton dan Siklus Karbon Global*. *Oseana* 44, 35–48.
- Gao, Y., Wei, J., Yuan, J., Zhang, X., Li, F., & Xiang, J. (2017). Transcriptome analysis on the exoskeleton formation in early developmental stages and reconstruction scenario in growth-moulting in *Litopenaeus vannamei*. *Scientific Reports*, 7(1). <https://doi.org/10.1038/s41598-017-01220-6>
- Ghufron, M., Lamid, M., Desi, P., Sari, W., & Suprpto, H. (2017). Enlargement Technique of Vannamei Shrimp (*Litopenaeus vannamei*) on Mentoring Pond of PT Central Proteina Prima Tbk in Randutatah Village, Paiton, Probolinggo, East Java. *Journal of Aquaculture and Fish Health* (Vol. 7, Issue 2).
- Goldman, C. R., & Horne, A. J. (1983). *Limnology: USA*. McGraw-Hill. pp. 197-238.
- Gutiérrez, J. C. S., Ponce-Palafox, J. T., Pineda-Jaimes, N. B., Arenasfuentes, V., Arredondo-Figueroa, J. L., & Cifuentes-Lemus, J. L. (2016). The feeding ecology of penaeid shrimp in tropical lagoon-estuarine systems. *Gayana*, 80(1), 16–28.  
<https://doi.org/10.4067/S0717-65382016000100003>

- Hajong, P., & Ramanujam, P. (2021). Algal community structure and primary productivity of stream ecosystem of west Garo Hills, Meghalaya, India. *International Journal of Ecology and Environmental Sciences*, 3(3), 140–149. <https://www.researchgate.net/publication/354914774>
- Hu, Y., Ni, Q., Wu, Y., Zhang, Y., & Guan, C. (2011). Study on CO<sub>2</sub> removal method in recirculating aquaculture waters. *Procedia Engineering*, 15, 4780–4789. <https://doi.org/10.1016/j.proeng.2011.08.894>
- Jepsen, P. M., van Someren Gréve, H., Jørgensen, K. N., Kjær, K. G. W., & Hansen, B. W. (2021). Evaluation of high-density tank cultivation of the live-feed cyclopoid copepod *Apocyclops royi* (Lindberg 1940). *Aquaculture*, 533. <https://doi.org/10.1016/j.aquaculture.2020.736125>
- Kent, M., Browdy, C. L., & Leffler, J. W. (2011). Consumption and digestion of suspended microbes by juvenile Pacific white shrimp *Litopenaeus vannamei*. *Aquaculture*, 319(3–4), 363–368. <https://doi.org/10.1016/j.aquaculture.2011.06.048>
- Keputusan Menteri Kelautan dan Perikanan Nomor: 75/PERMEN-KP/2016 Tentang Pedoman Umum Pembesaran Udang Windu (*Penaeus Monodon*) dan Udang Vaname (*Litopenaeus Vannamei*).
- Keputusan Menteri Kelautan dan Perikanan Nomor: Kep. 28/Men/2004 Tentang Pedoman Umum Budidaya Udang Di Tambak.
- Krebs, C. J. (2014). *Ecology : The Experimental Analysis of Distribution and Abundance*. Pearson.
- Kreibich, T., Saborowski, R., Hagen, W., & Niehoff, B. (2008). Short-term variation of nutritive and metabolic parameters in *Temora longicornis* females (Crustacea, Copepoda) as a response to diet shift and starvation. *Helgoland Marine Research*, 62(3), 241–249. <https://doi.org/10.1007/s10152-008-0112-0>
- Kumar, V., al Momin, S., Kumar, V. v., Ahmed, J., Al-Musallam, L., Shajan, A. B., Al-Aqeel, H., Al-Mansour, H., & Al-Zakri, W. M. (2021). Distribution and diversity of eukaryotic microalgae in Kuwait waters assessed using 18S rRNA gene sequencing. *PLoS ONE*, 16(4 April). <https://doi.org/10.1371/journal.pone.0250645>
- Li, W., Yang, M., Wang, B., & Liu, C. Q. 2022. Regulation strategy for nutrient-dependent carbon and nitrogen stoichiometric homeostasis in freshwater phytoplankton. *Science of the Total Environment*, 823. <https://doi.org/10.1016/j.scitotenv.2022.153797>
- Llario, F., Rodilla, M., Escrivá, J., Falco, S., & Sebastiá-Frasquet, M. T. 2019. Phytoplankton evolution during the creation of a biofloc system for shrimp culture. *International Journal of Environmental Science and Technology*, 16(1): 211–222.

- Lu, Q., Liu, X., Qiu, X., Liang, T., Chen, J., Zhao, S., Ouyang, S., Jin, B., & Wu, X. (2021). Changes and drivers of zooplankton diversity patterns in the middle reach of Yangtze River floodplain lakes, China. *Ecology and Evolution*, 11(24), 17885–17900. <https://doi.org/10.1002/ece3.8353>
- Martins, T. G., Odebrecht, C., Jensen, L. v., D'Oca, M. G., & Wasielesky, W. (2016). The contribution of diatoms to bioflocs lipid content and the performance of juvenile *Litopenaeus vannamei* (Boone, 1931) in a BFT culture system. *Aquaculture Research*, 47(4), 1315–1326. <https://doi.org/10.1111/are.12592>
- Mawarni, A., Azizah, F. N. N., Sartika, H. W., Hadisusanto, S., Putri, D. M., & Reza, A. (2020). Short communication: Community of phytoplankton in peatland canal, Riau, and wet dune slacks of Parangtritis, Yogyakarta, Indonesia. *Biodiversitas*, 21(5), 1874–1879. <https://doi.org/10.13057/biodiv/d210513>
- Montagnes, D. J. S., & Franklin, D. J. (2001). Effect of temperature on diatom volume, growth rate, and carbon and nitrogen content: Reconsidering some paradigms. *Limnology and Oceanography*, 46(8).
- Nasution, A. K., Takarina, N. D., & Thoha, H. (2021). The presence and abundance of harmful dinoflagellate algae related to water quality in Jakarta bay, Indonesia. *Biodiversitas*, 22(5), 2909–2917. <https://doi.org/10.13057/biodiv/d220556>
- Neelgund, H. D., & Kadadevaru, G. G. (2021). A Study on Seasonal Variation in Zooplankton Abundance in Kadasgatti Minor Irrigation Tank of Bailhongal Taluk, Belagavi District, Karnataka State, INDIA. *Indian Journal of Science and Technology*, 14(27), 2238–2249. <https://doi.org/10.17485/IJST/v14i27.323>
- Niu, B., Zhai, M., Hernández-Becerril, D. U., & Li, Y. (2022). Diversity and phylogeny of Chaetoceros species (Bacillariophyceae) with a central valve linking protuberance. *Phycologia*, 61(1), 104–115. <https://doi.org/10.1080/00318884.2021.2007712>
- Nonwachai, T., Purivirojkul, W., Limsuwan, C., Chuchird, N., Velasco, M., & Dhar, A. K. (2010). Growth, nonspecific immune characteristics, and survival upon challenge with *Vibrio harveyi* in Pacific white shrimp (*Litopenaeus vannamei*) raised on diets containing algal meal. *Fish and Shellfish Immunology*, 29(2), 298–304. <https://doi.org/10.1016/j.fsi.2010.04.009>
- Nour, A. A., Zaki, M. A., Abdel-Rahim, M. M., & Srour, T. M. (2004). Growth Performance and Feed Utilization of Marine Shrimp *Penaeus Semisulcatus* Post-Larvae Reared in Two Nursery Systems with Different Stocking Sizes. *Egyptian Journal of Aquatic Research*, 30(B), 390–405. <https://www.researchgate.net/publication/309458547>
- Padmakumar, K. B., Thomas, L. C., Salini, T. C., Vijayan, A., & Sudhakar, M. (2018). Subsurface bloom of dinoflagellate *Gonyaulax polygramma* Stein in the shelf waters off Mangalore-South Eastern Arabian Sea. In *Indian Journal of Geo Marine Sciences*, 47(8).



- Panjaitan, A. S., Hadie, W., & Harijati, S. (2015). Penggunaan *Chaetoceros calcitrans*, *Thalassiosira weissflogii* dan Kombinasinya pada Pemeliharaan Larva Udang Vaname (*Litopenaeus vannamei*, Boone 1931). *Berita Biologi*, 14(3), 235–240.
- Pinho, S. M., & Emerenciano, M. G. C. (2021). Sensorial attributes and growth performance of whiteleg shrimp (*Litopenaeus vannamei*) cultured in biofloc technology with varying water salinity and dietary protein content. *Aquaculture*, 540. <https://doi.org/10.1016/j.aquaculture.2021.736727>
- Prihutomo, A., Triana, A., Herinto, Dongoran, R. K., & Nurdin. (2018). Pengaruh Pemuaan Pada Pemeliharaan Udang Vanamei (*Litopenaeus vannamei*, Boone) Terhadap Performa Produksi dan Kualitas Air. *Buletin Produksi Perikanan Budidaya*, 1(1), 1–10.
- Purnomo, E. A., Sutrisno, E., & Sumiyati, S. (2017). Pengaruh Variasi C/N Rasio Terhadap Produksi Kompos dan Kandungan Kalium (K), Fosfat (P) dari Batang Pisang dengan Kombinasi Kotoran Sapi dalam Sistem Vermikomposting. In *Jurnal Teknik Lingkungan*, 6(2).
- Putri, M. A. (2015). Rasio C/N terhadap Bahan Organik dan Total Bakteri pada Sedimen di Habitat Rajungan (*Portunus pelagicus*) Pantai Betahwalang, Kabupaten Demak. *Diponegoro Journal of Management of Aquatic Resources*, 4(4), 51–57.
- Rahayu, N. T., Sudrajat, S., & Hendra, M. (2022). Trophic Status of Phytoplankton as Bioindicator of Eutrophication Level of Flood Retention Ponds in Samarinda City. *Jurnal Pendidikan Matematika Dan IPA*, 13(1), 141–155. <https://doi.org/10.26418/jpmipa.v13i1.51663>
- Raja, K. (2015). Loose Shell Syndrome (LSS) in *Litopenaeus vannamei* grow-out Ponds and its Effect on Growth and Production. *Fisheries and Aquaculture Journal*, 04(04). <https://doi.org/10.4172/2150-3508.1000151>
- Ren, W., Li, L., Dong, S., Tian, X., & Xue, Y. (2019). Effects of C/N ratio and light on ammonia nitrogen uptake in *Litopenaeus vannamei* culture tanks. *Aquaculture*, 498, 123–131. <https://doi.org/10.1016/j.aquaculture.2018.08.043>
- Reynolds, C. S. (2006). *The Ecology of Phytoplankton - Ecology, Biodiversity and Conservation*. Cambridge University Press.
- Rihi, A. P. (2019). Pengaruh Pemberian Pakan Alami dan Buatan terhadap Pertumbuhan dan Kelangsungan Hidup Benih Ikan Lele Dumbo (*Clarias gariepinus* Burchell.) di Balai Benih Sentral Noekele Kabupaten Kupang. *Bioedu*, 4(2), 56–62.
- Roy, S., & Chattopadhyay, J. (2007). Towards a resolution of “the paradox of the plankton”: A brief overview of the proposed mechanisms. *Ecological Complexity*, 4(1), 26–33. <https://doi.org/10.1016/j.ecocom.2007.02.016>

- Salcher, M. M. (2014). Same same but different: Ecological niche partitioning of planktonic freshwater prokaryotes. *Journal of Limnology*, 73(1), 74–87. <https://doi.org/10.4081/jlimnol.2014.813>
- Sediadi, A. (1999). *Ekologi Dinoflagellata*. XXIV(4), 21–30. [www.oseanografi.lipi.go.id](http://www.oseanografi.lipi.go.id)
- Shirota, A. (1966). *The Plankton of South Vietnam - Fresh Water and Marine Plankton*. Overseas Technical Cooperation Agency.
- Singh, N. K., Singh, J., Bhatia, A., & Kazmi, A. A. (2016). A pilot-scale study on PVA gel beads based integrated fixed film activated sludge (IFAS) plant for municipal wastewater treatment. *Water Science and Technology*, 73(1), 113–123. <https://doi.org/10.2166/wst.2015.466>
- Supono. (2017). *Teknologi Produksi Udang*. Yogyakarta: Plantaxia. pp.1-97.
- Supono, Pinem, R. T., & Harpeni, E. (2021). Performa Udang Vaname *Litopenaeus vannamei* (Boone, 1931) yang Dipelihara pada Sistem Biofloc dengan Sumber Karbon Berbeda. *Jurnal Kelautan*, 14(2), 192–202. <https://journal.trunojoyo.ac.id/jurnalkelautan/index>
- Sweeny, B. M., & Hasting, J. W. (1958). Rhythmic Cell Division in Populations of *Gonyaulax polyedra*. *Journal of Protozoology*, 5(3), 217–224.
- ter Braak, C. J. E., & Verdonschot, P. E. M. (1995). Canonical correspondence analysis and related multivariate methods in aquatic ecology. *Aquatic Sciences*, 57.
- Triyatmo, B. (2012). *Pola Pengembangan Perikanan Budidaya Tambak Berbasis Karakteristik Lingkungan di Pesisir antara Sungai Bogowonto Kabupaten Kulonprogo dan Sungai Jali Kabupaten Purworejo*. Universitas Gadjah Mada.
- van Nieuwerburgh, L., Wänstrand, I., & Snoeijs, P. (2004). Growth and C:N:P ratios in copepods grazing on N-or Si-limited phytoplankton blooms. In *Hydrobiologia*, 514.
- Wahyu Setyaningrum, E., Dewi Masithah, E., Yuniartik, M., Tri-Kusuma Dewi, A., & Putri Nugrahani, M. (2020). Comparison of Plankton Abundance, Water Condition, Organism Growth Performance of Vaname Shrimp (*Litopenaeus vannamei*) on Intensive and Extensive Culture Systems in Banyuwangi Districts. *J Aquac Res Development*, 11(6), 597. <https://doi.org/10.35248/2155-9546.20.11.597>
- Wei, J., Zhang, X., Yu, Y., Huang, H., Li, F., & Xiang, J. (2014). Comparative transcriptomic characterization of the early development in Pacific white shrimp *Litopenaeus vannamei*. *PLoS ONE*, 9(9). <https://doi.org/10.1371/journal.pone.0106201>
- Wetzel, R. G. (2001). *Limnology: Lake and River Ecosystems*. Academic Press. pp. 205-219.
- Wetzel, R. G., & Likens, G. E. (2000b). *Limnological Analyses Third Edition*. 191p.

Widigdo, B. (2013). *Bertambak udang dengan teknologi biocrete*. Penerbit Buku Kompas. pp. 52-75.

World Register of Marine Species. (2022). *WoRMS name details Litopenaeus vannamei (Boone, 1931)*. <https://www.marinespecies.org>

WWF Indonesia. (2014). *Budidaya Udang Vaname*. [www.wwf.or.id](http://www.wwf.or.id)

Zhang, K., Yu, D., Li, Z., Xie, J., Wang, G., Gong, W., Yu, E., & Tian, J. (2020). Influence of eco-substrate addition on organic carbon, nitrogen, and phosphorus budgets of intensive aquaculture ponds of the Pearl River, China. *Aquaculture*, 520. <https://doi.org/10.1016/j.aquaculture.2019.734868>