

TABLE OF CONTENTS

| | |
|--|-------------|
| RATIFICATION PAGE | iii |
| STATEMENT | iv |
| PREFACE | vi |
| TABLE OF CONTENTS | viii |
| LIST OF FIGURES | x |
| LIST OF TABLES | xi |
| LIST OF APPENDICES | xii |
| ABSTRACT | xiii |
| INTISARI | xiv |
| CHAPTER I INTRODUCTION | 1 |
| I.1 Background | 1 |
| I.2 Research Purposes | 3 |
| I.3 Research Benefits | 3 |
| CHAPTER II LITERATURE REVIEW AND HYPOTHESIS FORMULATION | 4 |
| II.1 Literature Review | 4 |
| II.1.1 <i>Melaleuca alternifolia</i> | 4 |
| II.1.2 Essential oil | 4 |
| II.1.3 Breast cancer | 8 |
| II.1.4 Molecular docking | 10 |
| II.2 Hypothesis Formulation | 12 |
| II.2.1 Hypothesis formulation 1 | 12 |
| II.2.2 Hypothesis formulation 2 | 12 |
| II.2.3 Research plan | 13 |
| CHAPTER III RESEARCH METHODOLOGY | 14 |
| III.1 Materials | 14 |
| III.2 Equipment | 14 |
| III.3 Molecular Docking Software | 14 |
| III.4 Work Procedure | 14 |
| III.4.1 Extraction of essential oil | 14 |
| III.4.2 Molecular docking studies | 15 |
| CHAPTER IV RESULT AND DISCUSSION | 16 |
| IV.1 Tea Tree Oil's Component | 16 |

| | |
|--|-----------|
| IV.2 In Silico Anti-breast Cancer Activities of Tea Tree Oil | 22 |
| CHAPTER V CONCLUSIONS | 29 |
| V.1 Conclusion | 29 |
| V.2 Suggestion | 29 |
| REFERENCES | 30 |
| APPENDICES | 35 |

LIST OF FIGURES

| | | |
|-------------|---|----|
| Figure II.1 | Monoterpene synthesis | 6 |
| Figure II.2 | Steam distillation technique | 7 |
| Figure II.3 | Free energy bond calculation | 11 |
| Figure II.4 | Complete free energy bond calculation | 11 |
| Figure IV.1 | (a) <i>Melaleuca alternifolia</i> tree and (b) leaves and terminal branches of <i>Melaleuca alternifolia</i> planted in Kebun Raya Cibodas, Bogor | 16 |
| Figure IV.2 | A pale to the yellow color of obtained TTO | 17 |
| Figure IV.3 | Chromatogram and spectral mass of TTO extracted from <i>Melaleuca alternifolia</i> | 18 |
| Figure IV.4 | Fragmentation ion of methyl eugenol compound | 19 |
| Figure IV.5 | Fragmentation ion of methyl cis cinnamate compound | 19 |
| Figure IV.6 | Fragmentation ion of linalool compound | 19 |
| Figure IV.7 | Bond distance between (a) tamoxifen and (b) terpinene-4-ol with amino acid residues of ER α | 25 |

LIST OF TABLES

| | | |
|------------|---|----|
| Table IV.1 | The composition of TTO, which has a Similarity Index (SI) with Library Mass Spectra | 20 |
| Table IV.2 | Comparison of the percentage composition of TTO essential oil's compounds of research results with ISO 4730:2017 standards and Cibodas Botanical Garden | 21 |
| Table IV.3 | Results on docking simulation of compound obtained from TTO with tamoxifen on ER α | 24 |
| Table IV.4 | Hydrogen bond present in obtained TTO | 26 |
| Table IV.5 | Hydrophobic interaction between tamoxifen and TTO compound, and amino acid residues | 27 |

LIST OF APPENDICES

| | | |
|------------|--|----|
| Appendix 1 | Yield of Essential Oil Calculation | 35 |
| Appendix 2 | Relative Density Calculation | 35 |
| Appendix 3 | Volume of Essential Oil Calculation | 35 |
| Appendix 4 | GCMS Instrumentation Condition | 36 |
| Appendix 5 | GCMS Chromatography Result | 37 |
| Appendix 6 | 2D diagram interactions and docked poses of ligand complex of (a) tamoxifen, (b) linalool-L, (c) terpinen-4-ol, (d) p-allylanisole, (e) β -citronellol, (f) geranial, (g) methyl eugenol, (h) elemicin, (i) 1-naphthalenol, and (j) benzoic acid | 45 |