

REFERENCES

- Adey, W. H., & Loveland, K. (2007). The Input of Organic Energy. In *Dynamic Aquaria* (pp. 93–100). Elsevier. <https://doi.org/10.1016/b978-0-12-370641-6.50015-7>
- Aiken, G. R., McKnight, D. M., Wershaw, R. L., & MacCarthy, P. (1986). Humic Substances in Soil, Sediment, and Water. 1985. *Soil Sci.*, 142(5), 323. <https://doi.org/10.1097/00010694-198611000-00011>
- Amoozgar, Z., Park, J., Lin, Q., & Yeo, Y. (2012). Low Molecular-Weight Chitosan as a pH-Sensitive Stealth Coating for Tumor-Specific Drug Delivery. *Mol. Pharm.*, 9(5), 1262–1270. <https://doi.org/10.1021/mp2005615>
- Armon, R., Zolkov, C. H., & Laor, Y. (2000). Entrapment of Humic Acid in a Sol-Gel Matrix-A New Tool for Sorption Studies. In *J. Solgel Sci. Technol.* (Vol. 19).
- Auerbach, S. S., Bristol, D. W., Peckham, J. C., Travlos, G. S., Hébert, C. D., & Chhabra, R. S. (2010). Toxicity and carcinogenicity studies of methylene blue trihydrate in F344N rats and B6C3F1 mice. *Food Chem. Toxicol.*, 48(1), 169–177. <https://doi.org/10.1016/j.fct.2009.09.034>
- Baglieri, A., Vindrola, D., Gennari, M., & Negre, M. (2014). Chemical and spectroscopic characterization of insoluble and soluble humic acid fractions at different pH values. *Chem. Biol. Technol*, 1(1). <https://doi.org/10.1186/s40538-014-0009-x>
- Baigorri, R., Fuentes, M., González-Gaitano, G., & García-Mina, J. M. (2007). Simultaneous presence of diverse molecular patterns in humic substances in solution. *J. Phys. Chem. B*, 111(35), 10577–10582. <https://doi.org/10.1021/jp0738154>
- Baker, J. J. (2019). Extraction and Purification of Humic Acid from Leonardite as a Graphene Precursor for Lithium Ion Battery Cathodes. *Theses and Dissertations*, 2447. <https://commons.und.edu/theses/2447>
- Basuki, R., Santosa, S. J., & Rusdiarso, B. (2017). *The novel kinetics expression of Cadmium (II) removal using green adsorbent horse dung humic acid (Hd-Ha)*. 020001. <https://doi.org/10.1063/1.4978074>

- Berneth, H. (2008). Azine Dyes. In *Ullmann's Encyclopedia of Industrial Chemistry*. Wiley-VCH Verlag GmbH & Co. KGaA.
https://doi.org/10.1002/14356007.a03_213.pub3
- Bernoux, M., & Cerri, C. E. P. (2005). GEOCHEMISTRY | Soil, Organic Components. In *Encyclopedia of Analytical Science* (pp. 203–208). Elsevier.
<https://doi.org/10.1016/B0-12-369397-7/00245-4>
- Bleam, W. (2017). Natural Organic Matter. In *Soil and Environmental Chemistry* (pp. 333–384). Elsevier. <https://doi.org/10.1016/b978-0-12-804178-9.00007-0>
- Budnyak, T. M., Pylypchuk, I. v., Tertykh, V. A., Yanovska, E. S., & Kolodynska, D. (2015). Synthesis and adsorption properties of chitosan-silica nanocomposite prepared by sol-gel method. *Nanoscale Res. Lett.*, 10(1).
<https://doi.org/10.1186/s11671-014-0722-1>
- Cameron, R. S., Thornton, B. K., Swift, R. S., & Posner, A. M. (n.d.). *Molecular Weight And Shape Of Humic Acid From Sedimentation And Diffusion Measurements On Fractionated Extracts*.
- Capasso, S., Chianese, S., Musmarra, D., & Iovino, P. (2020). Macromolecular structure of a commercial humic acid sample. *Environments - MDPI*, 7(4).
<https://doi.org/10.3390/environments7040032>
- Cesco, C. T., Valente, A. J. M., & Paulino, A. T. (2021). Methylene blue release from chitosan/pectin and chitosan/dna blend hydrogels. *Pharmaceutics*, 13(6).
<https://doi.org/10.3390/pharmaceutics13060842>
- Chen, H., Berndtsson, R., Ma, M., & Zhu, K. (2009). Characterization of insolubilized humic acid and its sorption behaviors. *Environ. Geol.*, 57(8), 1847–1853.
<https://doi.org/10.1007/s00254-008-1472-0>
- Chen, R. P., Zhang, Y. L., Wang, X. Y., Zhu, C. Y., Ma, A. J., & Jiang, W. M. (2015). Removal of methylene blue from aqueous solution using humic-acid coated magnetic nanoparticles. *Desalin. Water Treat.*, 55(2), 539–548.
<https://doi.org/10.1080/19443994.2014.916233>

- Chen, X., Yang, X., Xiang, Y., Xu, L., & Liu, G. (2020). Humic Acid Derived Resin: An Efficient Adsorbent for Au(III) Uptake from Aqueous Acidic Solution. *J. Chem. Eng. Data*, 65(4), 1824–1832. <https://doi.org/10.1021/acs.jced.9b01085>
- Chianese, S., Fenti, A., Iovino, P., Musmarra, D., & Salvestrini, S. (2020). Sorption of organic pollutants by humic acids: A review. In *Molecules* (Vol. 25, Issue 4). MDPI AG. <https://doi.org/10.3390/molecules25040918>
- Choi, H.-J., & Yu, S.-W. (2018). Biosorption of methylene blue from aqueous solution by agricultural bioadsorbent corncob. *Environ. Eng. Res.*, 24(1), 99–106. <https://doi.org/10.4491/eer.2018.107>
- Côa, F., Strauss, M., Clemente, Z., Rodrigues Neto, L. L., Lopes, J. R., Alencar, R. S., Souza Filho, A. G., Alves, O. L., Castro, V. L. S. S., Barbieri, E., & Martinez, D. S. T. (2017). Coating carbon nanotubes with humic acid using an eco-friendly mechanochemical method: Application for Cu(II) ions removal from water and aquatic ecotoxicity. *Sci. Total Environ.*, 607–608, 1479–1486. <https://doi.org/10.1016/j.scitotenv.2017.07.045>
- Contreras, M., Grande-Tovar, C. D., Vallejo, W., & Chaves-López, C. (2019). Bio-Removal of Methylene Blue from Aqueous Solution by *Galactomyces geotrichum* KL20A. *Water*, 11(2), 282. <https://doi.org/10.3390/w11020282>
- Cychosz, K. A., & Thommes, M. (2018). Progress in the Physisorption Characterization of Nanoporous Gas Storage Materials. In *Engineering* (Vol. 4, Issue 4, pp. 559–566). Elsevier Ltd. <https://doi.org/10.1016/j.eng.2018.06.001>
- Dai, H., He, H., Lai, C., Xu, Z., Zheng, X., Yu, G., Huang, B., Pan, X., & Dionysiou, D. D. (2021). Modified humic acids mediate efficient mineralization in a photo-bio-electro-Fenton process. *Water Res.*, 190, 116740. <https://doi.org/10.1016/j.watres.2020.116740>
- de Melo, B. A. G., Motta, F. L., & Santana, M. H. A. (2016). Humic acids: Structural properties and multiple functionalities for novel technological developments. In *Mater. Sci. Eng. C* (Vol. 62, pp. 967–974). Elsevier Ltd. <https://doi.org/10.1016/j.msec.2015.12.001>

- de Souza, F., & Bragança, S. R. (2018). Extraction and characterization of humic acid from coal for the application as dispersant of ceramic powders. *J. Mater. Res. Technol.*, 7(3), 254–260. <https://doi.org/10.1016/j.jmrt.2017.08.008>
- Doolin, P. K., Alerasool, S., Zalewski, D. J., & Hoffman, J. F. (1994). Acidity studies of titania-silica mixed oxides. *Catal. Lett.*, 25(3–4), 209–223. <https://doi.org/10.1007/BF00816302>
- Duan, M., Wu, J., Xiong, Y., Fang, S., & Chen, J. (2018). Characterization and differentiation of the adsorption behavior of crystal violet and methylene blue at the silica/water interface using near field evanescent wave. *Soft Matter*, 14(36), 7516–7525. <https://doi.org/10.1039/c8sm01385c>
- Ebisike, K., Okoronkwo, A. E., & Alaneme, K. K. (2020). Synthesis and characterization of Chitosan–silica hybrid aerogel using sol-gel method. *J. King Saud Univ. Sci.*, 32(1), 550–554. <https://doi.org/10.1016/j.jksus.2018.08.005>
- El-Wakeel, S. T., Radwan, E. K., Ghafar, H. H. A., & Moursy, A. S. (2017). Humic acid-carbon hybrid material as lead (II) ions adsorbent. *Desalin. Water Treat.*, 74, 216–223. <https://doi.org/10.5004/dwt.2017.20584>
- Eshwar, M., Srilatha, M., Rekha, K. B., & Sharma, S. H. K. (2017). Characterization of Humic Substances by Functional Groups and Spectroscopic Methods. *Int. J. Curr. Microbiol. Appl. Sci.*, 6(10), 1768–1774. <https://doi.org/10.20546/ijcmas.2017.610.213>
- Franciski, M. A., Peres, E. C., Godinho, M., Perondi, D., Foletto, E. L., Collazzo, G. C., & Dotto, G. L. (2018). Development of CO₂ activated biochar from solid wastes of a beer industry and its application for methylene blue adsorption. *Waste Manage.*, 78, 630–638. <https://doi.org/10.1016/j.wasman.2018.06.040>
- Gezici, O., Guven, I., Özcan, F., Ertul, S., & Bayrakci, M. (2016). Humic-makeup approach for simultaneous functionalization of polyacrylonitrile nanofibers during electrospinning process, and dye adsorption study. *Soft Materials*, 14(4), 278–287. <https://doi.org/10.1080/1539445X.2016.1201506>

- Gezici, O., & Kara, H. (2011). Towards multimodal HPLC separations on humic acid-bonded aminopropyl silica: RPLC and HILIC behavior. *Talanta*, 85(3), 1472–1482. <https://doi.org/10.1016/j.talanta.2011.06.026>
- Ginimuge, P. R., & Jyothi, S. D. (2010). Methylene blue: revisited. *J. Anaesthesiol. Clin. Pharmacol.* 26(4), 517–520.
- Gjessing, E. T. (1976). *Physical and chemical characteristics of aquatic humus*. Ann Arbor Science.
- Guy, R. D., Narine, D. R., & deSilva, S. (1980). Organocation speciation. I. A comparison of the interactions of methylene blue and paraquat with bentonite and humic acid. *Can. J. Chem.*, 58(6), 547–554. <https://doi.org/10.1139/v80-085>
- Han, Q., Wang, J., Goodman, B. A., Xie, J., & Liu, Z. (2020). High adsorption of methylene blue by activated carbon prepared from phosphoric acid treated eucalyptus residue. *Powder Technol.*, 366, 239–248. <https://doi.org/10.1016/j.powtec.2020.02.013>
- Hassanpour, M., Safardoust-Hojaghan, H., & Salavati-Niasari, M. (2017). Degradation of methylene blue and Rhodamine B as water pollutants via green synthesized Co₃O₄/ZnO nanocomposite. *J. Mol. Liq.*, 229, 293–299. <https://doi.org/10.1016/j.molliq.2016.12.090>
- Haworth, R. D. (1971). The Chemical Nature of Humic Acid. *Soil Sci.*, 111(1), 71–79.
- Hayes, M. H. B. (1991). Concepts of the origins, composition, and structures of humic substances. In W. S. Wilson (Ed.), *Advances in Soil Organic Matter Research: The Impact on Agriculture and the Environment* (pp. 3–22). The Royal Society of Chemistry.
- Hayes, M. H. B., MacCarthy, P., Malcolm, R. L., & Swift, R. S. (1989). *Humic substances II*. John Wiley and Sons.
- Horvath, R. S., & Ropp, M. E., 1974, Mechanism of Action of Eosin-Methylene Blue Agar in the Differentiation of *Escherichia coli* and *Enterobacter aerogenes*. *Int. J. Sys. Bacteriol.*, 24(2), 221–224. <https://doi.org/10.1099/00207713-24-2-221>

- Ilharco, L. M. (1992). NON-CRYSTALLINE SOLIDS The evolution of TEOS to silica gel and glass by vibrational spectroscopy. In *J. Non-Cryst. Solids* (Vol. 147, Issue 148).
- Isolation of IHSS Soil Fulvic and Humic Acids*. (2022). International Humic Substances Society. <https://humic-substances.org/isolation-of-ihss-soil-fulvic-and-humic-acids/>
- Jarupanavet, P., & Piyamongkala, K. (2021). *Chitosan modified with sodium lauryl ether sulfate used as adsorbent of soluble oil*. 070005.
<https://doi.org/10.1063/5.0064061>
- Jawad, A. H., Abdulhameed, A. S., & Mastuli, M. S. (2020). Acid-fractionalized biomass material for methylene blue dye removal: a comprehensive adsorption and mechanism study. *J. Taibah Univ. Sci.*, 14(1), 305–313.
<https://doi.org/10.1080/16583655.2020.1736767>
- Jawad, A. H., Abdulhameed, A. S., Reghioua, A., & Yaseen, Z. M. (2020). Zwitterion composite chitosan-epichlorohydrin/zeolite for adsorption of methylene blue and reactive red 120 dyes. *Int. J. Biol. Macromol.*, 163, 756–765.
<https://doi.org/10.1016/j.ijbiomac.2020.07.014>
- Jones, M. N., & Bryan, N. D. (1998). *Colloidal properties of humic substances*.
- Joseph, A., Vellayan, K., González, B., Vicente, M. A., & Gil, A. (2019). Effective degradation of methylene blue in aqueous solution using Pd-supported Cu-doped Ti-pillared montmorillonite catalyst. *Appl. Clay Sci.*, 168, 7–10.
<https://doi.org/10.1016/j.clay.2018.10.009>
- Joseph, J., & Jemmis, E. D. (2007). Red-, blue-, or no-shift in hydrogen bonds: A unified explanation. *J. Am. Chem. Soc.*, 129(15), 4620–4632.
<https://doi.org/10.1021/ja067545z>
- Khan, I., Saeed, K., Zekker, I., Zhang, B., Hendi, A. H., Ahmad, A., Ahmad, S., Zada, N., Ahmad, H., Shah, L. A., Shah, T., & Khan, I. (2022). Review on Methylene Blue: Its Properties, Uses, Toxicity and Photodegradation. *Water*, 14(2), 242.
<https://doi.org/10.3390/w14020242>

- Kirk-Othmer Encyclopedia of Chemical Technology. (2007). *Reference Reviews*, 21(2), 49–50. <https://doi.org/10.1108/09504120710728879>
- Kleber, M., & Lehmann, J. (2019). Humic Substances Extracted by Alkali Are Invalid Proxies for the Dynamics and Functions of Organic Matter in Terrestrial and Aquatic Ecosystems. *J. Environ. Qual.*, 48(2), 207–216. <https://doi.org/10.2134/jeq2019.01.0036>
- Klučáková, M., & Kalina, M. (2015). Composition, particle size, charge, and colloidal stability of pH-fractionated humic acids. *J. Soils Sediments*, 15(9), 1900–1908. <https://doi.org/10.1007/s11368-015-1142-2>
- Klučáková, M., & Pekař, M. (2005). Solubility and dissociation of lignitic humic acids in water suspension. *Colloids Surf. A: Physicochem. Eng. Asp.*, 252(2–3), 157–163. <https://doi.org/10.1016/j.colsurfa.2004.10.019>
- Kolokassidou, C., Pashalidis, I., Costa, C. N., Efstathiou, A. M., & Buckau, G. (2007). Thermal stability of solid and aqueous solutions of humic acid. *Thermochim. Acta*, 454(2), 78–83. <https://doi.org/10.1016/j.tca.2006.12.022>
- Koopal, L. K., Yang, Y., Minnaard, A. J., Theunissen, P. L. M., & van Riemsdijk, W. H. (1998). Chemical immobilisation of humic acid on silica. *Colloids Surf. A: Physicochem. Eng. Asp.*, 141(3), 385–395. [https://doi.org/10.1016/S0927-7757\(97\)00170-2](https://doi.org/10.1016/S0927-7757(97)00170-2)
- Kumar Gautam, R., Navaratna, D., Muthukumaran, S., Singh, A., Islamuddin, & More, N. (2021). Humic Substances: Its Toxicology, Chemistry and Biology Associated with Soil, Plants and Environment. In *Humic Substance [Working Title]*. IntechOpen. <https://doi.org/10.5772/intechopen.98518>
- Kurt, E., Koseoglu-Imer, D. Y., Dizge, N., Chellam, S., & Koyuncu, I. (2012). Pilot-scale evaluation of nanofiltration and reverse osmosis for process reuse of segregated textile dyewash wastewater. *Desalination*, 302, 24–32. <https://doi.org/10.1016/j.desal.2012.05.019>
- Leong, Y. K. (2005). Yield stress and zeta potential of nanoparticulate silica dispersions under the influence of adsorbed hydrolysis products of metal ions—Cu(II), Al(III)

- and Th(IV). *J. Colloid Interface Sci.*, 292(2), 557–566.
<https://doi.org/10.1016/j.jcis.2005.06.004>
- Li, C., Zhao, S., Huang, X., Xie, D., Li, X., Ma, J., & Liao, Y. (2022). Development of humic acid based adsorbents for fast and efficient removal of ammonia and organic nitrogen from super magnetic separation treated wastewater. *J. Environ. Chem. Eng.*, 10(2), 107223. <https://doi.org/10.1016/j.jece.2022.107223>
- Li, D., Hua, T., Yuan, J., & Xu, F. (2021). Methylene blue adsorption from an aqueous solution by a magnetic graphene oxide/humic acid composite. *Colloids Surf. A: Physicochem. Eng. Asp.*, 627. <https://doi.org/10.1016/j.colsurfa.2021.127171>
- Lin, J., Zhan, Y., & Zhu, Z. (2011). Adsorption characteristics of copper (II) ions from aqueous solution onto humic acid-immobilized surfactant-modified zeolite. *Colloids Surf. A: Physicochem. Eng. Asp.*, 384(1–3), 9–16.
<https://doi.org/10.1016/j.colsurfa.2011.02.044>
- Liu, J. F., Zhao, Z. S., & Jiang, G. bin. (2008). Coating Fe₃O₄ magnetic nanoparticles with humic acid for high efficient removal of heavy metals in water. *Environ. Sci. Technol.*, 42(18), 6949–6954. <https://doi.org/10.1021/es800924c>
- Liu, T., Li, Y., Du, Q., Sun, J., Jiao, Y., Yang, G., Wang, Z., Xia, Y., Zhang, W., Wang, K., Zhu, H., & Wu, D. (2012). Adsorption of methylene blue from aqueous solution by graphene. *Colloids Surf. B Biointerfaces*, 90, 197–203.
<https://doi.org/10.1016/j.colsurfb.2011.10.019>
- Liu, Y. guo, Li, T. ting, Zeng, G. ming, Zheng, B. hong, Xu, W. hua, & Liu, S. bo. (2016). Removal of Pb(II) from aqueous solution by magnetic humic acid/chitosan composites. *J. Cent. South Univ.*, 23(11), 2809–2817.
<https://doi.org/10.1007/s11771-016-3344-1>
- Lodygin, E. D., Alekseev, I. I., Vasilevich, R. S., & Abakumov, E. V. (2020). Complexation of lead and cadmium ions with humic acids from arctic peat soils. *Environ. Res.*, 191. <https://doi.org/10.1016/j.envres.2020.110058>
- Lu, S., Liu, W., Wang, Y., Zhang, Y., Li, P., Jiang, D., Fang, C., & Li, Y. (2019). An adsorbent based on humic acid and carboxymethyl cellulose for efficient dye

- removal from aqueous solution. *Int. J. Biol. Macromol.*, *135*, 790–797.
<https://doi.org/10.1016/j.ijbiomac.2019.05.095>
- Luo, W. J., Gao, Q., Wu, X. L., & Zhou, C. G. (2014). Removal of Cationic Dye (Methylene Blue) from Aqueous Solution by Humic Acid-Modified Expanded Perlite: Experiment and Theory. *Sep. Sci. Technol.*, *49*(15), 2400–2411.
<https://doi.org/10.1080/01496395.2014.920395>
- Lv, S. W., Liu, J. M., Ma, H., Wang, Z. H., Li, C. Y., Zhao, N., & Wang, S. (2019). Simultaneous adsorption of methyl orange and methylene blue from aqueous solution using amino functionalized Zr-based MOFs. *Microporous Mesoporous Mater.*, *282*, 179–187. <https://doi.org/10.1016/j.micromeso.2019.03.017>
- MacCarthy, P. (2001). THE PRINCIPLES OF HUMIC SUBSTANCES. *Soil Sci.*, *166*(11), 738–751. <https://doi.org/10.1097/00010694-200111000-00003>
- Maccarthy, P., Clapp, C. E., & Bloom, P. R. (1990). *Chapter 1 An Introduction to Soil Humic Substances*.
- Malatji, N., Makhado, E., Modibane, K. D., Ramohlola, K. E., Maponya, T. C., Monama, G. R., & Hato, M. J. (2021). Removal of methylene blue from wastewater using hydrogel nanocomposites: A review. In *Nanomaterials and Nanotechnology* (Vol. 11). SAGE Publications Ltd. <https://doi.org/10.1177/18479804211039425>
- Marques Neto, J. de O., Bellato, C. R., Milagres, J. L., Pessoa, K. D., & Alvarenga, E. S. de. (2013). Preparation and evaluation of chitosan beads immobilized with Iron(III) for the removal of As(III) and As(V) from water. *J. Braz. Chem. Soc.*, *24*(1), 121–132. <https://doi.org/10.1590/S0103-50532013000100017>
- Masini, J. C., Abate, G., Lima, E. C., Hahn, L. C., Nakamura, M. S., Lichtig, J., & Nagatomy, Â. R. (1998). *Comparison of methodologies for determination of carboxylic and phenolic groups in humic acids*. 223–233.
- Methylene Blue. (2016). In *Some Drugs and Herbal Products: IARC Monographs on the Evaluation of Carcinogenic Risks to Humans* (Vol. 108, pp. 155–183). International Agency for Research on Cancer.

- Mir, S. H., Nagahara, L. A., Thundat, T., Mokarian-Tabari, P., Furukawa, H., & Khosla, A. (2018). Review—Organic-Inorganic Hybrid Functional Materials: An Integrated Platform for Applied Technologies. *J. Electrochem. Soc.*, *165*(8), B3137–B3156. <https://doi.org/10.1149/2.0191808jes>
- Mohammed, M. I., Ismael, M. K., & Gönen, M. (2020). Synthesis of Chitosan-Silica Nanocomposite for Removal of Methyl Orange from Water: Composite Characterization and Adsorption Performance. *IOP Conference Series: Mater. Sci. Eng.*, *745*(1). <https://doi.org/10.1088/1757-899X/745/1/012084>
- Mouni, L., Belkhir, L., Bollinger, J. C., Bouzaza, A., Assadi, A., Tirri, A., Dahmoune, F., Madani, K., & Remini, H. (2018). Removal of Methylene Blue from aqueous solutions by adsorption on Kaolin: Kinetic and equilibrium studies. *Appl. Clay Sci.*, *153*, 38–45. <https://doi.org/10.1016/j.clay.2017.11.034>
- Munir, M., Nazar, M. F., Zafar, M. N., Zubair, M., Ashfaq, M., Hosseini-Bandegharai, A., Khan, S. U.-D., & Ahmad, A. (2020). Effective Adsorptive Removal of Methylene Blue from Water by Didodecyldimethylammonium Bromide-Modified Brown Clay. *ACS Omega*, *5*(27), 16711–16721. <https://doi.org/10.1021/acsomega.0c01613>
- Nageeb, M. (2013). Adsorption Technique for the Removal of Organic Pollutants from Water and Wastewater. In *Organic Pollutants - Monitoring, Risk and Treatment*. InTech. <https://doi.org/10.5772/54048>
- Ononye, A. I., & Graveel, J. G. (1994). Modeling the reactions of 1-naphthylamine and 4-methylaniline with humic acids: Spectroscopic investigations of the covalent linkages. *Environ. Toxicol. Chem.*, *13*(4), 537–541. <https://doi.org/10.1002/etc.5620130401>
- Ononye, A. I., Graveel, J. G., & Wolt, J. D. (1989). Kinetic Studies of the Reactions of Benzidine with Humic Acid. *Soil Sci Soc Am J*, *53*(3), 981–983. <https://doi.org/10.2136/sssaj1989.03615995005300030060x>

- Orsi, M. (2014). REVIEW Open Access Molecular dynamics simulation of humic substances. In *Orsi Chem. Biol. Technol* (Vol. 1).
<http://www.chembioagro.com/content/>
- Pawlak, A., & Mucha, M. (2003). Thermogravimetric and FTIR studies of chitosan blends. In *Thermochim. Acta* (Vol. 396).
- Peng, L., Qin, P., Lei, M., Zeng, Q., Song, H., Yang, J., Shao, J., Liao, B., & Gu, J. (2012). Modifying Fe₃O₄ nanoparticles with humic acid for removal of Rhodamine B in water. *J. Hazard. Mater.*, 209–210, 193–198.
<https://doi.org/10.1016/j.jhazmat.2012.01.011>
- Peres, E. C., Slaviero, J. C., Cunha, A. M., Hosseini-Bandegharai, A., & Dotto, G. L. (2018). Microwave synthesis of silica nanoparticles and its application for methylene blue adsorption. *J. Environ. Chem. Eng.*, 6(1), 649–659.
<https://doi.org/10.1016/j.jece.2017.12.062>
- Piccolo, A. (2001). The Supramolecular Structure of Humic Substances. *Soil Sci.*, 166(11), 810–832.
- Piccolo, A., & Camici, L. (1990). A comparison of two methods for the determination of total acidity of humic substances. *Int J Environ Anal Chem.*, 41(1–2), 65–69.
<https://doi.org/10.1080/03067319008030530>
- Piccolo, A., & Mirabella, A. (1987). Molecular weight distribution of peat humic substances extracted with different inorganic and organic solutions. *Sci. Total Environ.*, 62, 39–46. [https://doi.org/10.1016/0048-9697\(87\)90479-7](https://doi.org/10.1016/0048-9697(87)90479-7)
- Pota, G., Venezia, V., Vitiello, G., di Donato, P., Mollo, V., Costantini, A., Avossa, J., Nuzzo, A., Piccolo, A., Silvestri, B., & Luciani, G. (2020). Tuning Functional Behavior of Humic Acids through Interactions with Stöber Silica Nanoparticles. *Polymers*, 12(4), 982. <https://doi.org/10.3390/polym12040982>
- Prado, A. G. S., Miranda, B. S., & Jacintho, G. V. M. (2003). Interaction of indigo carmine dye with silica modified with humic acids at solid/liquid interface. *Surf. Sci.*, 542(3), 276–282. [https://doi.org/10.1016/S0039-6028\(03\)00990-7](https://doi.org/10.1016/S0039-6028(03)00990-7)

- Prado, A. G. S., Miranda, B. S., & Zara, L. F. (2005). Adsorption and thermochemical data of divalent cations onto silica gel surface modified with humic acid at solid/liquid interface. *J. Hazard. Mater.*, 120(1–3), 243–247.
<https://doi.org/10.1016/j.jhazmat.2005.01.011>
- Prado, A. G. S., Pertusatti, J., & Nunes, A. R. (2011). Aspects of protonation and deprotonation of humic acid surface on molecular conformation. *J. Braz. Chem. Soc.*, 22(8), 1478–1483. <https://doi.org/10.1590/S0103-50532011000800011>
- Prasetyo, E. (2018). Humic acid provenance influence to the adsorption capacity in uranium and thorium removal. *IOP Conference Series: Mater. Sci. Eng.*, 285, 012011. <https://doi.org/10.1088/1757-899X/285/1/012011>
- Prasetyo, E., & Toyoda, K. (2016). Sol–gel synthesis of a humic acid-silica gel composite material as low-cost adsorbent for thorium and uranium removal. *J. Radioanal. Nucl. Chem.*, 310(1), 69–80. <https://doi.org/10.1007/s10967-016-4861-y>
- Prasetyo, E., & Toyoda, K. (2021). Humic acid attachment on chitosan-modified silica gel as an economical, efficient, and selective adsorbent for thorium and uranium removal. *Environ. Technol.*, 1–15.
<https://doi.org/10.1080/09593330.2021.1968038>
- Proctor, A., & Toro-Vazquez, J. F. (2009). The Freundlich Isotherm in Studying Adsorption in Oil Processing. In *Bleaching and Purifying Fats and Oils: Theory and Practice* (pp. 209–219). Elsevier Inc. <https://doi.org/10.1016/B978-1-893997-91-2.50016-X>
- Qian, F., Sun, X., & Liu, Y. (2013). Removal characteristics of organics in bio-treated textile wastewater reclamation by a stepwise coagulation and intermediate GAC/O₃ oxidation process. *Chem. Eng. J.*, 214, 112–118.
<https://doi.org/10.1016/j.cej.2012.09.130>
- Queiroz, M. F., Melo, K. R. T., Sabry, D. A., Sassaki, G. L., & Rocha, H. A. O. (2015). Does the use of chitosan contribute to oxalate kidney stone formation? *Marine Drugs*, 13(1), 141–158. <https://doi.org/10.3390/md13010141>

- Rashed, M. N., & Palanisamy, P. N. (2018). Introductory Chapter: Adsorption and Ion Exchange Properties of Zeolites for Treatment of Polluted Water. In *Zeolites and Their Applications*. InTech. <https://doi.org/10.5772/intechopen.77190>
- Revellame, E. D., Fortela, D. L., Sharp, W., Hernandez, R., & Zappi, M. E. (2020). Adsorption kinetic modeling using pseudo-first order and pseudo-second order rate laws: A review. In *Cleaner Engineering and Technology* (Vol. 1). Elsevier Ltd. <https://doi.org/10.1016/j.clet.2020.100032>
- Rubio, F., Rubio, J., & Oteo, J. L. (1998). A FT-IR study of the hydrolysis of Tetraethylorthoselicate (TEOS). *Spectrosc. Lett.*, 31(1), 199–219. <https://doi.org/10.1080/00387019808006772>
- Ruiz-Hitzky, E., Ariga, K., & Lvov, Y. M. (Eds.). (2007). *Bio-inorganic Hybrid Nanomaterials*. Wiley. <https://doi.org/10.1002/9783527621446>
- Rusdiarso, B., & Basuki, R. (2020). Stability Improvement of Humic Acid as Sorbent through Magnetite and Chitin Modification. *J. Kim. Sains Apl.*, 23(5), 152–159. <https://doi.org/10.14710/jksa.23.5.152-159>
- Salazar-Rabago, J. J., Leyva-Ramos, R., Rivera-Utrilla, J., Ocampo-Perez, R., & Cerino-Cordova, F. J. (2017). Biosorption mechanism of Methylene Blue from aqueous solution onto White Pine (*Pinus durangensis*) sawdust: Effect of operating conditions. *Sustain. Environ. Res.*, 27(1), 32–40. <https://doi.org/10.1016/j.serj.2016.11.009>
- Sanchez, C., Julián, B., Belleville, P., & Popall, M. (2005). Applications of hybrid organic-inorganic nanocomposites. *J. Mater. Chem.*, 15(35–36), 3559–3592. <https://doi.org/10.1039/b509097k>
- Sánchez- Monedero, M. A., Roig, A., Cegarra, J., Bernal, M. P., & Paredes, C. (2002). Effects of HCl-HF purification treatment on chemical composition and structure of humic acids. *European Journal of Soil Sci.*, 53(3), 375–381. <https://doi.org/10.1046/j.1365-2389.2002.00464.x>
- Sándor, M., Nistor, C. L., Szalontai, G., Stoica, R., Nicolae, C. A., Alexandrescu, E., Fazakas, J., Oancea, F., & Donescu, D. (2016). Aminopropyl-Silica Hybrid Particles

- as Supports for Humic Acids Immobilization. *Materials (Basel, Switzerland)*, 9(1).
<https://doi.org/10.3390/ma9010034>
- Santos, R. R., Vermeulen, S., Haritova, A., & Fink-Gremmels, J. (2011). Isotherm modeling of organic activated bentonite and humic acid polymer used as mycotoxin adsorbents. *Food Addit. Contam.: Part A*, 28(11), 1578–1589.
<https://doi.org/10.1080/19440049.2011.595014>
- Santosa, S. J., Siswanta, D., Sudiono, S., & Utarianingrum, R. (2008). Chitin–humic acid hybrid as adsorbent for Cr(III) in effluent of tannery wastewater treatment. *Applied Surf. Sci.*, 254(23), 7846–7850.
<https://doi.org/10.1016/j.apsusc.2008.02.102>
- Santosa, S.J., Krisbiantoro, P.A., Yuniarti, M., Kustomo, & Koesnarpardi, S. (2021). Magnetically separable humic acid-functionalized magnetite for reductive adsorption of tetrachloroaurate(III) ion in aqueous solution. *Environ. Nanotechnol. Monit. Manag.*, 15. <https://doi.org/10.1016/j.enmm.2021.100454>
- Saparpakorn, P., Kim, J. H., & Hannongbua, S. (2007). molecules Investigation on the Binding of Polycyclic Aromatic Hydrocarbons with Soil Organic Matter: A Theoretical Approach. In *Molecules* (Vol. 12). <http://www.mdpi.org>
- Senesi, N., & Testini, C. (1984). Theoretical aspects and experimental evidence of the capacity of humic substances to bind herbicides by charge-transfer mechanisms (electron donor-acceptor processes). *Chemosphere*, 13(3), 461–468.
[https://doi.org/10.1016/0045-6535\(84\)90104-8](https://doi.org/10.1016/0045-6535(84)90104-8)
- Singare, P. U., & Dhabarde, S. S. (2013). Pollution Discharge Scenario of Dyeing Industries along Dombivali Industrial Belt of Mumbai, India. *Int. Lett. Chem. Phys. Astron.*, 22, 48–55. <https://doi.org/10.18052/www.scipress.com/ilcpa.22.48>
- Slama, H. ben, Chenari Bouket, A., Pourhassan, Z., Alenezi, F. N., Silini, A., Cherif-Silini, H., Oszako, T., Luptakova, L., Golińska, P., & Belbahri, L. (2021). Diversity of Synthetic Dyes from Textile Industries, Discharge Impacts and Treatment Methods. *Appl. Sci.*, 11(14), 6255. <https://doi.org/10.3390/app11146255>

- Smilek, J., Sedláček, P., Kalina, M., & Klučáková, M. (2015). On the role of humic acids' carboxyl groups in the binding of charged organic compounds. *Chemosphere*, 138, 503–510. <https://doi.org/10.1016/j.chemosphere.2015.06.093>
- Song, M., Song, B., Meng, F., Chen, D., Sun, F., & Wei, Y. (2019). Incorporation of humic acid into biomass derived carbon for enhanced adsorption of phenol. *Sci. Rep.*, 9(1). <https://doi.org/10.1038/s41598-019-56425-8>
- Spark, K. M., Wells, J. D., & Johnson, B. B. (1997). The interaction of a humic acid with heavy metals. *Soil Res.*, 35(1), 89. <https://doi.org/10.1071/S96008>
- Sparks, D. L. (2003). Chemistry of Soil Organic Matter. In *Environmental Soil Chemistry* (pp. 75–113). Elsevier. <https://doi.org/10.1016/B978-012656446-4/50003-7>
- Stathi, P., & Deligiannakis, Y. (2010). Humic acid-inspired hybrid materials as heavy metal absorbents. *J. Colloid Interface Sci.*, 351(1), 239–247. <https://doi.org/10.1016/j.jcis.2010.06.023>
- Steelink, C. (1963). What is humic acid? *J. Chem. Educ.*, 40(7), 379. <https://doi.org/10.1021/ed040p379>
- Steelink, C. (1985). Implication of elemental characteristics of humic substances. In *Humic Substances in Soil, Sediment and Water: Geochemistry, Isolation, and Characterization* (pp. 457–476). Wiley-Interscience Publication.
- Stevenson, F. J. (1994). *Humus Chemistry: Genesis, Composition, Reactions* (2nd ed.). John Wiley & Sons, Inc.
- Stevenson, F. J., & Goh, K. M. (1971). Infrared spectra of humic acids and related substances. *Geochim. Cosmochim. Acta*, 35(5), 471–483. [https://doi.org/10.1016/0016-7037\(71\)90044-5](https://doi.org/10.1016/0016-7037(71)90044-5)
- Tan, K. H. (2014). *Humic Matter in Soil and the Environment* (2nd ed.). CRC Press.
- Tang, X., Ran, G., Li, J., Zhang, Z., & Xiang, C. (2021). Extremely efficient and rapidly adsorb methylene blue using porous adsorbent prepared from waste paper: Kinetics and equilibrium studies. *J. Hazard. Mater.*, 402, 123579. <https://doi.org/10.1016/j.jhazmat.2020.123579>

- Tejeda-Agredano, M.-C., Mayer, P., & Ortega-Calvo, J.-J. (2014). The effect of humic acids on biodegradation of polycyclic aromatic hydrocarbons depends on the exposure regime. *Environ. Pollut.*, 184, 435–442.
<https://doi.org/10.1016/j.envpol.2013.09.031>
- Thommes, M., Kaneko, K., Neimark, A. v., Olivier, J. P., Rodriguez-Reinoso, F., Rouquerol, J., & Sing, K. S. W. (2015). Physisorption of gases, with special reference to the evaluation of surface area and pore size distribution (IUPAC Technical Report). *Pure Appl. Chem.*, 87(9–10), 1051–1069.
<https://doi.org/10.1515/pac-2014-1117>
- Thompson, A., & Goyne, K. W. (2012). Introduction to the Sorption of Chemical Constituents in Soils. *Nat. Sci. Educ.*, 4(4), 7.
- Thurman, E. M. (1985). Aquatic Humic Substances. In *Organic Geochemistry of Natural Waters* (pp. 273–361). Springer Netherlands. https://doi.org/10.1007/978-94-009-5095-5_11
- van Stempvoort, D. R., Lesage, S., Novakowski, K. S., Millar, K., Brown, S., & Lawrence, J. R. (2002). Humic acid enhanced remediation of an emplaced diesel source in groundwater. *J. Contam. Hydrol.*, 54(3–4), 249–276.
[https://doi.org/10.1016/S0169-7722\(01\)00182-6](https://doi.org/10.1016/S0169-7722(01)00182-6)
- von Wandruszka, R. (2000). Humic acids: Their detergent qualities and potential uses in pollution remediation. *Geochem. Trans.*, 1, 10–15.
<https://doi.org/10.1039/b001869o>
- WAKSMAN, S. A. (1936). Humus Origin, Chemical Composition, and Importance in Nature. *Soil Sci.*, 41(5), 395. <https://doi.org/10.1097/00010694-193605000-00010>
- Wang, Y., Huang, X., Lv, X., Sun, J., Zheng, R., & Hu, K. (2020). The Metabolism of Methylene Blue and Its Derivatives in Japanese Eel (*Anguilla Japonica*). *J. Ocean Univ. China*, 19(3), 653–658. <https://doi.org/10.1007/s11802-020-4166-1>
- Wei, W., Han, X., Zhang, M., Zhang, Y., & Zheng, C. (2020). Macromolecular humic acid modified nano-hydroxyapatite for simultaneous removal of Cu(II) and methylene

- blue from aqueous solution: Experimental design and adsorption study. *Int. J. Biol. Macromol.*, 150, 849–860. <https://doi.org/10.1016/j.ijbiomac.2020.02.137>
- Wershaw, R. L. (1999). MOLECULAR AGGREGATION OF HUMIC SUBSTANCES. *Soil Sci.*, 164(11), 803–813. <https://doi.org/10.1097/00010694-199911000-00004>
- Wershaw, R. L. (2000). THE STUDY OF HUMIC SUBSTANCES—IN SEARCH OF A PARADIGM. In *Humic Substances* (pp. 1–7). Elsevier. <https://doi.org/10.1016/B978-1-85573-807-2.50005-9>
- William Kajjumba, G., Emik, S., Öngen, A., Kurtulus Özcan, H., & Aydın, S. (2019). Modelling of Adsorption Kinetic Processes—Errors, Theory and Application. In *Advanced Sorption Process Applications*. IntechOpen. <https://doi.org/10.5772/intechopen.80495>
- Xu, J.-Z., Dai, L., Wu, B., Ding, T., Zhu, J.-J., Lin, H., Chen, H.-L., Shen, C.-Y., & Jiang, Y. (2009). Determination of methylene blue residues in aquatic products by liquid chromatography-tandem mass spectrometry. *J. Sep. Sci.*, 32(23–24), 4193–4199. <https://doi.org/10.1002/jssc.200900364>
- Xu, P., Wang, Y., Li, X., Chen, Q., Hao, L., Zhang, J., Zhu, X., & Jia, B. (2020). An acidic-groups detection method and its application to analysis of Chinese humic acid samples. *PLOS ONE*, 15(8), e0238061. <https://doi.org/10.1371/journal.pone.0238061>
- Yan, W. L., & Bai, R. (2005). Adsorption of lead and humic acid on chitosan hydrogel beads. *Water Res.*, 39(4), 688–698. <https://doi.org/10.1016/j.watres.2004.11.007>
- Yang, H. C., Gong, J. L., Zeng, G. M., Zhang, P., Zhang, J., Liu, H. Y., & Huan, S. Y. (2017). Polyurethane foam membranes filled with humic acid-chitosan crosslinked gels for selective and simultaneous removal of dyes. *J. Colloid Interface Sci.*, 505, 67–78. <https://doi.org/10.1016/j.jcis.2017.05.075>
- Yang, S., Zong, P., Ren, X., Wang, Q., & Wang, X. (2012). Rapid and highly efficient preconcentration of Eu(III) by core-shell structured Fe₃O₄@Humic acid magnetic nanoparticles. *ACS Appl. Mater. Interfaces*, 4(12), 6891–6900. <https://doi.org/10.1021/am3020372>

- Yu, X. Z., Feng, Y. X., & Yue, D. (2015). Phytotoxicity of methylene blue to rice seedlings. *Glob. J. Environ. Sci. Manag.*, 1, 199–204.
- Zakirov, A. S., Navamathavan, R., Kang, T. W., Kim, C. Y., Choi, C. K., Ihm, J., & Cheong, H. (2011). *Effect of Ultraviolet Irradiation on the Defect States and Charge Transport Properties of Low-k SiOC(-H) Dielectric Films Deposited by UV-Assisted PECVD*. 71–72. <https://doi.org/10.1063/1.3666262>
- Zhang, W., Zhang, L. Y., Zhao, X. J., & Zhou, Z. (2016). Citrus pectin derived porous carbons as a superior adsorbent toward removal of methylene blue. *J. Solid State Chem.*, 243, 101–105. <https://doi.org/10.1016/j.jssc.2016.08.014>
- Zhao, P., Huang, Z., Ma, Q., Zhang, B., & Wang, P. (2022). Artificial humic acid synthesized from food wastes: An efficient and recyclable adsorbent of Pb (II) and Cd (II) from aqueous solution. *Environ. Technol. Innov.*, 27. <https://doi.org/10.1016/j.eti.2022.102399>