

## DAFTAR PUSTAKA

- Abdel-Aal, E.S.M., Young, J.C., and Rabalski, I. (2006). Anthocyanin composition in black, blue, pink, purple and red cereal grains. *Journal Agriculture Food Chemistry*, 54:4696–4704.
- Alessandra, C. P., Daniel, G., and Neiva, D. R. (2016). Extaction of anthocyanins and polyphenols from black rice (*Oryza sativa* L.) by modeling and assessing their reversibility and stability. *Food Chemistry*, 191, 12–20.
- Apridamayanti, P., Pratiwi, R., Purwestri, Y. A., Tunjung, W. A. S., and Rumiya, R. (2017). Anthocyanin, nutrient contents, and antioxidant activity of black rice bran of *Oryza sativa* L. ‘Cempo Ireng’ from Sleman, Yogyakarta, Indonesia. *Indonesian Journal of Biotechnology*, 22(1): 49-54.
- Bridgers, E.N., Chinn, M. S., and Truong, V.D. (2010). Extraction of anthocyanins from industrial purple-fleshed sweetpotatoes and enzymatic hydrolysis of residues for fermentable sugars. *Industrial Crops and Products*, 32:613-620.
- Castañeda-Ovando, A., Pacheco-Hernández, M. D. L., Páez-Hernández, M. E., Rodríguez, J. A., and Galán-Vidal, C. A. (2009). Chemical studies of anthocyanins: A review. *Food Chemistry*, 113: 859-871.
- Chaiyasut, C., Sivamaruthi, B. S., Pengkumsri, N., Sirilun, S., Peerajan, S., and Kesika, P. (2016). Anthocyanin profile and its antioxidant activity of widely used fruits, vegetables, and flowers in Thailand. *Asian Journal of Pharmaceutical and Clinical Research*, 9(6): 218-224.
- Chandrasekhar, J., Madhusudhan, M. C., and Raghavarao, K. (2012). Extraction of anthocyanins from red cabbage and purification using adsorption. *Food and Bioproducts Processing*, 90: 615-623.
- Deng, J., Chen, S., Yin, X., Wang, K., Liu, Y., Li, S., and Yang, P. (2013). Systematic qualitative and quantitative assessment of anthocyanins, flavones and flavonols in the petals of 108 lotus (*Nelumbo nucifera*) varieties. *Food chemistry*, 139(1-4):307-312.
- Farahmandazad, H. (2015). *Recovery and purification of anthocyanins from purple-blue potato. Thesis.*
- Fatchiyah, F., Sari, D. R. T., Safitri, A., and Cairns, J. R. (2020). Phytochemical compound and nutritional value in black rice from Java Island, Indonesia. *Sys Rev Pharm*, 11(7): 414-421.
- Fatonah, N., Nora, I., and Harlia. (2016). Uji Stabilitas Zat Warna Ekstrak Buah Senggani (*Melastoma malabathricum* L.). *Jurnal Kimia Khatulistiwa*, 5(1).
- Giusti, M. M., and Wrolstad, R. E. (2003). Acylated anthocyanins from edible sources and their applications in food systems. *Biochemical Engineering Journal*, 14: 217-225.
- Harborne, J. B. (1962). Anthocyanins and their sugar components. In *Fortschritte der Chemie Organischer Naturstoffe/Progress in the Chemistry of Organic Natural Products/Progrès dans la Chimie des Substances Organiques Naturelles*. Springer, Vienna. pp. 165-199.
- Hellström, J., Mattila, P., and Karjalainen, R. (2013). Stability of anthocyanins in berry juices stored at different temperatures. *Journal of Food Composition and Analysis*, 31(1): 12–19.

- Hosseinian, F. S., Li, W., and Beta, T. (2008). Measurement of anthocyanins and other phytochemicals in purple wheat. *Food Chemistry*, 109: 916-924.
- Hou, Z., Qin, P., Zhang, Y., Cui, S., and Ren, G. (2013). Identification of anthocyanins isolated from black rice (*Oryza sativa* L.) and their degradation kinetics. *Food Research International*, 50(2): 691–697.
- Ignat, I., Volf, I., and Popa, V. I. (2011). A critical review of methods for characterisation of polyphenolic compounds in fruits and vegetables. *Food Chemistry*, 126: 1821-35.
- Ilmi, W., Pratiwi, R., and Purwestri, Y. A. (2018). Total anthocyanin content and antioxidant activity of Brown rice, endosperm, and rice bran of three Indonesian Black rice (*Oryza sativa* L.) cultivars. In *Proceeding of the 2nd International Conference on Tropical Agriculture* (pp. 205-216). Springer, Cham.
- Iosub, I., Kajzar, F., Makowska-Janusik, M., Meghea, A., Tane, A., and Rau, I. (2012). Electronic structure and optical properties of some anthocyanins extracted from grapes. *Optical Materials*, 34(10): 1644-1650.
- Jamei, Rashid, and Farideh, B. (2017). Stability of blueberry (*Cornus mas* Yulyush) anthocyanin pigment under pH and co-pigment treatments. *International Journal of Food Properties*, 20(9): 2128-2133.
- Kang, Y. J., Jung, S. W., and Lee, S. J. (2014). An optimal extraction solvent and purification adsorbent to produce anthocyanins from black rice (*Oryza sativa* cv. Heugjinjubyeo). *Food Science and Biotechnology*, 23: 97-106.
- Kaneda, I., Iwai, K., Kubo, H., and Sakurai, H. (2004). Antioxidative properties of extracts from ancient rice brans. *Food Science and Technology Research*, 10: 374–382.
- Kim, M. K., Kim, H. A., Koh, K., Kim, H. S., Lee, Y. S., and Kim, Y. H. (2008). Identification and quantification of anthocyanin pigments in colored rice. *Nutrition Research and Practice*, 2(1):46.
- Kim, C. K., Kikuchi, S., Kim, Y.K., Park, S.H., Yoon, U., Lee, G.S., Choi, J.W., Kim, Y.H., and Park, S.C. (2010) Computational identification of seed specific transcription factors involved in anthocyanin production in black rice. *Biochip Journal*, 4(3):247–255.
- Kristamtini, K., Wiranti, E.W., and Sutarno, S. (2018). Variasi Warna dan Kandungan Antosianin Kultivar Lokal Beras Hitam Yogyakarta pada Dua Ketinggian. *Buletin Plasma Nutfah*, 24(2): 97-106.
- Kushwaha, U. K. S. (2016). Black rice. In *Black Rice*. Springer. pp. 21-47.
- Lapornik, B., Prošek, M. and Golc W. A. (2005). Comparison of extracts prepared from plant by-products using different solvents and extraction time. *Journal of Food Engineering*, 71: 214-222.
- Lee, J. H. (2010). Identification and quantification of anthocyanins from the grains of black rice (*Oryza sativa* L.) varieties. *Food Science and Biotechnology*, 19(2): 391–397.
- Lima, Á. S., Soares, C. M. F., Paltram, R., Halbwirth, H., and Bica, K. (2017). Extraction and consecutive purification of anthocyanins from grape pomace using ionic liquid solutions. *Fluid Phase Equilibria*, 451: 68-78.
- Markakis, P and Jurd, L. (1974). Anthocyanins and their stability in foods, *C R C Critical Reviews in Food Technology*, 4:4: 437-456

- Moldovan, B., David, L., Chisbora, C., and Cimpoiu, C. (2012). Degradation kinetics of anthocyanins from European cranberrybush (*Viburnum opulus* L.) fruit extracts, Effects of temperature, pH and storage solvent. *Molecules*, 17: 11655-66.
- Noviyanto, F. 2020. *Penetapan Kadar Ketoprofen dengan Metode Spektrofotometri UV-Vis*. Media Sains Indonesia. Bandung. pp. 2-6.
- Park, Y. S., Kim, S. J., and Chang, H. I. (2008). Isolation of anthocyanin from black rice (Heugjinjubyeo) and screening of its antioxidant activities. *Microbiology and Biotechnology Letters*, 36(1): 55-60.
- Patil, G., Madhusudhan, M. C., Ravindra Babu, B., and Raghavarao, K. S. M. S. (2009). Extraction, dealcoholization and concentration of anthocyanin from red radish. *Chemical Engineering and Processing: Process Intensification*, 48(1): 364–369.
- Pedro, A. C., Granato, D., and Rosso, N. D. (2016). Extraction of anthocyanins and polyphenols from black rice (*Oryza sativa* L.) by modeling and assessing their reversibility and stability. *Food Chemistry*, 191: 12-20.
- Pratiwi, R., Purwestri, Y. A., Tunjung, W. A. S. (2014). Effect of diet rice pellets (*Oryza sativa* L.) ‘Cempo Ireng’, ‘Cempo Abang’, and ‘IR-64’ to lipid serum profile of white rat (*Rattus norvegicus* Berkenhout, 1769) hyperllipidemia. In *Proceeding of Seminar and Workshop on Nutrigenomic, Yogyakarta March* (pp. 12-18).
- Pratiwi, R., Tunjung, W. A. S., Rumiati., and Amalia, A. R. (2015). Black rice bran extracts and fractions containing Cyanidin 3-glucoside and Peonidin 3-glucoside induce apoptosis in human cervical cancer cells. *Indonesian Journal of Biotechnology*, 20(1): 69-76.
- Pratiwi, R., and Purwestri, Y. A. (2017). Black rice as a functional food in Indonesia. *Functional Foods in Health and Disease*, 7(3): 182-194.
- Priska, M., Peni, N., Carvallo, L., and Ngapa, Y. D. (2018). Antosianin dan Pemanfaatannya. *Cakra Kimia (Indonesian E-Journal of Applied Chemistry)*, 6(2): 79-97.
- Puertolas, E., Cregenzan, O., Luengo, E., Alvarez, I., and Raso, J. (2013). Pulsedelectric-field-assisted extraction of anthocyanins from purple-fleshed potato. *Food Chemistry*, 136: 1330-6.
- Raharjo, S., Purwandari, F. A., Hastuti, P., and Olsen, K. (2019). Stabilization of black rice (*Oryza sativa*, L. *indica*) anthocyanins using plant extracts for copigmentation and maltodextrin for encapsulation. *Journal of food science*, 84(7): 1712-1720.
- Rakkimuthu, R., Palmurugan, S., & Shanmugapriya, A. (2016). Effect of temperature, light, pH on the stability of anthocyanin pigments in cocculus hirsutus fruits. *International Journal of Multidisciplinary Research and Modern Education*, 2(2): 91-96.
- Sa’adah, I. R., Supriyanto., and Subejo. (2013). Keragaman warna gabah dan warna beras kultivar lokal padi beras hitam (*Oryza Sativa* L.) yang dibudidayakan oleh petani Kabupaten Sleman, Bantul, dan Magelang. *Vegetalika*, 2(3): 13-20.
- Setyaningrum, E. N. (2010). Efektivitas Penggunaan Jenis Asam dalam Proses Ekstraksi Pigmen Antosianin Kulit Manggis (*Garcinia mangostana* L.) dengan Penambahan Aseton 60%. *Skripsi*.



- Sipahli, S., Mohanlall, V., and Mellem, J. J. (2017). Stability and Degradation Kinetics of Crude Anthocyanin Extract from *Hibiscus sabdariffa*. *Food Science and Technology*, 37 (2): 209 – 215.
- Siregar, Y. D. I., and Nurlela, N. (2012). Ekstraksi dan uji stabilitas zat warna alami dari bunga kembang sepatu (*Hibiscus rosa-sinensis* L) dan bunga rosela (*Hibiscus sabdariffa* L.). *Jurnal Kimia Valensi*: 2(3).
- Spagna, G., Barbagallo, R. N., Todaro, A., Durante, M. J., and Pifferi, P. G. (2003). A method for anthocyanin extraction from fresh grape skin. *Italian journal of food science*, 15(3): 337-346.
- Sui, X., Dong, X., and Zhou, W. (2014). Combined effect of pH and high temperature on the stability and antioxidant capacity of two anthocyanins in aqueous solution. *Food Chemistry*, 163: 163–170.
- Surh, J., and Eunmi, K. (2014). Effects of four different cooking methods on anthocyanins, total phenolics and antioxidant activity of black rice. *Journal of the Science of Food and Agriculture*, 94(15): 3296-3304.
- Tai, L., Huang, S., Zhao, Z., and Huang, G. (2020). Chemical composition analysis and antioxidant activity of black rice pigment. *Chemical Biology and Drug Design*, 97(3): 711-720.
- Tjitrosoepomo, G. 2005. *Taksonomi Tumbuhan*. Gadjah Mada University Press. Yogyakarta.
- Wang, B. C., He, R., and Li, Z. M. (2010). The stability and antioxidant activity of anthocyanins from blueberry. *Food technology and biotechnology*, 48(1): 42.
- Xue, L., Wang, Z., Zhang, W., Li, Y., Wang, J., and Lei, J. (2016). Flower pigment inheritance and anthocyanin characterization of hybrids from pink-flowered and white-flowered strawberry. *Scientia Horticulturae*, 200: 143-150.