

## DAFTAR PUSTAKA

- Acosta-Cárdenas, A., Alcaraz-Zapata, W., & Cardona-Betancur, M. 2018. Sugarcane molasses and vinasse as a substrate for polyhydroxyalkanoates (PHA) production. *DYNA (Colombia)*, 85(206), 220–225.
- Albuquerque, M. G. E., Eiroa, M., Torres, C., Nunes, B. R., & Reis, M. A. M. (2007). Strategies for the development of a side stream process for polyhydroxyalkanoate (PHA) production from sugar cane molasses. *Journal of Biotechnology*, 130(4), 411–421.
- Ali, W. S., Zaki, N. H., & Obiad, S. Y. N. (2017). Production of bioplastic by bacteria isolated from local soil and organic wastes. *Current Research in Microbiology and Biotechnology*, 5(2), 1012–1017.
- Aljuraifani, A. A., Berekaa, M. M., & Ghazwani, A. A. (2019). Bacterial biopolymer (polyhydroxyalkanoate) production from low-cost sustainable sources. *MicrobiologyOpen*, 8(6), 1–7.
- Alkotaini, B., Sathiyamoorthi, E., & Kim, B. S. (2015). Potential of *Bacillus megaterium* for production of polyhydroxyalkanoates using the red algae *Gelidium amansii*. *Biotechnology and Bioprocess Engineering*, 20(5), 856–860.
- Amache, R., Sukan, A., Safari, M., Roy, I., & Keshavarz, T. (2013). Advances in PHAs production. *Chemical Engineering Transactions*, 32, 931–936.
- Bengtsson, S., Pisco, A. R., Reis, M. A. M., & Lemos, P. C. (2010). Production of polyhydroxyalkanoates from fermented sugar cane molasses by a mixed culture enriched in glycogen accumulating organisms. *Journal of Biotechnology*, 145(3), 253–263.
- Bharti, S., & Swetha, G. (2016). Need for Bioplastics and Role of Biopolymer PHB: A Short Review. *Journal of Petroleum & Environmental Biotechnology*, 07(02), 2–4.
- Bozorg, A., Vossoughi, M., Kazemi, A., & Alemzadeh, I. (2015). Optimal medium composition to enhance poly- $\beta$ -hydroxybutyrate production by *Ralstonia eutropha* using cane molasses as sole carbon source. *Applied Food Biotechnology*, 2(3), 39–47.
- Bresan, S., Sznajder, A., Hauf, W., Forchhammer, K., Pfeiffer, D., & Jendrosseck, D. (2016). Polyhydroxyalkanoate (PHA) granules have no phospholipids. *Scientific Reports*, 6(March), 1–13.
- Brinda Devi, A., Valli Nachiyar, C., Kaviyarasi, T., & Samrot, A. V. (2015). Characterization of polyhydroxybutyrate synthesized by *Bacillus Cereus*. *International Journal of Pharmacy and Pharmaceutical Sciences*, 7(3), 140–144.
- Castilho, L. R., Mitchell, D. A., & Freire, D. M. G. (2009). Production of polyhydroxyalkanoates (PHAs) from waste materials and by-products by submerged and solid-state fermentation. *Bioresource Technology*, 100(23), 5996–6009.
- Catone, M. V., Ruiz, J. A., Castellanos, M., Segura, D., Espin, G., & López, N. I. (2014). High polyhydroxybutyrate production in *Pseudomonas extremaustralis* is associated with differential expression of horizontally acquired and core genome polyhydroxyalkanoate synthase genes. *PLoS ONE*, 9(6).

- Ceyhan, N., & Ozdemir, G. (2011). Poly--hydroxybutyrate (PHB) production from domestic wastewater using *Enterobacter aerogenes* 12Bi strain. *African Journal of Microbiology Research*, 5(6), 690–702.
- Chen, Z., Huang, L., Wen, Q., & Guo, Z. (2015). Efficient polyhydroxyalkanoate (PHA) accumulation by a new continuous feeding mode in three-stage mixed microbial culture (MMC) PHA production process. *Journal of Biotechnology*, 209(Mmc), 68–75.
- Chung, H. Y., Lee, K. H., Ryu, S., Yoon, H., Lee, J. H., Kim, H. B., Kim, H., Jeong, H. G., Choi, S. H., & Kim, B. S. (2016). Genome sequence of *Bacillus cereus* FORC\_021, a food-borne pathogen isolated from a knife at a sashimi restaurant. *Journal of Microbiology and Biotechnology*, 26(12), 2030–2035.
- Das, R., Pal, A., & Paul, A. K. (2019). Bioconversion of sugarcane molasses to poly(3-hydroxybutyrate-co-3-hydroxyvalerate) by endophytic *Bacillus cereus* RCL 02. *Journal of Applied Biology and Biotechnology*, 7(2), 20–24.
- Desetty, R. D., Mahajan, V. S., Khan, B. M., & Rawal, S. K. (2008). Isolation and heterologous expression of PHA synthesising genes from *Bacillus thuringiensis* R1. *World Journal of Microbiology and Biotechnology*, 24(9), 1769–1774.
- Furuta, Y., Tsujinouchi, M., Shawa, M., Zorigt, T., Miyajima, Y., Paudel, A., Suzuki, S., & Higashi, H. (2022). Complete Genome Sequences of 24 Strains of *Bacillus cereus* Isolated from Nosocomial Infection and Bacteremia Cases in Japan. *Microbiology Resource Announcements*, 11(4), 11–13.
- Gatea, Iman H., Abbas, Alaa S., Abid Ameena G., Halob Amal A., Maied, Sana K and Abidali, A. S. (2018). Isolation And Characterization Of *Pseudomonas Putida* Producing Bioplastic. *Pak. J. Biotechnol*, 15(2), 469–473.
- Gomaa, E. Z. (2014). Production of Polyhydroxyalkanoates (PHAs) By *Bacillus subtilis* and *Escherichia coli* Grown on Cane Molasses Fortified with Ethanol. *Brazilian Archives Of Biology And Technology Arch. Biol. Technol.* V, 57157(1), 145–154.
- Gozan, M., Hermansyah, H., & Alamsyah, R. (2016). Pengaruh Jenis Sumber Nitrogen Pada Pembuatan Polyhydroxybutyrate Dari Glukosa Menggunakan Bakteri *Bacillus cereus*. *Seminar Nasional Sains Dan Teknologi, UMJ, November*, 1–5.
- Haedar, N., Gobel, R. B., Umar, M. R., & Ambeng. (2014). Isolation and characterization of bacteria from Waste Sugar Mill Arasoe-Kab.Bone As Raw Material Producing Bioplastics Degraded (Poly-B-hydroxybutyrate). *Jurnal Alam Dan Ilmu Lingkungan, Vol.5*(8), 1–7.
- Hammami, K., Souissi, Y., Souii, A., Ouertani, A., El-Hidri, D., Jabberi, M., Chouchane, H., Mosbah, A., Masmoudi, A. S., Cherif, A., & Neifar, M. (2022). Extremophilic Bacterium *Halomonas desertis* G11 as a Cell Factory for Poly-3-Hydroxybutyrate-co-3-Hydroxyvalerate Copolymer's Production. *Frontiers in Bioengineering and Biotechnology*, 10(May), 1–16.
- Haruhiro, F., Wataru, I., Katsuyuki, N., Joko, P., Hiroe, T., & Genshiro, K. (2015). A Comparative Study of Waste Cooking Oil Recycling Programs in Bogor and Niigata Cities and GHG Emission Reduction by Recycling. *Proceedings of the 2015 AASRI International Conference on Circuits and Systems*, 9(Cas), 169–172.

- Hendrawan, Y., Alvianto, D., Sumarlan, S. H., & Wibisono, Y. (2020). Characterization of *Pseudomonas fluorescens* polyhydroxyalkanoate. *Advances in Food Science, Sustainable Agriculture and Agroindustrial Engineering (AFSSAAE)*, 3(1), 1–10.
- Huang, Chien-Jui; Zheng, Po-Xing; Ou, Heng-Yang; Lin, Yao-Cheng; Chen, C.-Y. (2017). crossm Complete Genome Sequence of *Bacillus Rhizobacterium* from the Rhizosphere of. *American Society of Microbiology*, 5(46), 1–2.
- Janaki, R. R. V., & Kamala, K. S. S. (2021). Bioconversion of agro - industrial effluents for polyhydroxyalkanoates production using *Bacillus subtilis* RS1. *International Journal of Environmental Science and Technology*, 16(10), 5725–5734.
- Jun, L., Wu, Y., Yuan, J., Yuan, J., Wang, Z., Gao, J., & Chen, H. (2021). Characterization of *Bacillus cereus* AFA01 Capable of Degrading. *MDPI*, 10(1725), 1–14.
- Kamilah, H., Al-Gheethi, A., Yang, T. A., & Sudesh, K. (2018). The Use of Palm Oil-Based Waste Cooking Oil to Enhance the Production of Polyhydroxybutyrate [P(3HB)] by *Cupriavidus necator* H16 Strain. *Arabian Journal for Science and Engineering*, 43(7), 3453–3463.
- Kanso, S., & Patel, B. K. C. (2003). *Microvirga subterranea* gen. nov., sp. nov., a moderate thermophile from a deep subsurface Australian thermal aquifer. *International Journal of Systematic and Evolutionary Microbiology*, 53(2), 401–406.
- Khatami, K., Perez-Zabaleta, M., Owusu-Agyeman, I., & Cetecioglu, Z. (2020). Waste to bioplastics: How close are we to sustainable polyhydroxyalkanoates production? *Waste Management*, 119, 374–388.
- Kongpeng, C., Iewkittayakorn, J., & Chotigeat, W. (2017). Effect of Storage Time and concentration of used cooking oil on polyhydroxyalkanoates (PHAs) production by *Cupriavidus necator* H16. *Sains Malaysiana*, 46(9), 1465–1469.
- Kourmentza, C., & Kornaros, M. (2016). Biotransformation of volatile fatty acids to polyhydroxyalkanoates by employing mixed microbial consortia: The effect of pH and carbon source. *Bioresource Technology*, 222, 388–398.
- Kresnawaty, I., Prakoso, H. T., Eris, D. D., & Mulyatni, A. S. (2016). Penapisan bakteri penghasil bioplastik polihidroksialkanoat dari tanah tempat pembuangan sampah dan limbah cair pabrik kelapa sawit Screening of bioplastics polyhydroxyalkanoic producing bacteria from landfill and palm oil mill effluents. *E-Journal Menara Perkebunan*, 82(1).
- Lee, D. H., Kim, H. R., Chung, H. Y., Lim, J. G., Kim, S., Kim, S. K., Ku, H. J., Kim, H., Ryu, S., Choi, S. H., & Lee, J. H. (2015). Complete genome sequence of *Bacillus cereus* FORC\_005, a food-borne pathogen from the soy sauce braised fish-cake with quail-egg. *Standards in Genomic Sciences*, 10(1), 4–11.
- Lee, K. M., & Gilmore, D. F. (2005). Formulation and process modeling of biopolymer (polyhydroxyalkanoates: PHAs) production from industrial wastes by novel crossed experimental design. *Process Biochemistry*, 40(1), 229–246.
- Liu, C. C., Zhang, L. L., An, J., Chen, B., & Yang, H. (2016). Recent strategies for efficient production of polyhydroxyalkanoates by micro-organisms. *Letters in Applied Microbiology*,

- Liu, F., Li, J., & Zhang, X. L. (2019). Bioplastic production from wastewater sludge and application. *IOP Conference Series: Earth and Environmental Science*, 344(1).
- Lopes, M., Miranda, S. M., & Belo, I. (2020). Microbial valorization of waste cooking oils for valuable compounds production—a review. *Critical Reviews in Environmental Science and Technology*, 50(24), 2583–2616.
- López-Cortés, A., Rodríguez-Fernández, O., Latisnere-Barragán, H., Mejía-Ruiz, H. C., González-Gutiérrez, G., & Lomelí-Ortega, C. (2010). Characterization of polyhydroxyalkanoate and the *phaC* gene of *Paracoccus seriniphilus* E71 strain isolated from a polluted marine microbial mat. *World Journal of Microbiology and Biotechnology*, 26(1), 109–118.
- Luengo, J. M., García, B., Sandoval, A., Naharro, G., & Olivera, E. R. (2003). Bioplastics from microorganisms. *Current Opinion in Microbiology*, 6(3), 251–260.
- Maheshwari, N., Kumar, M., Thakur, I. S., & Srivastava, S. (2018). Production, process optimization and molecular characterization of polyhydroxyalkanoate (PHA) by CO<sub>2</sub> sequestering *B. cereus* SS105. *Bioresource Technology*, 254(November 2017), 75–82.
- Melani, A., Herawati, N., & Kurniawan, A. F. (2017). Bioplastik Pati Umbi Talas Melalui Proses Melt Intercalation (Kajian Pengaruh Jenis Filler, Konsentrasi Filler dan Jenis Plasticizer). *Distilasi*, 2(2), 53–67.
- Merugu, R., Girisham, S., & Reddy, S. M. (2010). Production of PHB (Polyhydroxybutyrate) by *Rhodospirillum rubrum* KU003 under Nitrogen Limitation. *International Journal of Applied Biology and Pharmaceutical Technology*, 1(3), 847–850.
- Mitra, R., Xu, T., Xiang, H., & Han, J. (2020). Current developments on polyhydroxyalkanoates synthesis by using halophiles as a promising cell factory. *Microbial Cell Factories*, 19(1), 1–30.
- Mostafa, Y. S., Alrumman, S. A., Otaif, K. A., Alamri, S. A., Mostafa, M. S., & Sahlabji, T. (2020). Production and characterization of bioplastic by polyhydroxybutyrate accumulating *Erythrobacter aquimaris* isolated from mangrove rhizosphere. *Molecules*, 25(1).
- Naheed, N., & Jamil, N. (2014). Optimization of biodegradable plastic production on sugar cane molasses in *Enterobacter sp.* SEL2. *Brazilian Journal of Microbiology*, 45(2), 417–426.
- Narita, V., Arum, A. L. A., M, S. I., & Fawzya, N. Y. (2012). Analisis Bioinformatika Berbasis WEB untuk Eksplorasi Enzim Kitosanase Berdasarkan Kemiripan Sekuens. *Jurnal Al-Azhar Indonesia Seri Sains Dan Teknologi*, 1(4), 197–203.
- Nielsen, C., Rahman, A., Rehman, A. U., Walsh, M. K., & Miller, C. D. (2017). Food waste conversion to microbial polyhydroxyalkanoates. *Microbial Biotechnology*, 10(6), 1338–1352.
- Noda, I., Satkowski, M. M., Dowrey, A. E., & Marcott, C. (2004). Polymer alloys of nodax copolymers and poly(lactic acid). *Macromolecular Bioscience*, 4(3), 269–275.
- Obruca, S., Marova, I., Snajdar, O., Mravcova, L., & Svoboda, Z. (2010). Production of poly(3-

- hydroxybutyrate-co-3-hydroxyvalerate) by *Cupriavidus necator* from waste rapeseed oil using propanol as a precursor of 3-hydroxyvalerate. *Biotechnology Letters*, 32(12), 1925–1932.
- Pagliano, G., Ventorino, V., Panico, A., & Pepe, O. (2017). Integrated systems for biopolymers and bioenergy production from organic waste and by-products: A review of microbial processes. *Biotechnology for Biofuels*, 10(1), 1–24.
- Patel, M., Gapes, D. J., Newman, R. H., & Dare, P. H. (2009). Physico-chemical properties of polyhydroxyalkanoate produced by mixed-culture nitrogen-fixing bacteria. *Applied Microbiology and Biotechnology*, 82(3), 545–555.
- Penido, A., Mendes, P., Campos, I., & Mendes, L. (2013). Waste cooking oil as substrate for biosynthesis of poly(3-hydroxybutyrate) and poly(3-hydroxybutyrate-co-3-hydroxyhexanoate): Turning waste into a value-added product. *Malaysian Journal of Microbiology*, 9(2), 166–175.
- Rehm, B. H. A. (2006). Genetics and biochemistry of polyhydroxyalkanoate granule self-assembly: The key role of polyester synthases. *Biotechnology Letters*, 28(4), 207–213.
- Salmiati, Ujang, Z., Salim, M. R., & Olsson, G. (2009). Recovery of Polyhydroxyalkanoates (PHAs) from Mixed Microbial Cultures by Simple Digestion and Saponification. *Proceedings of the 3rd International Water Association (IWA)-ASPIRE Conference and Exhibition*, 8–15.
- Sehgal, R., & Gupta, R. (2020). Polyhydroxyalkanoate and its efficient production: an eco-friendly approach towards development. *3 Biotech*, 10(12), 1–14.
- Shrivastav, A., Kim, H. Y., & Kim, Y. R. (2013). Advances in the applications of polyhydroxyalkanoate nanoparticles for novel drug delivery system. *BioMed Research International*, 2013(Mcl).
- Sistem informasi pengelolaan sampah nasional (SIPSN). 2021. (<https://sipsn.menlhk.go.id/sipsn/public/data/komposisi>)
- Surendran, A., Lakshmanan, M., Chee, J. Y., Sulaiman, A. M., Thuoc, D. Van, & Sudesh, K. (2020). Can Polyhydroxyalkanoates Be Produced Efficiently From Waste Plant and Animal Oils? *Frontiers in Bioengineering and Biotechnology*, 8(March), 1–15.
- Taniguchi, I., Kagotani, K., & Kimura, Y. (2003). Microbial production of poly(hydroxyalkanoate)s from waste edible oils. *Green Chemistry*, 5(5), 545–548.
- Tariq, A., Hameed, A., Bokhari, H., & Masood, F. (2014). Is atomic rearrangement of type IV PHA synthases responsible for increased PHA production? *Journal of Biomolecular Structure and Dynamics*, 33(6), 1225–1238.
- Teeka, J., Imai, T., Reungsang, A., Cheng, X., Yuliani, E., Thiantanankul, J., Poomipuk, N., Yamaguchi, J., Jeenanong, A., Higuchi, T., Yamamoto, K., & Sekine, M. (2012). Characterization of polyhydroxyalkanoates (PHAs) biosynthesis by isolated *Novosphingobium* sp. THA-AIK7 using crude glycerol. *Journal of Industrial Microbiology and Biotechnology*, 39(5), 749–758.



- Thode Filho, S., Paiva, J. L. de, Franco, H. A., Perez, D. V., & Marques, M. R. da C. (2017). Environmental Impacts Caused By Residual Vegetable Oil in the Soil-Plant System. *Ciência e Natura*, 39(3), 748.
- Valappil, S. P., Rai, R., Bucke, C., & Roy, I. (2008). Polyhydroxyalkanoate biosynthesis in *Bacillus cereus* SPV under varied limiting conditions and an insight into the biosynthetic genes involved. *Journal of Applied Microbiology*, 104(6), 1624–1635.
- Verlinden, R. A. J., Hill, D. J., Kenward, M. A., Williams, C. D., Piotrowska-Seget, Z., & Radecka, I. K. (2011). Production of polyhydroxyalkanoates from waste frying oil by *Cupriavidus necator*. *AMB Express*, 1(1), 1–8.
- Wagle, A. R., Dixit, Y. M., & Vakil, B. V. (2019). Scale Up Studies for Polyhydroxyalkanoate Production by a *Bacillus flexus* Strain with Industrial Potential. *Indian Journal of Microbiology*, 59(3), 383–386.
- Wei, Y. H., Chen, W. C., Huang, C. K., Wu, H. S., Sun, Y. M., Lo, C. W., & Janarthanan, O. M. (2011). Screening and evaluation of polyhydroxybutyrate-producing strains from indigenous isolate *Cupriavidus taiwanensis* strains. *International Journal of Molecular Sciences*, 12(1), 252–265.
- Yanti, N. A., & Margino, S. (2010). Optimasi Produksi Poli-  $\beta$  -Hidroksibutirat ( PHB ) oleh *Bacillus sp* . PSA10 Pendahuluan Metode Penelitian. 15(3), 331–339.
- Yang, C., Zhang, W., Liu, R., Zhang, C., Gong, T., Li, Q., Wang, S., & Song, C. (2013). Analysis of polyhydroxyalkanoate (PHA) synthase gene and PHA-producing bacteria in activated sludge that produces PHA containing 3-hydroxydodecanoate. *FEMS Microbiology Letters*, 346(1), 56–64.