

### Daftar Pustaka

- Achard, C.S., & Laybutt, D.R., 2012. Lipid-induced endoplasmic reticulum stress in liver cells results in two distinct outcomes: Adaptation with enhanced insulin signaling or insulin resistance. *Endocrinology* 153: 2164–2177. doi:10.1210/en.2011-1881
- Al-Khalili, L., Forsgren, M., Kannisto, K., Zierath, J.R., Lönnqvist, F., & Krook, A., 2005. Enhanced insulin-stimulated glycogen synthesis in response to insulin, metformin or rosiglitazone is associated with increased mRNA expression of GLUT4 and peroxisomal proliferator activator receptor gamma co-activator 1. *Diabetologia* 48: 1173–1179. doi:10.1007/s00125-005-1741-3
- Al-Saraj, F., 2015. Pathogenesis of type 2 diabetes mellitus, in: Intech. *IntechOpen*, pp. 61–104. doi:10.5772/59183
- Alaaeldin, R., Abdel-Rahman, I.A.M., Hassan, H.A., Youssef, N., Allam, A.E., Abdelwahab, S.F., et al., 2021. Carpachromene ameliorates insulin resistance in HepG2 cells via modulating IR/IRS1/PI3K/Akt/GSK3/Foxo1 pathway. *Molecules* 26(24): 1-13. doi:10.3390/molecules26247629
- Alabsi, A.M., Lim, K.L., Paterson, I.C., Ali-Saeed, R., & Muharram, B.A., 2016. Cell cycle arrest and apoptosis induction via modulation of mitochondrial integrity by Bcl-2 family members and caspase dependence in *Dracaena cinnabari*-treated H400 human oral squamous cell carcinoma. *Biomed Res. Int.* 1–13. doi:10.1155/2016/4904016
- Alkhateeb, H., Chabowski, A., Glatz, J.F.C., Luiken, J.F.P., & Bonen, A., 2007. Two phases of palmitate-induced insulin resistance in skeletal muscle: Impaired GLUT4 translocation is followed by a reduced GLUT4 intrinsic activity. *Am. J. Physiol. - Endocrinol. Metab* 293: 783–793. doi:10.1152/ajpendo.00685.2006
- American Diabetes Association (ADA), 2010. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 33. doi:10.2337/dc10-S062
- Amorim, R., Simões, I.C.M., Veloso, C., Carvalho, A., Simões, R.F., Pereira, F.B., et al., 2021. Exploratory data analysis of cell and mitochondrial high-fat, high-sugar toxicity on human HepG2 cells. *Nutrients* 13: 1–27. doi:10.3390/nu13051723
- Ariga, M., Nedachi, T., Katagiri, H., & Kanzaki, M., 2008. Functional role of sortilin in myogenesis and development of insulin-responsive glucose transport system in C2C12 myocytes. *J. Biol. Chem* 283: 10208–10220. doi:10.1074/jbc.M710604200
- Arumugasamy, K., Latha, K., & Kumar, N., 2004. Studies on some pharmacognostic profiles of *Swietenia Macrophylla*. King. *Anc. Sci. Life* 24: 97–102.
- Arzumanian, V.A., Kiseleva, O.I., & Poverennaya, E. V., 2021. The curious case of the HepG2 cell line: 40 years of expertise. *Int. J. Mol. Sci.* 22: 13135. doi:10.3390/ijms222313135
- Asmat, U., Abad, K., & Ismail, K., 2016. Diabetes mellitus and oxidative stress - A concise review. *Saudi Pharm. J.* 24: 547–553. doi:10.1016/j.jsps.2015.03.013
- Atkinson, B.J., Griesel, B.A., King, C.D., Josey, M.A., & Olson, A.L., 2013. Moderate GLUT4 overexpression improves insulin sensitivity and fasting triglyceridemia in high-fat diet-fed transgenic mice. *Diabetes* 62: 2249–2258. doi:10.2337/db12-1146
- Auyeung, K.K., Law, P.C., & Shun Ko, J.K., 2009. *Astragalus saponins* induce apoptosis via an ERK-independent NF- $\kappa$ B signaling pathway in human hepatocellular HepG2 cell line. *Int. J. Mol. Med.* 23: 189-196. doi: [https://doi.org/10.3892/ijmm\\_00000116](https://doi.org/10.3892/ijmm_00000116)
- Azzahra, R.M.I., 2018. Analisis morfofisiologis mahoni (*Swietenia macrophylla* King.). *Kehutanan*. Universitas Hassanudin.
- Bai, J., Zhu, Y., Li, J., Zhang, Y., Dong, Y., & Xiao, X., 2020. Effects of *Bitter melon saponin* on the glucose and lipid metabolism in HepG2 cell and *C. elegans*. *J. Food*



- Balijepalli, M.K., Suppaiah, V., Chin, A.M., Buru, A.S., Sagineedu, S.R., & Pichika, M.R., 2015. Acute oral toxicity studies of *Swietenia macrophylla* seeds in sprague dawley rats. *Pharmacognosy Res.* 7: 38–44. doi:10.4103/0974-8490.147197
- Barky, A. El, & Hussein, S.A., 2017. Saponins and their potential role in diabetes mellitus. *Diabetes Manag.* 7: 148–158.
- Belfiore, A., Frasca, F., Pandini, G., Sciacca, L., & Vigneri, R., 2009. Insulin receptor isoforms and insulin receptor/insulin-like growth factor receptor hybrids in physiology and disease. *Endocr. Rev.* 30: 586–623. doi:10.1210/er.2008-0047
- Bhavsar, S.K., Föller, M., Gu, S., Vir, S., Shah, M.B., Bhutani, K.K., *et al.*, 2009a. Involvement of the PI3K/AKT pathway in the hypoglycemic effects of saponins from *Helicteres isora*. *J. Ethnopharmacol.* 126: 386–396. doi:10.1016/j.jep.2009.09.027
- Bhavsar, S.K., Singh, S., Giri, S., Jain, M.R., & Santani, D.D., 2009b. Effect of saponins from *Helicteres isora* on lipid and glucose metabolism regulating genes expression. *J. Ethnopharmacol.* 124: 426–433. doi:10.1016/j.jep.2009.05.041
- Bilous, R., & Donnelly, R., 2014. Buku Pegangan Diabetes (translated from Handbook of Diabetes), 4th ed. Bumi Medika, Jakarta.
- Björnholm, M., & Zierath, J.R., 2005. Insulin signal transduction in human skeletal muscle: Identifying the defects in type II diabetes. *Biochem. Soc. Trans.* 33: 354–357. doi:10.1042/BST0330354
- Blahova, J., Martiniakova, M., Babikova, M., Kovacova, V., Mondockova, V., & Omelka, R., 2021. Pharmaceutical drugs and natural therapeutic products for the treatment of type 2 diabetes mellitus, *Pharmaceuticals* 14(8): 1-32. doi: 10.3390/ph14080806
- Boncler, M., Golanski, J., Lukasiak, M., Redzynia, M., Dastyh, J., & Watala, C., 2017. A new approach for the assessment of the toxicity of polyphenol-rich compounds with the use of high content screening analysis. *PLoS One* 12: 1–25. doi:10.1371/journal.pone.0180022
- Bonora, E., Kiechl, S., Willeit, J., Oberhollenzer, F., Egger, G., Targher, G., *et al.*, 1998. Prevalence of insulin resistance in metabolic disorders: The Bruneck Study. *Diabetes* 47: 1643–1649. doi:10.2337/diabetes.47.10.1643
- Boura-Halfon, S., & Zick, Y., 2009. Phosphorylation of IRS proteins, insulin action, and insulin resistance. *Am. J. Physiol. - Endocrinol. Metab.* 296(4):E581-591. doi:10.1152/ajpendo.90437.2008
- BPOM RI, 2014. Peraturan Kepala Badan Pengawas Obat dan Makanan Republik Indonesia Nomor 13 Tahun 2014, BPOM.
- Bryner, R.W., Woodworth-Hobbs, M.E., Williamson, D.L., & Alway, S.E., 2012. Docosahexaenoic acid protects muscle cells from palmitate-induced Aarophy. *ISRN Obes.* 2012: 1–14. doi:10.5402/2012/647348
- Bunel, V., Ouedraogo, M., Nguyen, A.T., Stévigny, C., & Duez, P., 2014. Methods applied to the in vitro primary toxicology testing of natural products: State of the art, strengths, and limits. *Planta Med.* 80: 1210–1226. doi:10.1055/s-0033-1360273
- Burattini, S., Ferri, R., Battistelli, M., Curci, R., Luchetti, F., & Falcieri, E., 2004. C2C12 murine myoblasts as a model of skeletal muscle development: Morpho-functional characterization. *Eur. J. Histochem.* 48: 223–233. doi:10.4081/891
- Burry, R.W., 2011. Controls for immunocytochemistry: An update. *J. Histochem. Cytochem.* 59: 6–12. doi:10.1369/jhc.2010.956920
- Cai, X., Hu, X., Cai, B., Wang, Q., Li, Y., Tan, X., *et al.*, 2013. Metformin suppresses hepatocellular carcinoma cell growth through induction of cell cycle G1/G0 phase arrest and p21CIP and p27KIP expression and downregulation of cyclin D1 in vitro and in vivo. *Oncol. Rep.* 30: 2449–2457. doi:10.3892/or.2013.2718



- N.A.D.P., Saldanha, L.L., Vareda, P.M.P., Rezende-Neto, J.M., Senger, M.R., Delgado, A.Q., *et al.*, 2019. Hypoglycaemic activity of *Bauhinia holophylla* through GSK3- $\beta$  inhibition and glycogenesis activation. *Pharm. Biol.* 57: 269–279. doi:10.1080/13880209.2019.1599962
- Cao, J., Meng, S., Chang, E., Beckwith-Fickas, K., Xiong, L., Cole, R.N., *et al.*, 2014. Low concentrations of metformin suppress glucose production in hepatocytes through AMP-activated protein kinase (AMPK). *J. Biol. Chem.* 289: 20435–20446. doi:10.1074/jbc.M114.567271
- CCRC UGM, 2012. Prosedur tetap. *Uji Sitotoksik Metod. MTT* 1–7.
- Chabowski, A., Zendzian-Piotrowska, M., Konstantynowicz, K., Pankiewicz, W., Mikłosz, A., Łukaszuk, B., *et al.*, 2013. Fatty acid transporters involved in the palmitate and oleate induced insulin resistance in primary rat hepatocytes. *Acta Physiol.* 207: 346–357. doi:10.1111/apha.12022
- Chavez, J.A., & Summers, S.A., 2003. Characterizing the effects of saturated fatty acids on insulin signaling and ceramide and diacylglycerol accumulation in 3T3-L1 adipocytes and C2C12 myotubes. *Arch. Biochem. Biophys.* 419: 101–109. doi:10.1016/j.abb.2003.08.020
- Chen, C., Han, X., Dong, P., Li, Z., Yanagita, T., Xue, C., *et al.*, 2018a. *Sea cucumber saponin* liposomes ameliorate obesity-induced inflammation and insulin resistance in high-fat-diet-fed mice. *Food Funct.* 9: 861–870. doi:10.1039/c7fo01599b
- Chen, J.W., Kong, Z.L., Tsai, M.L., Lo, C.Y., Ho, C.T., & Lai, C.S., 2018b. *Tetrahydrocurcumin* ameliorates free fatty acid-induced hepatic steatosis and improves insulin resistance in HepG2 cells. *J. Food Drug Anal.* 26: 1075–1085. doi:10.1016/j.jfda.2018.01.005
- Chen, X., & Thibeault, S., 2013. Effect of DMSO concentration, cell density and needle gauge on the viability of cryopreserved cells in three dimensional hyaluronan hydrogel. *Proc. Annu. Int. Conf. IEEE Eng. Med. Biol. Soc. EMBS* 6228–6231. doi:10.1109/EMBC.2013.6610976
- Cho, H.-K., Lee, J.-Y., Jang, Y., & Kwon, Y.H., 2008. Involvement of endoplasmic reticulum stress in almitate-induced apoptosis in HepG2 Cells. *Toxicol. Res.* 24: 129–135.
- Choi, K., & Kim, Y.B., 2010. Molecular mechanism of insulin resistance in obesity and type 2 diabetes. *Korean J. Intern. Med.* 25: 119–129. doi:10.3904/kjim.2010.25.2.119
- Chopra, I., Li, H.F., Wang, H., & Webster, K.A., 2012. Phosphorylation of the insulin receptor by AMP-activated protein kinase (AMPK) promotes ligand-independent activation of the insulin signalling pathway in rodent muscle. *Diabetologia* 55: 783–794. doi:10.1007/s00125-011-2407-y
- Chua, B.T., Gallego-Ortega, D., de Molina, A.R., Ullrich, A., Lacal, J.C., & Downward, J., 2009. Regulation of Akt(ser473) phosphorylation by choline kinase in breast carcinoma cells. *Mol. Cancer* 8: 1–12. doi:10.1186/1476-4598-8-131
- Colhoun, H.M., Livingstone, S.J., Looker, H.C., Morris, A.D., Wild, S.H., Lindsay, R.S., *et al.*, 2012. Hospitalised hip fracture risk with rosiglitazone and pioglitazone use compared with other glucose-Lowering drugs. *Diabetologia* 55: 2929–2937. doi:10.1007/s00125-012-2668-0
- Collino, M., Aragno, M., Castiglia, S., Tomasinelli, C., Thiemermann, C., Boccuzzi, G., *et al.*, 2009. Insulin reduces cerebral ischemia/reperfusion injury in the hippocampus of diabetic rats: A role for glycogen synthase kinase-3 $\beta$ . *Diabetes* 58: 235–242. doi:10.2337/db08-0691
- Cook, M.N., Girman, C.J., Stein, P.P., & Alexander, C.M., 2007. Initial monotherapy with either metformin or sulphonylureas often fails to achieve or maintain current glycaemic



- goals in patients with type 2 diabetes in UK primary care. *Diabet. Med.* 24: 350-358. doi:10.1111/j.1464-5491.2007.02078.x
- Cusi, K., Consoli, A., & DeFronzo, R.A., 1996. Metabolic effects of metformin on glucose and lactate metabolism in noninsulin-dependent diabetes mellitus. *Endocrinol. Metab.* 81: 4059-4067.
- Daisy, P., Jasmine, R., Ignacimuthu, S., & Murugan, E., 2009. A novel steroid from *Elephantopus scaber* L. an ethnomedicinal plant with antidiabetic activity. *Phytomedicine* 16: 252–257. doi:10.1016/j.phymed.2008.06.001
- De Abreu Costa, L., Ottoni, M.H.F., Dos Santos, M.G., Meireles, A.B., De Almeida, V.G., De Fátima Pereira, W., *et al.*, 2017. Dimethyl sulfoxide (DMSO) decreases cell proliferation and TNF- $\alpha$ , IFN- $\gamma$ , and IL-2 cytokines production in cultures of peripheral blood lymphocytes. *Molecules* 22: 1–10. doi:10.3390/molecules22111789
- De, D., Chatterjee, K., Ali, K.M., Bera, T.K., & Ghosh, D., 2011. Antidiabetic potentiality of the aqueous-methanolic extract of seed of *Swietenia mahagoni* (L.) Jacq. in streptozotocin-induced diabetic male albino rat: A correlative and evidence-based approach with antioxidative and antihyperlipidemic activities. *Evidence-based Complement. Altern. Med.* 2011: 1-11. doi:10.1155/2011/892807
- DeFronzo, R.A., 2009. From the triumvirate to the ominous octet: A new paradigm for the treatment of type 2 diabetes mellitus. *Diabetes* 58: 773–795. doi:10.2337/db09-9028
- Den Hartogh, D.J., Vlaveciski, F., Giacca, A., & Tsiani, E., 2020. Attenuation of free fatty acid (FFA)-induced skeletal muscle cell insulin resistance by resveratrol is linked to activation of AMPK and inhibition of mTOR and p70 S6K. *Int. J. Mol. Sci.* 21: 1–17. doi:10.3390/ijms21144900
- Denes, L.T., Riley, L.A., Mijares, J.R., Arboleda, J.D., McKee, K., Esser, K.A., *et al.*, 2019. Culturing C2C12 myotubes on micromolded gelatin hydrogels accelerates myotube maturation. *Skelet. Muscle* 9: 1–10. doi:10.1186/s13395-019-0203-4
- Departemen Kesehatan RI, 2008. Pedoman Pengendalian Diabetes Melitus dan Penyakit Metabolik. *Indones. Sehat 2010*.
- Deshpande, V., Zen, Y., Chan, J.K.C., Yi, E.E., Sato, Y., Yoshino, T., *et al.*, 2012. Consensus statement on the pathology of IgG4-related disease. *Mod. Pathol.* 25: 1181–1192. doi:10.1038/modpathol.2012.72
- Dewanjee, S., Maiti, A., Das, A.K., Mandal, S.C., & Dey, S.P., 2009. Swietenine: A potential oral hypoglycemic from *Swietenia macrophylla* seed. *Fitoterapia* 80: 249–251.
- Dimopoulos, N., Watson, M., Sakamoto, K., & Hundal, H.S., 2006. Differential effects of palmitate and palmitoleate on insulin action and glucose utilization in rat L6 skeletal muscle cells. *Biochem. J.* 399: 473–481. doi:10.1042/BJ20060244
- Ding, X., Jian, T., Wu, Y., Zuo, Y., Li, J., Lv, H., *et al.*, 2019. *Ellagic acid* ameliorates oxidative stress and insulin resistance in high glucose-treated HepG2 cells via miR-223/keap1-Nrf2 pathway. *Biomed. Pharmacother.* 110: 85–94. doi:10.1016/j.biopha.2018.11.018
- DiNicolantonio, J.J., Bhutani, J., & O’Keefe, J.H., 2015. Acarbose: safe and effective for lowering postprandial hyperglycaemia and improving cardiovascular outcomes. *Open Hear.* 2: 1–13. doi:10.1136/openhrt-2015-000327
- Draznin, B., 2006. Molecular mechanisms of insulin resistance: Serine phosphorylation of insulin receptor substrate-1 and increased expression of p85  $\alpha$ : The two sides of a coin. *Diabetes* 55: 2392–2397. doi:10.2337/db06-0391
- Durai, M. V., Balamuniappan, G., & Geetha, S., 2016. Phytochemical screening and antimicrobial activity of leaf, seed and central-fruit-axis crude extract of *Swietenia macrophylla* King. *J. Pharmacogn. Phytochem.* 5: 181–186.
- Dutta, M., Biswas, U., Chakraborty, R., Banerjee, P., & Raychaudhuri, U., 2012.

- Regeneration of pancreatic  $\beta$ -cells on streptozotocin induced diabetic rats under the effect of *Swietenia macrophylla* seeds. *Int. J. Green Pharm.* 6: 336–339. doi:10.4103/0973-8258.108253
- ECACC, 2018. Cell line profile C2C12 catalogue no.910031101.
- Eid, A.M.M., Elmarzugi, N.A., & El-Enshasy, H.A., 2013. A review on the phytopharmacological effect of *Swietenia macrophylla*. *Int. J. Pharm. Pharm. Sci.* 5: 47–53.
- Elekofehinti, O.O., 2015. Saponins: Anti-diabetic principles from medicinal plants - A review. *Pathophysiology* 22: 95–103. doi:10.1016/j.pathophys.2015.02.001
- Esposito, D.L., Li, Y., Cama, A., & Quon, M.J., 2001. Tyr612 and Tyr632 in human insulin receptor substrate-1 are important for full activation of insulin-stimulated phosphatidylinositol 3-kinase activity and translocation of GLUT4 in adipose cells. *Endocrinology* 142: 2833–2840. doi:10.1210/endo.142.7.8283
- Faizal, A., & Geelen, D., 2013. Saponins and their role in biological processes in plants. *Phytochem. Rev.* 12: 877–893. doi:10.1007/s11101-013-9322-4
- Fan, X., Tao, J., Zhou, Y., Hou, Y., Wang, Y., Gu, D., *et al.*, 2019. Investigations on the effects of ginsenoside-Rg1 on glucose uptake and metabolism in insulin resistant HepG2 cells. *Eur. J. Pharmacol.* 843: 277–284. doi:10.1016/j.ejphar.2018.11.024
- Feng, M., Liu, F., Xing, J., Zhong, Y., & Zhou, X., 2021. *Anemarrhena saponins* attenuate insulin resistance in rats with high-fat diet-induced obesity via the IRS1/PI3K/AKT pathway. *J. Ethnopharmacol.* 277: 114251. doi:10.1016/j.jep.2021.114251
- Feng, X.T., Wang, T.Z., Leng, J., Chen, Y., Liu, J.B., Liu, Y., *et al.*, 2012. Palmitate contributes to insulin resistance through downregulation of the src-mediated phosphorylation of Akt in C2C12 myotubes. *Biosci. Biotechnol. Biochem.* 76: 1356–1361. doi:10.1271/bbb.120107
- Foretz, M., Hébrard, S., Leclerc, J., Zarrinpashneh, E., Soty, M., Mithieux, G., *et al.*, 2010. Metformin inhibits hepatic gluconeogenesis in mice independently of the LKB1/AMPK pathway via a decrease in hepatic energy state. *J. Clin. Invest.* 120: 2355–2369. doi:10.1172/JCI40671
- Fujimoto, W.Y., 2000. The importance of insulin resistance in the pathogenesis of type 2 diabetes mellitus. *Am. J. Med.* 108: 9–14. doi:10.1016/s0002-9343(00)00337-5
- Gao, D., Nong, S., Huang, X., Lu, Y., Zhao, H., Lin, Y., *et al.*, 2010. The effects of palmitate on hepatic insulin resistance are mediated by NADPH oxidase 3-derived reactive oxygen species through JNK and p38 MAPK pathways. *J. Biol. Chem.* 285: 29965–29973. doi:10.1074/jbc.M110.128694
- Gentile, M.T., Vecchione, C., Marino, G., Aretini, A., Pardo, A. Di, Antenucci, G., *et al.*, 2008. Resistin impairs insulin-evoked vasodilation. *Diabetes* 57: 577–583. doi:10.2337/db07-0557
- Gerich, J.E., 2003. Contributions of insulin-resistance and insulin-secretory defects to the pathogenesis of type 2 diabetes mellitus. *Mayo Clin. Proc.* 78: 447–456. doi:10.4065/78.4.447
- Gibbs, P.E.M., Miralem, T., Lerner-Marmarosh, N., & Maines, M.D., 2016. Nanoparticle delivered human biliverdin reductase-based peptide increases glucose uptake by activating IRK/Akt/GSK3 Axis: The peptide is effective in the cell and wild-type and diabetic ob/b Mice. *J. Diabetes Res.* 1–15. doi:10.1155/2016/4712053
- Goedeke, L., Perry, R.J., & Shulman, G.I., 2019. Emerging pharmacological targets for the treatment of nonalcoholic fatty liver disease, insulin resistance, and type 2 diabetes. *Annu Rev Pharmacol Toxicol.* 59: 65–87. doi:10.1146/annurev-pharmtox-010716-104727.
- Goh, B.H., & Kadir, H.A., 2011. In vitro cytotoxic potential of *Swietenia macrophylla* King

- Gonzalez, E., & McGraw, T.E., 2009. Insulin-modulated Akt subcellular localization determines Akt isoform-specific signaling. *Proc. Natl. Acad. Sci. U. S. A.* 106: 7004–7009. doi:10.1073/pnas.0901933106
- Guo, S., 2014. Insulin signaling, resistance, and metabolic syndrome: insights from mouse models into disease mechanisms. *J. Endocrinol.* 220: T1–T23. doi:10.1530/JOE-13-0327
- Guo, Xuan, Sun, W., Luo, G., Wu, L., Xu, G., Hou, D., *et al.*, 2019. *Panax notoginseng saponins* alleviate skeletal muscle insulin resistance by regulating the IRS1–PI3K–AKT signaling pathway and GLUT4 expression. *FEBS Open Bio* 9: 1008–1019. doi:10.1002/2211-5463.12635
- Han, M.S., Lim, Y.M., Quan, W., Kim, J.R., Chung, K.W., Kang, M., *et al.*, 2011. Lysophosphatidylcholine as an effector of fatty acid-induced insulin resistance. *J. Lipid Res.* 52: 1234–1246. doi:10.1194/jlr.M014787
- Hasan, M.M., Ahmed, Q.U., Soad, S.Z.M., Latip, J., Taher, M., Syafiq, T.M.F., *et al.*, 2017. Flavonoids from *Tetracera indica* Merr. induce adipogenesis and exert glucose uptake activities in 3T3-L1 adipocyte cells. *BMC Complement. Altern. Med.* 17: 1–14. doi:10.1186/s12906-017-1929-3
- Hashim, M.A., Yam, M.F., Hor, S.Y., Lim, C.P., Asmawi, M.Z., & Sadikun, A., 2013. Anti-hyperglycaemic activity of *Swietenia macrophylla* King (meliaceae) seed extracts in normoglycaemic rats undergoing glucose tolerance tests. *Chin. Med.* 8: 1–8. doi:10.1186/1749-8546-8-11
- Hassan, R.H., De Sousa, A.C.P., Mahfouz, R., Hainault, I., Blachnio-Zabielska, A., Bourron, O., *et al.*, 2016. Sustained action of ceramide on the insulin signaling pathway in muscle cells: Implication of the double-stranded RNA-activated protein kinase. *J. Biol. Chem.* 291: 3019–3029. doi:10.1074/jbc.M115.686949
- He, L., & Wondisford, F.E., 2015. Metformin action: Concentrations matter. *Cell Metab.* 21: 159–162. doi:10.1016/j.cmet.2015.01.003
- Henao-Duque, A.M., Rodriguez, B. de J., Pereanez, J.A., Lobo-Echeverri, T., & Nunez Rangel, V., 2017. Acute oral toxicity from a fraction rich in phenolic compounds from the leaf extract of *Swietenia macrophylla* King in a murine model. *Rev. Vitae* 24: 23–29. doi:10.17533/udea.vitae.v24n1a03
- Henriksen, E.J., & Dokken, B.B., 2006. Role of glycogen synthase kinase-3 in insulin resistance and type 2 diabetes. *Curr. Drug Targets* 7: 1435–1441. doi:10.2174/1389450110607011435
- Hernandez, R., Teruel, T., & Lorenzo, M., 2001. Akt mediates insulin induction of glucose uptake and up-regulation of GLUT4 gene expression in brown adipocytes. *FEBS Lett.* 494: 225–231. doi:10.1016/S0014-5793(01)02353-5
- Hoffmeister, L., Diekmann, M., Brand, K., & Huber, R., 2020. GSK3: A kinase balancing promotion and resolution of inflammation. *Cells* 9(4): 1-34 doi:10.3390/cells9040820
- Hu, X., Wang, S., Xu, J., Wang, D.B., Chen, Y., & Yang, G.Z., 2014. Triterpenoid saponins from *Stauntonia chinensis* ameliorate insulin resistance via the AMP-activated protein kinase and IR/IRS1/PI3K/Akt pathways in insulin-resistant HepG2 cells. *Int. J. Mol. Sci.* 15: 10446–10458. doi:10.3390/ijms150610446
- Huang, S., & Czech, M.P., 2007. The GLUT4 glucose transporter. *Cell Metab.* 5: 237–252. doi:10.1016/j.cmet.2007.03.006
- Huang, X., Liu, G., Guo, J., & Su, Z.Q., 2018. The PI3K/AKT pathway in obesity and type 2 diabetes. *Int. J. Biol. Sci.* 14: 1483–1496. doi:10.7150/ijbs.27173
- Huang, Y., Huang, T., Zhen, X., Li, Y., Mo, M., Ye, D., *et al.*, 2019. A selective sphingomyelin synthase 2 inhibitor ameliorates diet induced insulin resistance via the

IRS1/Akt/GSK-3 $\beta$  signaling pathway. *Pharmazie* 74: 553–558.  
doi:10.1691/ph.2019.9310

- Iliya, I., Mohammed, B., Akuyam, S., Nok, A., Bauchi, Z., Tanko, M., *et al.*, 2016. Molecular mechanisms of insulin signaling. *Sub-Saharan African J. Med.* 3: 1–7. doi:10.4103/2384-5147.176267
- International Diabetes Federation (IDF), 2017. IDF Diabetes Atlas, 8th edition, IDF Diabetes Atlas, 8th edition.
- Inzucchi, S.E., Bergenstal, R.M., Buse, J.B., Diamant, M., Ferrannini, E., Nauck, M., *et al.*, 2015. Management of hyperglycemia in type 2 diabetes, 2015: A patient-centered approach: Update to a position statement of the American Diabetes Association and the European Association for the study of diabetes. *Diabetes Care* 38: 140–149. doi:10.2337/dc14-2441
- Ishii, M., Maeda, A., Tani, S., & Akagawa, M., 2015. Palmitate induces insulin resistance in human HepG2 hepatocytes by enhancing ubiquitination and proteasomal degradation of key insulin signaling molecules. *Arch. Biochem. Biophys.* 566: 26–35. doi:10.1016/j.abb.2014.12.009
- Jamalzadeh, L., Ghafoori, H., Sariri, R., Rabuti, H., Nasirzade, J., Hasani, H., *et al.*, 2016. Cytotoxic effects of some common organic solvents on MCF-7, RAW-264.7 and human umbilical vein endothelial cells. *Avicenna J. Med. Biochem* 4(1): 1-6. doi:10.17795/ajmb-33453
- Jessen, N., & Goodyear, L.J., 2005. Invited review: Contraction signaling to glucose transport in skeletal muscle. *J. Appl. Physiol.* 99: 330–337. doi:10.1152/jappphysiol.00175.2005
- Ji, Y., Ji, C., Yue, L., & Xu, H., 2012. Saponins isolated from Asparagus induce apoptosis in human hepatoma cell line HepG2 through a mitochondrial-mediated pathway. *Curr. Oncol.* 19: 1–9. doi:10.3747/co.19.1139
- Jiang, S., Xu, L., Xu, Y., Guo, Y., Wei, L., Li, X., *et al.*, 2020. Antidiabetic effect of *Momordica charantia* saponins in rats induced by high-fat diet combined with STZ. *Electron. J. Biotechnol.* 43: 41–47. doi:10.1016/j.ejbt.2019.12.001
- Jové, M., Planavila, A., Laguna, J.C., & Vázquez-Carrera, M., 2005. Palmitate-induced interleukin 6 production is mediated by protein kinase C and nuclear-factor  $\beta$  activation and leads to glucose transporter 4 down-regulation in skeletal muscle cells. *Endocrinology* 146: 3087–3095. doi:10.1210/en.2004-1560
- Jové, M., Planavila, A., Sánchez, R.M., Merlos, M., Laguna, J.C., & Vázquez-Carrera, M., 2006. Palmitate induces tumor necrosis factor- $\alpha$  expression in C2C12 skeletal muscle cells by a mechanism involving protein kinase C and nuclear factor- $\kappa\beta$  activation. *Endocrinology* 147: 552–561. doi:10.1210/en.2005-0440
- Jung, T.W., Kim, H.C., Abd El-Aty, A.M., & Jeong, J.H., 2017. Protectin DX ameliorates palmitate-or high-fat diet-induced insulin resistance and inflammation through an AMPK-PPAR $\alpha$ -dependent pathway in mice. *Sci. Rep.* 7: 1–9. doi:10.1038/s41598-017-01603-9
- Kaku, K., 2010. Pathophysiology of type 2 diabetes and its treatment policy. *Japan Med. Assoc. J.* 53: 41–46.
- Kalaivanan, K., & Pugalendi, K.V., 2011. Antihyperglycemic effect of the alcoholic seed extract of *Swietenia macrophylla* on streptozotocin-diabetic rats. *Pharmacognosy Res.* 3: 67–71. doi:10.4103/0974-8490.79119
- Kammerer, S., & Küpper, J.-H., 2018. Human hepatocyte systems for in vitro toxicology analysis. *J. Cell. Biotechnol.* 3: 85–93. doi:10.3233/jcb-179012
- Kang, J.S., Kim, S.O., Kim, G.Y., Hwang, H.J., Kim, B.W., Chang, Y.C., *et al.*, 2016. An exploration of the antioxidant effects of garlic saponins in mouse-derived C2C12



- Karnieli, E., & Armoni, M., 2008. Transcriptional regulation of the insulin-responsive glucose transporter GLUT4 gene: From physiology to pathology. *Am. J. Physiol. - Endocrinol. Metab.* 295: 38–45. doi:10.1152/ajpendo.90306.2008
- Karthikeyan, G., Lakshmi, T., Rajeshkumar, S., Roy, A., Gurunadhan, D., Geetha, R. V., *et al.*, 2019. Glucose uptake potential in L6 myotubes by ficus racemosa. *Indian J. Public Heal. Res. Dev.* 10: 3527–3531. doi:10.5958/0976-5506.2019.04132.9
- Kataoka, Y., Matsumura, I., Ezoe, S., Nakata, S., Takigawa, E., Sato, Y., *et al.*, 2003. Reciprocal inhibition between MyoD and STAT3 in the regulation of growth and differentiation of myoblasts. *J. Biol. Chem.* 278: 44178–44187. doi:10.1074/jbc.M304884200
- Kaur, P., Robin, Mehta, R.G., Arora, S., & Singh, B., 2018. Progression of conventional hepatic cell culture models to bioengineered HepG2 cells for evaluation of herbal bioactivities. *Biotechnol. Lett.* 40: 881–893. doi:10.1007/s10529-018-2547-y
- Kazyken, D., Magnuson, B., Bodur, C., Acosta-jaquez, H.A., Tong, X., Barnes, T.M., *et al.*, 2019. AMPK directly activates MTOR2 to promote cell survival during acute energetic stress. *Sci Signal* 12: 1–37. doi:10.1126/scisignal.aav3249.AMPK
- Keller, A.C., Ma, J., Kavalier, A., He, K., Brillantes, A.-M.B., & Kennelly, E.J., 2011. Saponins from the traditional medicinal plant *Momordica charantia* stimulate insulin secretion in vitro. *Phytomedicine* 19: 32–37. doi:<https://dx.doi.org/10.1016%2Fj.phymed.2011.06.019>
- Khalid, M., Alkaabi, J., Khan, M.A.B., & Adem, A., 2021. Insulin signal transduction perturbations in insulin resistance. *Int. J. Mol. Sci.* 22: 1–17. doi:10.3390/ijms22168590
- Kim, Do Sung, Jeong, S.K., Kim, H.R., Kim, Dal Sik, Chae, S.W., & Chae, H.J., 2010. Metformin regulates palmitate-induced apoptosis and ER stress response in HepG2 liver cells. *Immunopharmacol. Immunotoxicol.* 32: 251–257. doi:10.3109/08923970903252220
- Kim, M.H., Jee, J.H., Park, S., Lee, M.S., Kim, K.W., & Lee, M.K., 2014. Metformin enhances glucagon-like peptide 1 via cooperation between insulin and Wnt signaling. *J. Endocrinol.* 220: 117–128. doi:10.1530/JOE-13-0381
- Kim, Y.D., Park, K., Lee, Y., Park, Y., Kim, D., Nedumaran, B., *et al.*, 2008. Metformin inhibits hepatic gluconeogenesis through AMP-activated protein kinase – dependent regulation of the orphan nuclear receptor SHP. *Diabetes* 57: 306–14. doi:10.2337/db07-0381.
- Krishna, R.N., Anitha, R., & Ezhilarasan, D., 2020. Aqueous extract of *Tamarindus indica* fruit pulp exhibits antihyperglycaemic activity. *Avicenna J. phytomedicine* 10: 440–447. doi:10.22038/ajp.2020.14787
- Krisnawati, H., Varis, E., Kallio, M., & Markku, K., 2011. *Swietenia macrophylla* King.: Ecology, silviculture and productivity. *Cent. Int. For. Res.* 1–12.
- Krook, A., Wallberg-Henriksson, H., & Zierath, J.R., 2004. Sending the signal: Molecular mechanisms regulating glucose uptake. *Med. Sci. Sports Exerc.* 36: 1212–1217. doi:10.1249/01.MSS.0000132387.25853.3B
- Kumar, K., Ghosh, S., Canovatchel, W., Garodia, N., & Rajashekar, S., 2017. A review of clinical efficacy and safety of canagliflozin 300 mg in the management of patients with type 2 diabetes mellitus. *Indian J. Endocrinol. Metab.* 21: 196–209. doi:10.4103/2230-8210.196016
- Kumar, N., & Dey, C.S., 2002. Metformin enhances insulin signalling in insulin-dependent and -independent pathways in insulin resistant muscle cells. *Br. J. Pharmacol.* 137: 329–336. doi:10.1038/sj.bjp.0704878
- Kusumastuti, S.A., Nugrahaningsih, D.A.A., & Wahyuningsih, M.S.H., 2019. *Centella*

- asiatica* (L.) extract attenuates inflammation and improve insulin sensitivity in a coculture of lipopolysaccharide (LPS)-induced 3T3-L1 adipocytes and RAW 264.7 macrophages. *Drug Discov. Ther.* 13: 261–267. doi:10.5582/ddt.2019.01052
- Lamoia, T.E., & Shulman, G.I., 2021. Cellular and molecular mechanisms of metformin action. *Endocr. Rev.* 42: 77–96. doi:10.1210/edrv/bnaa023
- Lau, W.K., Goh, B.H., Kadir, H.A., Shu-Chien, A.C., Muhammad, T.S.T., & McPhee, D.J., 2015. Potent PPAR $\gamma$  ligands from *Swietenia macrophylla* are capable of stimulating glucose uptake in muscle cells. *Molecules* 20: 22301–22314. doi:10.3390/molecules201219847
- Leclercq, I.A., Da Silva Morais, A., Schroyen, B., Van Hul, N., & Geerts, A., 2007. Insulin resistance in hepatocytes and sinusoidal liver cells: Mechanisms and consequences. *J. Hepatol.* 47: 142–156. doi:10.1016/j.jhep.2007.04.002
- Lee, H.M., Lee, O.H., Kim, K.J., & Lee, B.Y., 2012. Ginsenoside Rg1 promotes glucose uptake through activated AMPK pathway in insulin-resistant muscle cells. *Phyther. Res.* 26: 1017–1022. doi:10.1002/ptr.3686
- Lee, J. young, Cho, H.K., & Kwon, Y.H., 2010. Palmitate induces insulin resistance without significant intracellular triglyceride accumulation in HepG2 cells. *Metabolism.* 59: 927–934. doi:10.1016/j.metabol.2009.10.012
- Lee, J.S., Park, J.M., Lee, S., Lee, H.J., Yang, H.S., Yeo, J., *et al.*, 2017. Hispidin rescues palmitate-induced insulin resistance in C2C12 myotubes. *Mol. Med. Rep.* 16: 4229–4234. doi:10.3892/mmr.2017.7042
- Leng, S., Zhang, W., Zheng, Y., Liberman, Z., Rhodes, C.J., Finkelman, H.E., *et al.*, 2010. Glycogen synthase kinase 3 $\beta$  mediates high glucose-induced ubiquitination and proteasome degradation of insulin receptor substrate 1. *J. Endocrinology* 206: 171–181. doi:10.1677/JOE-09-0456.Glycogen
- Li, H.B., Yang, Y.R.Y., Mo, Z.J., Ding, Y., & Jiang, W.J., 2015. Silibinin improves palmitate-induced insulin resistance in C2C12 myotubes by attenuating IRS1/PI3K/Akt pathway inhibition. *Brazilian J. Med. Biol. Res.* 48: 440–446. doi:10.1590/1414-431X20144238
- Li, Z., Lai, Z.W., Christiano, R., Gazos-Lopes, F., Walther, T.C., & Farese, R. V., 2018. Global analyses of selective insulin resistance in hepatocytes caused by palmitate lipotoxicity. *Mol. Cell. Proteomics* 17: 836–849. doi:10.1074/mcp.RA117.000560
- Liang, G., Wang, F., Song, X., Zhang, L., Qian, Z., & Jiang, G., 2016. 3-Deoxyglucosone induces insulin resistance by impairing insulin signaling in HepG2 cells. *Mol. Med. Rep.* 13: 4506–4512. doi:10.3892/mmr.2016.5081
- Lin, C.H., Kuo, Y.H., & Shih, C.C., 2018. Antidiabetic and hypolipidemic activities of eburicoic acid, a triterpenoid compound from *Antrodia camphorata* by regulation of Akt phosphorylation, gluconeogenesis, and PPAR $\alpha$  in streptozotocin-induced diabetic mice. *RSC Adv.* 8: 20462–20476. doi:10.1039/c8ra01841c
- Lin, Y., & Sun, Z., 2010. Current views on type 2 diabetes. *J. Endocrinology* 204: 1–18. doi:10.1677/JOE-09-0260
- Lipina, C., Huang, X., Finlay, D., McManus, E.J., Alessi, D.R., & Sutherland, C., 2005. Analysis of hepatic gene transcription in mice expressing insulin-insensitive GSK3. *Biochem. J.* 392: 633–639. doi:10.1042/BJ20051046
- Liu, J.Y., Zhang, Y.C., Song, L.N., Zhang, L., Yang, F.Y., Zhu, X.R., *et al.*, 2019a. Nifuroxazide ameliorates lipid and glucose metabolism in palmitate-induced HepG2 cells. *RSC Adv.* 9: 39394–39404. doi:10.1039/c9ra06527j
- Liu, Q., Zhang, F.G., Zhang, W.S., Pan, A., Yang, Y.L., Liu, J.F., *et al.*, 2017. Ginsenoside Rg1 inhibits glucagon-induced hepatic gluconeogenesis through Akt-Foxo1 interaction. *Theranostics* 7: 4001–4012. doi:10.7150/thno.18788



- Liu, T.Y., Shi, C.X., Gao, R., Sun, H.J., Xiong, X.Q., Ding, L., *et al.*, 2015. Irisin inhibits hepatic gluconeogenesis and increases glycogen synthesis via the PI3K/Akt pathway in type 2 diabetic mice and hepatocytes. *Clin. Sci.* 129: 839–850. doi:10.1042/CS20150009
- Liu, X., 2019. Research progresses of the mechanism of insulin resistance in type II diabetes. *E3S Web Conf.* 78: 1–6. doi:10.1051/e3sconf/20197801006
- Liu, Y., Deng, J., & Fan, D., 2019b. Ginsenoside Rk3 ameliorates high-fat-diet/streptozocin induced type 2 diabetes mellitus in mice: Via the AMPK/Akt signaling pathway. *Food Funct.* 10: 2538–2551. doi:10.1039/c9fo00095j
- Liu, Z., Qu, C.Y., Li, J.X., Wang, Y.F., Li, W., Wang, C.Z., *et al.*, 2021. Hypoglycemic and hypolipidemic effects of malonyl ginsenosides from American ginseng (*Panax quinquefolius* L.) on type 2 diabetic mice. *ACS Omega* 6: 33652–33664. doi:10.1021/acsomega.1c04656
- Lochhead, P.A., Coghlan, M., Rice, S.Q.J., & Sutherland, C., 2001. Inhibition of GSK-3 selectively reduces glucose-6-phosphatase and phosphoenolpyruvate carboxykinase gene expression. *Diabetes* 50: 937–946.
- Lu, J.M., Wang, Y.F., Yan, H.L., Lin, P., Gu, W., & Yu, J., 2016. Antidiabetic effect of total saponins from *Polygonatum kingianum* in streptozotocin-induced diabetic rats. *J. Ethnopharmacol.* 179: 291–300. doi:10.1016/j.jep.2015.12.057
- Luo, W., Ai, L., Wang, B. fa, & Zhou, Y., 2019. High glucose inhibits myogenesis and induces insulin resistance by down-regulating AKT signaling. *Biomed. Pharmacother.* 120: 109498. doi:10.1016/j.biopha.2019.109498
- MacAulay, K., & Woodgett, J.R., 2008. Targeting glycogen synthase kinase-3 (GSK-3) in the treatment of type 2 diabetes. *Expert Opin. Ther. Targets* 12: 1265–1274. doi:10.1517/14728222.12.10.1265
- Mackenzie, R.W.A., & Elliott, B.T., 2014. Akt/PKB activation and insulin signaling: A novel insulin signaling pathway in the treatment of type 2 diabetes. *Diabetes, Metab. Syndr. Obes. Targets Ther.* 7: 55–64. doi:10.2147/DMSO.S48260
- Maiti, A., Dewanjee, S., Jana, G., & Mandal, S., 2008. Hypoglycemic effect of *Swietenia macrophylla* seeds against type II diabetes. *Int. J. Green Pharm.* 2: 224–227. doi:10.4103/0973-8258.44738
- Maiti, A., Dewanjee, S., Kundu, M., & Mandal, S.C., 2009. Evaluation of antidiabetic activity of the seeds of *Swietenia macrophylla* in diabetic rats. *Pharm. Biol.* 47: 132–136. doi:10.1080/13880200802436703
- Maiti, A., Dewanjee, S., & Mandal, S., 2007a. Effect of *Swietenia macrophylla* seeds on normal and alloxan-induced diabetic rats. *Orient. Pharm. Exp. Med.* 7: 363–371. doi:10.3742/opem.2007.7.4.363
- Maiti, A., Dewanjee, S., Kundu, M., & Mandal, S., 2007b. Protective effect of methanol extract of *Swietenia macrophylla* seeds on oxidative stress associated with streptozotocin induced diabetic rats. *Nat. Prod. Sci.* 13(4): 295–299.
- Malik, S.A., Acharya, J.D., Mehendale, N.K., Kamat, S.S., & Ghaskadbi, S.S., 2019. *Pterostilbene* reverses palmitic acid mediated insulin resistance in HepG2 cells by reducing oxidative stress and triglyceride accumulation. *Free Radic. Res.* 53: 815–827. doi:10.1080/10715762.2019.1635252
- Mannucci, E., & Monami, M., 2017. Bone fractures with sodium glucose co-transporter 2 inhibitors: How real is the risk? *Drug Saf.* 40: 115–119. doi:10.1007/s40264-016-0470-5
- Mejia, C.F., 2006. Molecular basis of type-2 diabetes. *Mol. Endocrinol.* 87–108.
- Meshkani, R., Sadeghi, A., Taheripak, G., Zarghooni, M., Gerayesh-Nejad, S., & Bakhtiyari, S., 2014. Rosiglitazone, a PPAR $\gamma$  agonist, ameliorates palmitate-induced insulin resistance and apoptosis in skeletal muscle cells. *Cell Biochem. Funct.* 32: 683–691. doi:10.1002/cbf.3072



- Miklosz, A., Łukaszuk, B., Baranowski, M., Górski, J., & Chabowski, A., 2013. Effects of inhibition of serine palmitoyltransferase (SPT) and sphingosine kinase 1 (SphK1) on palmitate induced insulin resistance in L6 myotubes. *PLoS One* 8: 1–9. doi:10.1371/journal.pone.0085547
- Moghadamtousi, S.Z., Goh, B.H., Chan, C.K., Shabab, T., & Kadir, H.A., 2013. Biological activities and phytochemicals of *Swietenia macrophylla* King. *Molecules* 18: 10465–10483. doi:10.3390/molecules180910465
- Mohanam, P., Subramaniyam, S., Mathiyalagan, R., & Yang, D.C., 2018. Molecular signaling of ginsenosides Rb1, Rg1, and Rg3 and their mode of actions. *J. Ginseng Res.* 42: 123–132. doi:10.1016/j.jgr.2017.01.008
- Molinaro, A., Becattini, B., & Solinas, G., 2020. Insulin signaling and glucose metabolism in different hepatoma cell lines deviate from hepatocyte physiology toward a convergent aberrant phenotype. *Sci. Rep.* 10: 1–10. doi:10.1038/s41598-020-68721-9
- Monroy, M.L.L. de la V., & Mejia, C.F., 2013. Oxidative stress in diabetes mellitus and the role of vitamins with antioxidant actions in oxidative stress and chronic degenerative diseases - A role for antioxidants. *IntechOpen*, pp. 209–232. doi:10.5772/51788
- Montano, M., 2014. C2C12 cell line: Model systems in translational biology in medicine. doi:10.1016/C2013-0-18173-1
- Mora, S., & Pessin, J.E., 2000. The MEF2A isoform is required for striated muscle-specific expression of the insulin-responsive GLUT4 glucose transporter. *J. Biol. Chem.* 275: 16323–16328. doi:10.1074/jbc.M910259199
- Morón, E.B., Jiménez, Z.A., de Marañón, A.M., Iannantuoni, F., López, I.E., Domènech, S.L., *et al.*, 2019. Relationship between oxidative stress, ER stress, and inflammation in type 2 diabetes: The battle continues. *J. Clin. Med.* 8: 1–22. doi:10.3390/jcm8091385
- Mota, M., Banini, B.A., Cazanave, S.C., & Sanyal, A.J., 2016. Molecular mechanisms of lipotoxicity and glucotoxicity in nonalcoholic fatty liver disease. *Metabolism.* 65: 1049–1061. doi:10.1016/j.metabol.2016.02.014
- Murphy, S.M., Kiely, M., Jakeman, P.M., Kiely, P.A., & Carson, B.P., 2016. Optimization of an in vitro bioassay to monitor growth and formation of myotubes in real time. *Biosci. Rep.* 36: 1–10. doi:10.1042/BSR20160036
- Mursiti, S., 2009. Isolasi, karakterisasi, dan uji aktivitas hipoglikemik senyawa dalam biji mahoni bebas minyak dan minyak biji mahoni (*Swietenia macrophylla* King). Semarang.
- Mursiti, S., 2008. Isolation compound antidiabetes mellitus from the seeds of mahogany (*Swietenia macrophylla* King) (translate from Indonesia language) [doctoral's desertation]. Universitas Gadjah Mada, Yogyakarta.
- Muthmainah, 2019. Effect of seeds isolate from *Swietenia macrophylla* King in type 2 diabetic rats model: molecular study of antidiabetic effect of isolate 1,4-bis-(3,4,5-trimetoksi-fenil)-tetrahidro-furo(3,4-c) furan (translate from Indonesia language). Universitas Sebelas Maret, Surakarta.
- Muthmainah, M., Yarso, K.Y., Purwanto, B., Mudigdo, A., & Mustofa, M., 2019. 1,4-bis-3,4,5-trimethoxy-phenyl-tetrahydro-furo(3,4-C) furan from mahogany (*Swietenia Macrophylla* King) seed significantly reduces glucose and malondialdehyde levels in diabetic wistar rats. *Bali Med. J.* 8: 570–575. doi:10.15562/bmj.v8i2.1227
- Muthmainah, Nurwati, I., Handayani, S., Saptiwi, B., & Ma'rufah, S., 2021. Isolat biji mahoni (*Swietenia macrophylla* King) memperbaiki gambaran histopatologi hepar tikus model DM Tipe 2. *SMJ.* 4: 73–82. doi:10.13057/smj.v4i2.
- Negi, J.S., Negi, P.S., Pant, G.J., Rawat, M., & Negi, S.K., 2013. Naturally occurring saponins: Chemistry and biology. *J. Poisonous Med. Plant Res.* 1: 1–6.
- Newsholme, P., Haber, E.P., Hirabara, S.M., Rebelato, E.L.O., Procopio, J., Morgan, D., *et*



- al.*, 2007. Diabetes associated cell stress and dysfunction: Role of mitochondrial and non-mitochondrial ROS production and activity. *J. Physiol.* 583: 9–24. doi:10.1113/jphysiol.2007.135871
- Nikoulina, S.E., Ciaraldi, T.P., Mudaliar, S., Carter, L., Johnson, K., & Henry, R.R., 2002. Inhibition of glycogen synthase kinase 3 improves insulin action and glucose metabolism in human skeletal muscle. *Diabetes* 51: 2190–2198. doi:10.2337/diabetes.51.7.2190
- Nikoulina, S.E., Ciaraldi, T.P., Mudaliar, S., Mohideen, P., Carter, L., & Henry, R.R., 2000. Potential role of glycogen synthase kinase-3 in skeletal muscle insulin resistance of type 2 diabetes. *Diabetes* 49: 263–271. doi:10.2337/diabetes.49.2.263
- Nissar, A.U., Sharma, L., & Tasduq, S.A., 2015. Palmitic acid induced lipotoxicity is associated with altered lipid metabolism, enhanced CYP450 2E1 and intracellular calcium mediated ER stress in human hepatoma cells. *Toxicol. Res. (Camb)*. 4: 1344–1358. doi:10.1039/c5tx00101c
- Nolan, C.J., & Prentki, M., 2019. Insulin resistance and insulin hypersecretion in the metabolic syndrome and type 2 diabetes: Time for a conceptual framework shift. *Diabetes Vasc. Dis. Res.* 16: 118–127. doi:10.1177/1479164119827611
- Nordlie, R.C., & Foster, J.D., 1999. Regulation of Glucose Metabolism. *Annu. Rev. Nutr.* 19: 379–406. doi:10.1146/annurev.nutr.19.1.379
- Nugraha, A., 2012. Molecular docking and antihyperglycemic activity of active compounds which isolated from methanol extract of *Swietenia macrophylla* King seeds in diabetic rats induced by streptozotocin (translate from Indonesia language). Universitas Gadjah Mada, Yogyakarta.
- Oliveira, A.F., Cunha, D.A., Ladriere, L., Esteve, M.I., Bugliani, M., Marchetti, P., *et al.*, 2015. In vitro use of free fatty acids bound to albumin: A comparison of protocols. *Biotechniques* 58: 228–233.
- Oliveira, J.M., Rebuffat, S.A., Gasa, R., & Gomis, R., 2014. Targeting type 2 diabetes: lessons from a knockout model of insulin receptor substrate 2. *Can. J. Physiol. Pharmacol.* 92: 613–620. doi:10.1139/cjpp-2014-0114
- Olson, A.L., 2012. Regulation of GLUT4 and insulin-dependent glucose flux. *ISRN Mol. Biol.* 1–12. doi:10.5402/2012/856987
- Ouyang, J., Parakhia, R.A., & Ochs, R.S., 2011. Metformin activates AMP kinase through inhibition of AMP deaminase. *J. Biol. Chem.* 286: 1–11. doi:10.1074/jbc.M110.121806
- Ozougwu, J., Obimba, K., Belonwu, C., & Unakalamba, C., 2013. The pathogenesis and pathophysiology of type 1 and type 2 diabetes mellitus. *J. Physiol. Pathophysiol.* 4: 46–57. doi:10.5897/jpap2013.0001
- Pandey, A.K., Verma, G., Vig, S., Srivastava, S., Srivastava, A.K., & Datta, M., 2011. MiR-29a levels are elevated in the db/db mice liver and its overexpression leads to attenuation of insulin action on PEPCK gene expression in HepG2 cells. *Mol. Cell. Endocrinol.* 332: 125–133. doi:10.1016/j.mce.2010.10.004
- Pardo, V., González-Rodríguez, Á., Muntané, J., Kozma, S.C., & Valverde, Á.M., 2015. Role of hepatocyte S6K1 in palmitic acid-induced endoplasmic reticulum stress, lipotoxicity, insulin resistance and in oleic acid-induced protection. *Food Chem. Toxicol.* 80: 298–309. doi:10.1016/j.fct.2015.03.029
- Paritala, V., Chiruvella, K.K., Thammineni, C., Ghanta, R.G., & Mohammed, A., 2015. Phytochemicals and antimicrobial potentials of mahogany family. *Rev. Bras. Farmacogn.* 25: 61–83. doi:10.1016/j.bjp.2014.11.009
- Park, S.Y., Kim, M.H., Kwon, H.Y., Ahn, J.H., Lee, S.J., Lee, J.H., *et al.*, 2014. The stimulatory effect of essential fatty acids on glucose uptake involves both Akt and AMPK activation in C2C12 skeletal muscle cells. *Korean J. Physiol. Pharmacol.* 18:



- Pearce, N.J., Arch, J.R.S., Clapham, J.C., Coghlan, M.P., Corcoran, S.L., Lister, C.A., *et al.*, 2004. Development of glucose intolerance in male transgenic mice overexpressing human glycogen synthase kinase-3 $\beta$  on a muscle-specific promoter. *Metabolism*. 53: 1322–1330. doi:10.1016/j.metabol.2004.05.008
- Peiffer, S.C., Craig, D.L., & Biden, T.J., 1999. Ceramide generation is sufficient to account for the inhibition of the insulin-stimulated PKB pathway in C2C12 skeletal muscle cells pretreated with palmitate. *J. Biol. Chem.* 274(34): 24202-24210 doi:10.1074/jbc.274.34.24202
- PERKENI, 2015. Konsensus pengelolaan pencegahan diabetes melitus tipe 2 di Indonesia.
- Petersen, M.C., & Shulman, G.I., 2018. Mechanisms of insulin action and insulin resistance. *Physiol. Rev.* 98: 2133–2223. doi:10.1152/physrev.00063.2017
- Pinel, A., Rigaudière, J.P., Jouve, C., & Capel, F., 2018. Modulation of insulin resistance and the adipocyte-skeletal muscle cell cross-talk by LCn-3PUFA. *Int. J. Mol. Sci.* 19: 0–15. doi:10.3390/ijms19092778
- Plomgaard, P., Nielsen, A.R., Fischer, C.P., Mortensen, O.H., Broholm, C., Penkowa, M., *et al.*, 2007. Associations between insulin resistance and TNF- $\alpha$  in plasma, skeletal muscle and adipose tissue in humans with and without type 2 diabetes. *Diabetologia* 50: 2562–2571. doi:10.1007/s00125-007-0834-6
- Ponnusamy, R., Giriraj, K., AM, M.S., Thangaraj, P., Sathyanarayanan, S., Araújo, A.A. de S., *et al.*, 2022. Glycogen synthase kinase 3 $\beta$  inhibition and insulin-receptor binding enhancement of compounds isolated from wild leafy vegetable *Acalypha alnifolia*. *Phytomedicine Plus* 2: 100216. doi:10.1016/j.phyplu.2022.100216
- Prasetyastuti, Sunarti, sadewa, A.H., & Mustofa, 2016. Effect of 7-hydroxy-2-(4-hydroxy-3-methoxy-phenyl)-chroman-4-one (*Swietenia macrophylla* King. seed) on retinol binding protein-4 and phosphoenolpyruvate carboxykinase gene expression in type 2 diabetic rats. *Rom. J. Diabetes, Nutr. Metab. Dis.* 23: 255-265. doi: 10.1515/rjdnmd-2016-0030
- Purintrapiban, J., & Ratanachaiyavong, S., 2003. The effects of insulin and metformin on glucose uptake in L8 m yotubes. *Science Asia* 29: 341–346. doi:10.2306/scienceasia1513-1874.2003.29.341
- Putra, I., & Berawi, K., 2015. Empat pilar penatalaksanaan pasien diabetes mellitus tipe 2. *Majority* 4: 8–12.
- Qaid, M.M., & Abdelrahman, M.M., 2016. Role of insulin and other related hormones in energy metabolism - A review. *Cogent Food Agric.* 2. doi:10.1080/23311932.2016.1267691
- Qin, H., Liu, Y., Lu, N.A., Li, Y., & Sun, C.H., 2009. Cis-9 trans-11 -conjugated linoleic acid activates AMP-activated protein kinase in attenuation of insulin resistance in C2C12 myotubes. *J. Agric. Food Chem.* 57: 4452–4458. doi:10.1021/jf900534w
- Quinn, P.G., & Yeagley, D., 2005. Insulin regulation of PEPCK gene expression: A model for rapid and reversible modulation. *Curr. Drug Targets Immune, Endocr. Metab. Disord.* 5: 423–437. doi:10.2174/156800805774912962
- Rachek, L.I., Musiyenko, S.I., LeDoux, S.P., & Wilson, G.L., 2007. Palmitate induced mitochondrial deoxyribonucleic acid damage and apoptosis in L6 rat skeletal muscle cells. *Endocrinology* 148: 293–299. doi:10.1210/en.2006-0998
- Ragheb, R., & Medhat, A.M., 2011. Mechanisms of fatty acid-induced insulin resistance in muscle and liver. *J. Diabetes Metab.* 02: 1–5. doi:10.4172/2155-6156.1000127
- Rahman, A.K.M.S., Chowdhury, A.K.A., Ali, H.A., Raihan, S.Z., Ali, M.S., Nahar, L., *et al.*, 2009. Antibacterial activity of two limonoids from *Swietenia mahagoni* against multiple-drug-resistant (MDR) bacterial strains. *J. Nat. Med.* 63: 41–45. doi:10.1007/s11418-008-0287-3



- Ramasamy, S., Wahab, N., Zainal Abidin, N., Manickam, S., & Zakaria, Z., 2012. Growth inhibition of human gynecologic and colon cancer cells by *Phyllanthus watsonii* through apoptosis induction. *PLoS One* 7: 1–15. doi:10.1371/journal.pone.0034793
- Rao, X., Huang, X., Zhou, Z., & Lin, X., 2013. An improvement of the 2<sup>-</sup>(-delta delta CT) method for quantitative real-time polymerase chain reaction data analysis. *Biostat. Bioinforma. Biomath.* 3: 71–85.
- Rena, G., Pearson, E.R., & Sakamoto, K., 2013. Molecular mechanism of action of metformin: Old or new insights? *Diabetologia* 56: 1898–1906. doi:10.1007/s00125-013-2991-0
- Ricchi, M., Odoardi, M.R., Carulli, L., Anzivino, C., Ballestri, S., Pinetti, A., *et al.*, 2009. Differential effect of oleic and palmitic acid on lipid accumulation and apoptosis in cultured hepatocytes. *J. Gastroenterol. Hepatol.* 24: 830–840. doi:10.1111/j.1440-1746.2008.05733.x
- Rodelo, C.G., Guiberna, A.R., & Reyes, J.A.O., 2017. Cellular mechanisms of insulin action. *Gac. Med. Mex.* 153: 197–209.
- Röder, P. V., Wu, B., Liu, Y., & Han, W., 2016. Pancreatic regulation of glucose homeostasis. *Exp. Mol. Med.* 48: 1-19. doi:10.1038/emm.2016.6
- Rui, L., 2014. Energy metabolism in the liver. *Compr Physiol.* 4(1): 177-197. doi:10.1002/cphy.c130024.Energy
- Russo, B., Picconi, F., Malandrucchio, I., & Frontoni, S., 2019. Flavonoids and insulin-resistance: from molecular evidences to clinical trials. *Int. J. Mol. Sci.* 20. doi:10.3390/ijms20092061
- Sahgal, G., Ramanathan, S., Sasidharan, S., Mordi, M.N., Ismail, S., & Mansor, S.M., 2009. In vitro antioxidant and xanthine oxidase inhibitory activities of methanolic *Swietenia mahagoni* seed extracts. *Molecules* 14: 4476–4485. doi:10.3390/molecules14114476
- Saini, V., 2010. Molecular mechanisms of insulin resistance in type 2 diabetes mellitus. *World J. Diabetes* 1: 68–75. doi:10.4239/wjd.v1.i3.68
- Salani, B., Del Rio, A., Marini, C., Sambucetti, G., Cordera, R., & Maggi, D., 2014. Metformin, cancer and glucose metabolism. *Endocr. Relat. Cancer* 21: R461–R471. doi:10.1530/ERC-14-0284
- Samuel, V.T., & Shulman, G.I., 2012. Mechanisms for insulin resistance: Common threads and missing links. *Cell* 148: 852–871. doi:10.1016/j.cell.2012.02.017
- Sanger, J.W., Wang, J., Holloway, B., Du, A., & Sanger, J.M., 2009. Myofibrillogenesis in skeletal muscle cells in zebrafish. *Cell Motil. Cytoskeleton* 66: 556–566. doi:10.1002/cm.20365
- Santoleri, D., & Titchenell, P.M., 2019. Resolving the paradox of hepatic insulin resistance. *CMGH* 7: 447–456. doi:10.1016/j.jcmgh.2018.10.016
- Santos, N.C., Figueira-Coelho, J., Martins-Silva, J., & Saldanha, C., 2003. Multidisciplinary utilization of dimethyl sulfoxide: Pharmacological, cellular, and molecular aspects. *Biochem. Pharmacol.* 65: 1035–1041. doi:10.1016/S0006-2952(03)00002-9
- Sari, S.N., 2016. Isolasi flavonoid dari biji mahoni (*Swietenia macrophylla* King.) dan uji aktivitasnya sebagai antibakteri. Universitas Negeri Semarang.
- Sariaydin, T., Çal, T., Aydın Dilsiz, S., Canpınar, H., & Ündeğer Bucurgat, Ü., 2021. In vitro assessment of cytotoxic, apoptotic and genotoxic effects of metformin. *Istanbul J. Pharm.* 51: 167–174. doi:10.26650/istanbuljpharm.2021.0079
- Sayyad, M., Tiang, N., Kumari, Y., Goh, B.H., Jaiswal, Y., Rosli, R., *et al.*, 2017. Acute toxicity profiling of the ethyl acetate fraction of *Swietenia macrophylla* seeds and in-vitro neuroprotection studies. *Saudi Pharm. J.* 25: 196–205. doi:10.1016/j.jsps.2016.05.002
- Scherthaner, G., Currie, C.J., & Scherthaner, G.H., 2013. Do we still need pioglitazone for

- the treatment of type 2 diabetes? A risk-benefit critique in 2013. *Diabetes Care* 36: 155–161. doi:10.2337/dcS13-2031
- Schinner, S., Scherbaum, W., Bornstein, S.R., & Barthel, A., 2005. Molecular Mechanisms of Insulin Resistance. *Diabet. Med.* 22: 674–682.
- Sefried, S., Häring, H.U., Weigert, C., & Eckstein, S.S., 2018. Suitability of hepatocyte cell lines HepG2, AML12 and THLE-2 for investigation of insulin signalling and hepatokine gene expression. *Open Biol.* 8. doi:10.1098/rsob.180147
- Shao, J., Qiao, L., Janssen, R.C., Pagliassotti, M., & Friedman, J.E., 2005. Chronic hyperglycemia enhances PEPCK gene expression and hepatocellular glucose production via elevated liver activating protein/liver inhibitory protein ratio. *Diabetes* 54: 976–984. doi:10.2337/diabetes.54.4.976
- Shaw, L.M., 2011. The insulin receptor substrate (IRS) proteins: At the intersection of metabolism and cancer. *Cell Cycle* 10: 1750–1756. doi:10.4161/cc.10.11.15824
- Shepherd, P.R., & Kahn, B.B., 1999. Glucose transporters and insulin action. *N. Engl. J. Med.* 341: 248–257. doi:10.1056/NEJM199907223410406
- Shi, J., & Kandrор, K. V., 2005. Sortilin is essential and sufficient for the formation of GLUT4 storage vesicles in 3T3-L1 adipocytes. *Dev. Cell* 9: 99–108. doi:10.1016/j.devcel.2005.04.004
- Shoelson, S.E., Lee, J., & Yuan, M., 2003. Inflammation and the IKK $\beta$ /I $\kappa$ B/NF- $\kappa$ B axis in obesity- and diet-induced insulin resistance. *Int. J. Obes.* 27: S49–S52. doi:10.1038/sj.ijo.0802501
- Singh, S., Sharma, R., Kumari, M., & Tiwari, S., 2019. Insulin receptors in the kidneys in health and disease. *World J. Nephrol.* 8: 11–22. doi:10.5527/wjn.v8.i1.11
- Singh, S., Usman, K., & Banerjee, M., 2016. Pharmacogenetic studies update in type 2 diabetes mellitus. *World J. Diabetes* 7: 302. doi:10.4239/wjd.v7.i15.302
- Sinha, S., Perdomo, G., Brown, N.F., & O’Doherty, R.M., 2004. Fatty acid-induced insulin resistance in L6 myotubes is prevented by inhibition of activation and nuclear localization of nuclear factor  $\kappa$ B. *J. Biol. Chem.* 279: 41294–41301. doi:10.1074/jbc.M406514200
- Soegondo, S., 2014. Farmakoterapi pada pengendalian glikemia diabetes melitus tipe 2, in: Setiati, S., Alwi, I., Sudoyo, A.W., Simadibrata, M., Setiyohadi, B., & Syam, A.F. (Eds.), *Buku Ajar Ilmu Penyakit Dalam*. Interna Publishing Pusat Penerbitan Ilmu Penyakit Dalam, Jakarta, pp. 2328–2333.
- Sokolowska, E., & Blachnio-Zabielska, A., 2019. The role of ceramides in insulin resistance. *Front. Endocrinol. (Lausanne)*. 10: 1–13. doi:10.3389/fendo.2019.00577
- Staels, B., & Fruchart, J.C., 2005. Therapeutic roles of peroxisome proliferator-activated receptor agonists. *Diabetes* 54: 2460–2470. doi:10.2337/diabetes.54.8.2460
- Suhono, B., & Wulan, Y., 2012. *Ensiklopedia biologi dunia tumbuhan*. PT Lentera Abadi, Jakarta.
- Sukardiman, & Ervina, M., 2020. The recent use of *Swietenia mahagoni* (L.) Jacq. as antidiabetes type 2 phytomedicine: A systematic review. *Heliyon* 6: e03536. doi:10.1016/j.heliyon.2020.e03536
- Sumida, K., Igarashi, Y., Toritsuka, N., Matsushita, T., Abe-Tomizawa, K., Aoki, M., *et al.*, 2011. Effects of DMSO on gene expression in human and rat hepatocytes. *Hum. Exp. Toxicol.* 30: 1701–1709. doi:10.1177/0960327111399325
- Sun, Y., Liu, S., Ferguson, S., Wang, L., Klepcyk, P., Yun, J.S., *et al.*, 2002. Phosphoenolpyruvate carboxykinase overexpression selectively attenuates insulin signaling and hepatic insulin sensitivity in transgenic mice. *J. Biol. Chem.* 277: 23301–23307. doi:10.1074/jbc.M200964200
- Sun, Y.P., Jin, W.F., Wang, Y.Y., Wang, G., Morris-Natschke, S.L., Liu, J.S., *et al.*, 2018.



- Chemical structures and biological activities of limonoids from the genus *swietenia* (meliaceae). *Molecules* 23: 1–17. doi:10.3390/molecules23071588
- Swiderska, E., Strycharz, J., Wroblewski, A., Szemraj, J., Drzewoski, J., & Sliwinska, A., 2018. Role of PI3K/AKT pathway in insulin-mediated glucose uptake, in: Blood Glucose Level. *IntechOpen*, pp. 1–18. doi:<http://dx.doi.org/19.5772/intechopen.80402>
- Syaiful, R.H., 2016. The effect of 1,4-bis-(3,4,5-trimetoksi-fenil)-tetrahidrofuro (3,4c)furan which isolated from the seed extract of *Swietenia macrophylla* King on cell viability: exploration on cell Vero's DNA (translate from Indonesia language). Universitas Gadjah Mada, Yogyakarta.
- Tahrani, A.A., 2017. Novel therapies in type 2 diabetes: insulin resistance. *Pract. Diabetes* 34: 161-166a. doi:10.1002/pdi.2109
- Tahrani, A.A., Bailey, C.J., Del Prato, S., & Barnett, A.H., 2011. Management of type 2 diabetes: New and future developments in treatment. *Lancet* 378: 182–197. doi:10.1016/S0140-6736(11)60207-9
- Tahrani, A.A., Barnett, A.H., & Bailey, C.J., 2013. SGLT inhibitors in management of diabetes. *Lancet Diabetes Endocrinol.* 1: 140–151. doi:10.1016/S2213-8587(13)70050-0
- Tao, R., Gong, J., Luo, X., Zang, M., Guo, W., Wen, R., *et al.*, 2010. AMPK exerts dual regulatory effects on the PI3K pathway. *J. Mol. Signal.* 5: 1–9. doi:10.1186/1750-2187-5-1
- Taylor, R., 2012. Insulin resistance and type 2 diabetes. *Diabetes* 61: 778–779. doi:10.2337/db12-0073
- Taylor, S.I., Blau, J.E., & Rother, K.I., 2015. SGLT2-inhibitors trigger downstream mechanisms that may exert adverse effects upon bone. *Lancet. Diabetes Endocrinol.* 3(1) : 1-6. doi:10.1016/S2213-8587(14)70227-X.SGLT2-inhibitors
- Thai, M. V., Guruswamy, S., Cao, K.T., Pessin, J.E., & Olson, A.L., 1998. Myocyte enhancer factor 2 (MEF2)-binding site is required for GLUT4 gene expression in transgenic mice: Regulation of MEF2 DNA binding activity in insulin-deficient diabetes. *J. Biol. Chem.* 273: 14285–14292. doi:10.1074/jbc.273.23.14285
- Thakur, M., Melzig, M.F., Fuchs, H., & Weng, A., 2011. Chemistry and pharmacology of saponins: special focus on cytotoxic properties. *Bot. Targets Ther.* 1: 19-29. doi:10.2147/btat.s17261
- Theeuwes, W.F., Gosker, H.R., Langen, R.C.J., Verhees, K.J.P., Pansters, N.A.M., Schols, A.M.W.J., *et al.*, 2017. Inactivation of glycogen synthase kinase-3 $\beta$  (GSK-3 $\beta$ ) enhances skeletal muscle oxidative metabolism. *Biochim. Biophys. Acta - Mol. Basis Dis.* 1863: 3075–3086. doi:10.1016/j.bbadis.2017.09.018
- Timm, M., Saaby, L., Moesby, L., & Hansen, E.W., 2013. Considerations regarding use of solvents in *in vitro* cell based assays. *Cytotechnology* 65: 887–894. doi:10.1007/s10616-012-9530-6
- Tong, T., Ren, N., Soomi, P., Wu, J., Guo, N., Kang, H., *et al.*, 2018. Theaflavins improve insulin sensitivity through regulating mitochondrial biosynthesis in palmitic acid-induced HepG2 cells. *Molecules* 23. doi:10.3390/molecules23123382
- Tsotetsi, T.N., Collins, N.E., Oosthuizen, M.C., & Sibeko-Matjila, K.P., 2018. Selection and evaluation of housekeeping genes as endogenous controls for quantification of mRNA transcripts in *Theileria parva* using quantitative real-time polymerase chain reaction (qPCR). *PLoS One* 13: 1–15. doi:10.1371/journal.pone.0196715
- Tsuchiya, Y., Hatakeyama, H., Emoto, N., Wagatsuma, F., Matsushita, S., & Kanzaki, M., 2010. Palmitate-induced down-regulation of sortilin and impaired GLUT4 trafficking in C2C12 myotubes. *J. Biol. Chem.* 285: 34371–34381. doi:10.1074/jbc.M110.128520
- Tumova, J., Andel, M., & Trnka, J., 2016. Excess of free fatty acids as a cause of metabolic dysfunction in skeletal muscle. *Physiol. Res.* 65: 193–207.



- Turner, M.C., 2015. Cell culture models of insulin signalling and glucose uptake. Loughborough University.
- Uhl, M., Helma, C., & Knasmüller, S., 1999. Single-cell gel electrophoresis assays with human-derived hepatoma (Hep G2) cells. *Mutat. Res. - Genet. Toxicol. Environ. Mutagen.* 441: 215–224. doi:10.1016/S1383-5718(99)00050-9
- van der Aa, M.P., Elst, M.A.J., van de Garde, E.M.W., van Mil, E.G.A.H., Knibbe, C.A.J., & van der Vorst, M.M.J., 2016. Long-term treatment with metformin in obese, insulin-resistant adolescents: results of a randomized double-blinded placebo-controlled trial. *Nutr. Diabetes* 6: 1–10. doi:10.1038/nutd.2016.37
- Varga, Z., Diebold, J., Dommann-Scherrer, C., Frick, H., Kaup, D., Noske, A., *et al.*, 2012. How reliable is Ki-67 immunohistochemistry in grade 2 breast carcinomas? A QA study of the Swiss working group of breast- and gynecopathologists. *PLoS One* 7: 1–12. doi:10.1371/journal.pone.0037379
- Vlavcheski, F., & Tsiani, E., 2018. Attenuation of free fatty acid-induced muscle insulin resistance by rosemary extract. *Nutrients* 10: 1–16. doi:10.3390/nu10111623
- Wang, J., Gao, Y., Duan, L., Wei, S., Liu, Jing, Tian, L., *et al.*, 2017. Metformin ameliorates skeletal muscle insulin resistance by inhibiting miR-21 expression in a high-fat dietary rat model. *Oncotarget* 8: 98029–98039. doi:10.18632/oncotarget.20442
- Wang, L., Waltenberger, B., Pferschy-Wenzig, E.M., Blunder, M., Liu, X., Malainer, C., *et al.*, 2014. Natural product agonists of peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ): A review. *Biochem. Pharmacol.* 92: 73–89. doi:10.1016/j.bcp.2014.07.018
- Wang, Q., Wang, N., Dong, M., Chen, F., Li, Z., & Chen, Y., 2014. GdCl<sub>3</sub> reduces hyperglycaemia through Akt/Foxo1-induced suppression of hepatic gluconeogenesis in type 2 diabetic mice. *Clin. Sci.* 127: 91–100. doi:10.1042/CS20130670
- Wang, T., Zheng, L., Zhao, T., Zhang, Q., Liu, Z., Liu, X., *et al.*, 2020. Anti-diabetic effects of sea cucumber (*Holothuria nobilis*) hydrolysates in streptozotocin and high-fat-diet induced diabetic rats via activating the PI3K/Akt pathway. *J. Funct. Foods* 75: 104224. doi:10.1016/j.jff.2020.104224
- Wang, Z.B., Zeng, H.C., Wei, H.S., Yi, G.H., Yu, J., Wang, Y.T., *et al.*, 2012. NO-1886 ameliorates glycogen metabolism in insulin-resistant HepG2 cells by GSK-3 $\beta$  signalling. *J. Pharm. Pharmacol.* 64: 293–301. doi:10.1111/j.2042-7158.2011.01402.x
- Wei, Y., Yang, H., Zhu, C., Deng, J., & Fan, D., 2020. Ginsenoside Rg5 relieves type 2 diabetes by improving hepatic insulin resistance in db/db mice. *J. Funct. Foods* 71: 104014. doi:10.1016/j.jff.2020.104014
- Weigert, C., Brodbeck, K., Staiger, H., Kausch, C., Machicao, F., Häring, H.U., *et al.*, 2004. Palmitate, but not unsaturated fatty acids, induces the expression of interleukin-6 in human myotubes through proteasome-dependent activation of nuclear factor- $\kappa$ B. *J. Biol. Chem.* 279: 23942–23952. doi:10.1074/jbc.M312692200
- Williams, V.L., Martin, R.E., Franklin, J.L., Hardy, R.W., & Messina, J.L., 2012. Injury-induced insulin resistance in adipose tissue. *Biochem. Biophys. Res. Commun.* 421: 442–448. doi:10.1016/j.bbrc.2012.03.146
- World Health Organization (WHO), 2016. Global Report on Diabetes. World Health Organization Press, Geneva.
- Wong, C.Y., Al-Salami, H., & Dass, C.R., 2020. C2C12 cell model: its role in understanding of insulin resistance at the molecular level and pharmaceutical development at the preclinical stage. *J. Pharm. Pharmacol.* 72: 1667–1693. doi:10.1111/jphp.13359
- World Health Organization (WHO), 1999. Definition, diagnosis, and classification of diabetes mellitus and its complication. *Amelior. Ment. Disabil. Quest. Retard.* doi:10.5005/jp/books/12626\_7



- Wu, Y., Ding, Y., Tanaka, Y., & Zhang, W., 2014. Risk factors contributing to type 2 diabetes and recent advances in the treatment and prevention. *Int. J. Med. Sci.* 11: 1185–1200. doi:10.7150/ijms.10001
- Xi, M., Hai, C., Tang, H., Chen, M., Fang, K., & Liang, X., 2008. Antioxidant and antiglycation properties of total saponins extracted from traditional Chinese medicine used to treat diabetes mellitus. *Phyther. Res.* 22: 228–237. doi:<https://doi.org/10.1002/ptr.2297>
- Xiong, H., Zhang, S., Zhao, Z., Zhao, P., Chen, L., & Mei, Z., 2018. Antidiabetic activities of entagenic acid in type 2 diabetic db/db mice and L6 myotubes via AMPK/GLUT4 pathway. *J. Ethnopharmacol.* 211: 366–374. doi:10.1016/j.jep.2017.10.004
- Xiong, X., Song, Q., Han, C., Gan, W., Wei, S., Liu, H., *et al.*, 2016. Insulin promotes the expression of the gluconeogenic rate-limiting enzymes phosphoenolpyruvate carboxykinase (PEPCK) and glucose 6-phosphatase (G6pase) through PI3K/Akt/mTOR signaling pathway in goose hepatocytes. *Brazilian J. Poult. Sci.* 18: 395–400. doi:<http://dx.doi.org/10.1590/1806-9061-2015-0080>
- Xu, J., Wang, S., Feng, T., Chen, Y., & Yang, G., 2018. Hypoglycemic and hypolipidemic effects of total saponins from *Stauntonia chinensis* in diabetic db/db mice. *J. Cell. Mol. Med.* 22: 6026–6038. doi:10.1111/jcmm.13876
- Xu, Q., Wang, L., Luo, J., & Shi, D., 2017. The hot and potential targets of type 2 diabetes mellitus treatment in recent decade. *Curr. Drug Targets* 19: 55-69.
- Yadollah, S., Kazemipour, N., Bakhtiyari, S., & Nazifi, S., 2017. Palmitate-induced insulin resistance is attenuated by pioglitazone and EGCG through reducing the gluconeogenic key enzymes expression in HepG2 cells. *J. Med. Life* 10: 244–249.
- Yan, J., Wang, C., Jin, Y., Meng, Q., Liu, Q., Liu, Z., *et al.*, 2018. Catalpol ameliorates hepatic insulin resistance in type 2 diabetes through acting on AMPK/NOX4/PI3K/AKT pathway. *Pharmacol. Res.* 130: 466–480. doi:10.1016/j.phrs.2017.12.026
- Yang, J., Kalhan, S.C., & Hanson, R.W., 2009. What is the metabolic role of phosphoenolpyruvate carboxykinase? *J. Biol. Chem.* 284: 27025–27029. doi:10.1074/jbc.R109.040543
- Yang, M., Wei, D., Mo, C., Zhang, J., Wang, X., Han, X., *et al.*, 2013. Saturated fatty acid palmitate-induced insulin resistance is accompanied with myotube loss and the impaired expression of health benefit myokine genes in C2C12 myotubes. *Lipids Health Dis.* 12: 1. doi:10.1186/1476-511X-12-104
- Yang, R., Wang, L., Xie, J., Li, X., Liu, S., Qiu, S., *et al.*, 2018. Treatment of type 2 diabetes mellitus via reversing insulin resistance and regulating lipid homeostasis in vitro and in vivo using cajanonic acid A. *Int. J. Mol. Med.* 42: 2329–2342. doi:10.3892/ijmm.2018.3836
- Yang, S.J., Choi, J.M., Park, S.E., Rhee, E.J., Lee, W.Y., Oh, K.W., *et al.*, 2015. Preventive effects of bitter melon (*Momordica charantia*) against insulin resistance and diabetes are associated with the inhibition of NF-κB and JNK pathways in high-fat-fed OLETF rats. *J. Nutr. Biochem.* 26: 234–240. doi:10.1016/j.jnutbio.2014.10.010
- Yin, J., Zuberi, A., Gao, Z., Liu, D., Liu, Z., & Ye, J., 2009. *Shilianhua* extract inhibits GSK-3β and promotes glucose metabolism. *Am. J. Physiol. - Endocrinol. Metab.* 296: 1275–1280. doi:10.1152/ajpendo.00092.2009
- Yogalakshmi, B., Bhuvaneshwari, S., Sreeja, S., & Anuradha, C.V., 2014. Grape seed proanthocyanidins and metformin act by different mechanisms to promote insulin signaling in rats fed high calorie diet. *J. Cell Commun. Signal.* 8: 13–22. doi:10.1007/s12079-013-0210-x
- Yu, Y., & Chai, J., 2015. The function of miRNAs and their potential as therapeutic targets in burn-induced insulin resistance (Review). *Int. J. Mol. Med.* 35: 305–310.



- Yuan, C., Gao, J., Guo, J., Bai, L., Marshall, C., Cai, Z., *et al.*, 2014. Dimethyl sulfoxide damages mitochondrial integrity and membrane potential in cultured astrocytes. *PLoS One* 9(9): 1-9 doi:10.1371/journal.pone.0107447
- Yuan, H.D., Kim, D.Y., Quan, H.Y., Kim, S.J., Jung, M.S., & Chung, S.H., 2012. Ginsenoside Rg2 induces orphan nuclear receptor SHP gene expression and inactivates GSK3 $\beta$  via AMP-activated protein kinase to inhibit hepatic glucose production in HepG2 cells. *Chem. Biol. Interact.* 195: 35–42. doi:10.1016/j.cbi.2011.10.006
- Yuan, Y.L., Lin, B.Q., Zhang, C.F., Cui, L.L., Ruan, S.X., Yang, Z.L., *et al.*, 2016. Timosaponin B-II ameliorates palmitate-induced insulin resistance and inflammation via IRS1/PI3K/Akt and IKK/NF- $\kappa$ B Pathways. *Am. J. Chin. Med.* 44: 755–769. doi:10.1142/S0192415X16500415
- Yung, H.W., Charnock-Jones, D.S., & Burton, G.J., 2011. Regulation of AKT phosphorylation at Ser473 and Thr308 by endoplasmic reticulum stress modulates substrate specificity in a severity dependent manner. *PLoS One* 6(3): 1-12. doi:10.1371/journal.pone.0017894
- Yusuf, M., 2016. The effects of 1,4-bis-(3,4,5-trimetoxy-fenyl)-tetrahydrofuro (3,4-c) furan isolate on insulin resistance and expression of IRS1 serine 307 protein in skeletal tissues of DM type 2 rats model (translate from Indonesia language). Universitas Gadjah Mada.
- Yuzefovych, L., Wilson, G., & Rachek, L., 2010. Different effects of oleate vs. palmitate on mitochondrial function, apoptosis, and insulin signaling in L6 skeletal muscle cells: Role of oxidative stress. *Am. J. Physiol. - Endocrinol. Metab.* 299: 1096–1105. doi:10.1152/ajpendo.00238.2010
- Zang, Y., Fan, L., Chen, J., Huang, R., & Qin, H., 2018. Improvement of lipid and glucose metabolism by capsate in palmitic acid-treated HepG2 Cells via activation of the AMPK/SIRT1 signaling pathway. *J. Agric. Food Chem.* 66: 6772–6781. doi:10.1021/acs.jafc.8b01831
- Zendarto, D., 2015. Mahoni (*Swietenia mahagoni* (L.) Jacq [www document]. URL <https://deslisumatran.wordpress.com/2015/03/23/mahoni-swietenia-mahagoni-1-jacq/>
- Zha, L., Chen, J., Sun, S., Mao, L., Chu, X., Deng, H., *et al.*, 2014. Soyasaponins can blunt inflammation by inhibiting the reactive oxygen species-mediated activation of PI3K/Akt/NF- $\kappa$ B pathway. *PLoS One* 9: 1–9. doi:10.1371/journal.pone.0107655
- Zhang, G., Cai, X., He, L., Qin, D., Li, H., & Fan, X., 2020a. Skimmin improves insulin resistance via regulating the metabolism of glucose: In vitro and in vivo models. *Front. Pharmacol.* 11: 1–9. doi:10.3389/fphar.2020.00540
- Zhang, H., Ge, Z., Ang, S., Meng, R., Bi, Y., & Zhu, D., 2017. Erythropoietin ameliorates PA-induced insulin resistance through the IRS/AKT/FOXO1 and GSK-3 $\beta$  signaling pathway, and inhibits the inflammatory response in HepG2 cells. *Mol. Med. Rep.* 16: 2295–2301. doi:10.3892/mmr.2017.6810
- Zhang, J., & Liu, F., 2014. Tissue-specific insulin signaling in the regulation of metabolism and aging. *IUBMB Life* 66: 485–495. doi:10.1002/iub.1293
- Zhang, Q., Kong, X., Yuan, H., Guan, H., Li, Y., & Niu, Y., 2019a. Mangiferin improved palmitate-induced-insulin resistance by promoting free fatty acid metabolism in HepG2 and C2C12 cells via PPAR $\alpha$ : Mangiferin improved insulin resistance. *J. Diabetes Res.* 2019: 1-13. doi:10.1155/2019/2052675
- Zhang, X., Yang, S., Chen, J., & Su, Z., 2019b. Unraveling the regulation of hepatic gluconeogenesis. *Front. Endocrinol. (Lausanne)*. 10: 1–17. doi:10.3389/fendo.2018.00802
- Zhang, Y., Chen, J., Zeng, Y., Huang, D., & Xu, Q., 2019c. Involvement of AMPK activation in the inhibition of hepatic gluconeogenesis by *Ficus carica* leaf extract in diabetic mice



doi:10.1016/j.biopha.2018.10.077

- Zhang, Y., Yan, L.S., Ding, Y., Cheng, B.C.Y., Luo, G., Kong, J., *et al.*, 2020b. *Edgeworthia gardneri* (Wall.) Meisn. water extract ameliorates palmitate induced insulin resistance by regulating IRS1/GSK3b/FOXO1 signaling pathway in human HepG2 hepatocytes. *Front. Pharmacol.* 10: 1–16. doi:10.3389/fphar.2019.01666
- Zheng, Y., Ley, S.H., & Hu, F.B., 2018. Global aetiology and epidemiology of type 2 diabetes mellitus and its complications. *Nat. Rev. Endocrinol.* 14: 88–98. doi:10.1038/nrendo.2017.151
- Zhou, T., Meng, X., Che, H., Shen, N., Xiao, D., Song, X., *et al.*, 2016. Regulation of insulin resistance by multiple MiRNAs via targeting the GLUT4 signalling pathway. *Cell. Physiol. Biochem.* 38: 2063–2078. doi:10.1159/000445565
- Zhu, R., Zheng, J., Chen, L., Gu, B., & Huang, S., 2016. Astragaloside IV facilitates glucose transport in C2C12 myotubes through the IRS1/AKT pathway and suppresses the palmitate-induced activation of the IKK/IB pathway. *Int. J. Mol. Med.* 37: 1697–1705. doi:10.3892/ijmm.2016.2555
- Zhu, Y.X., Hu, H.Q., Zuo, M.L., Mao, L., Song, G.L., Li, T.M., *et al.*, 2021. Effect of oxymatrine on liver gluconeogenesis is associated with the regulation of pepck and g6pase expression and akt phosphorylation. *Biomed. Reports* 15: 1–10. doi:10.3892/br.2021.1432