

DAFTAR PUSTAKA

- Ai, F., Deng, M., Mao, M., Xu, T., & Zhu, L. (2018). Depressive symptoms screening in postmenopausal women with symptomatic pelvic organ prolapse. *Menopause*, 25(3), 314–319. <https://doi.org/10.1097/GME.0000000000001006>
- Ansari, M. K., Sharma, P. P., & Khan, S. (2021). Pelvic Organ Prolapse in Perimenopausal and Menopausal Women. *Journal of Obstetrics and Gynecology of India*, 0123456789. <https://doi.org/10.1007/s13224-021-01524-8>
- Barati, M., Akbari-heidari, H., Samadi-yaghin, E., Jenabi, E., Jormand, H., & Kamyari, N. (2021). The factors associated with the quality of life among postmenopausal women. *BMC Women's Health*, 21(1), 1–8. <https://doi.org/10.1186/s12905-021-01361-x>
- Barber, M. D. (2016). Pelvic organ prolapse. *BMJ (Online)*, 354(July), 1–9. <https://doi.org/10.1136/bmj.i3853>
- Barber, M. D., & Maher, C. (2013). Epidemiology and outcome assessment of pelvic organ prolapse. *International Urogynecology Journal and Pelvic Floor Dysfunction*, 24(11), 1783–1790. <https://doi.org/10.1007/s00192-013-2169-9>
- Caylan, A., Aydemir, I., Dagdeviren, N., Akturk, Z., Set, T., Oztora, S., & Enec Can, F. (2011). Evaluation of health related quality of life among perimenopausal Turkish women. *HealthMED*, 5(4), 741–752.
- Corton, M. M. (2009). Anatomy of Pelvic Floor Dysfunction. In *Obstetrics and Gynecology Clinics of North America* (Vol. 36, Issue 3, pp. 401–419). <https://doi.org/10.1016/j.ogc.2009.09.002>
- Digesu, G. A., Khullar, V., Cardozo, L., Robinson, D., & Salvatore, S. (2005). P-QOL: A validated questionnaire to assess the symptoms and quality of life of women with urogenital prolapse. *International Urogynecology Journal*, 16(3), 176–181. <https://doi.org/10.1007/s00192-004-1225-x>
- Erekson, E. A., Li, F. Y., Martin, D. K., & Fried, T. R. (2016). Vulvovaginal symptoms prevalence in postmenopausal women and relationship to other menopausal symptoms and pelvic floor disorders. *Menopause*, 23(4), 368–375. <https://doi.org/10.1097/GME.0000000000000549>
- Fornell, E. U., Wingren, G., & Kjølhedde, P. (2004). Factors associated with pelvic floor dysfunction with emphasis on urinary and fecal incontinence and genital prolapse: An epidemiological study. *Acta Obstetrica et Gynecologica Scandinavica*, 83(4), 383–389. <https://doi.org/10.1111/j.0001-6349.2004.00367.x>

- Frota, I. P. R., Rocha, A. B. O., Neto, J. A. V., Vasconcelos, C. T. M., De Magalhaes, T. F., Karbage, S. A. L., Augusto, K. L., Nascimento, S. L. Do, Haddad, J. M., & Bezerra, L. R. P. S. (2018). Pelvic floor muscle function and quality of life in postmenopausal women with and without pelvic floor dysfunction. *Acta Obstetrica et Gynecologica Scandinavica*, 97(5), 552–559. <https://doi.org/10.1111/aogs.13305>
- Hakim, S., & Maharani, C. R. (2017). Perubahan Skor Kualitas Hidup Pasien Prolaps Organ Panggul setelah Operasi Vagina yang Diukur dengan Kuesioner Pelvic Floor Distress Inventory (PFDI-20) dan Pelvic Floor Impact Questionnaire (PFIQ-7). *Indonesian Journal of Obstetrics and Gynecology*, 5(3), 164.
- Handa, V. L., Blomquist, J. L., Knoepf, L. R., Hoskey, K. A., McDermott, K. C., & Muñoz, A. (2011). Pelvic floor disorders 5-10 years after vaginal or cesarean childbirth. *Obstetrics and Gynecology*, 118(4), 777–784. <https://doi.org/10.1097/AOG.0b013e3182267f2f>
- Hilditch, J. R., Lewis, J., Peter, A., Van Maris, B., Ross, A., Franssen, E., Guyatt, G. H., Norton, P. G., & Dunn, E. (1996). A menopause-specific quality of life questionnaire: Development and psychometric properties. *Maturitas*, 24(3), 161–175. [https://doi.org/10.1016/S0378-5122\(96\)82006-8](https://doi.org/10.1016/S0378-5122(96)82006-8)
- Jelovsek, J. E., Chagin, K., Gyhagen, M., Hagen, S., Wilson, D., Kattan, M. W., Elders, A., Barber, M. D., Areskoug, B., MacArthur, C., & Milsom, I. (2018). Predicting risk of pelvic floor disorders 12 and 20 years after delivery. *American Journal of Obstetrics and Gynecology*, 218(2), 222.e1-222.e19. <https://doi.org/10.1016/j.ajog.2017.10.014>
- Jelovsek, J. E., Maher, C., & Barber, M. D. (2007). Pelvic organ prolapse. *Lancet*, 369(9566), 1027–1038. [https://doi.org/10.1016/S0140-6736\(07\)60462-0](https://doi.org/10.1016/S0140-6736(07)60462-0)
- Junizaf, & Santoso, B. I. (2013). Penatalaksanaan Prolaps Organ Panggul. *Himpunan Uroginekologi Indonesia*.
- Kalarhoudi M, Taebi M, Sadat Z, Saberi F. Assessment of quality of life in menopausal periods: a population study in kashan, iran. *Iran Red Crescent Med J*. 2011 Nov;13(11):811-7. Epub 2011 Nov 1. PMID: 22737420; PMCID: PMC3371893.
- Kim, C. M., Jeon, M. J., Chung, D. J., Kim, S. K., Kim, J. W., & Bai, S. W. (2007). Risk factors for pelvic organ prolapse. *International Journal of Gynecology and Obstetrics*, 98(3), 248–251. <https://doi.org/10.1016/j.ijgo.2007.02.019>
- Koirala, D., Thapa, N., & Shrestha, S. (2020). Quality of life of postmenopausal women of Kaski district. *Nepal Journal of Obstetrics and Gynaecology*, 15(1), 43–49. <https://doi.org/10.3126/njog.v15i1.29340>
- Kołodzyńska, G., Zalewski, M., & Rozek-Piechura, K. (2019). Urinary incontinence in postmenopausal women - Causes, symptoms, treatment. *Przegląd*

- Menopausalny*, 18(1), 46–50. https://doi.org/10.5114/pm.2019.84157_rfseq1
- Lobo, R. A., Davis, S. R., De Villiers, T. J., Gompel, A., Henderson, V. W., Hodis, H. N., Lumsden, M. A., Mack, W. J., Shapiro, S., & Baber, R. J. (2014). Prevention of diseases after menopause. *Climacteric*, 17(5), 540–556. <https://doi.org/10.3109/13697137.2014.933411>
- Lukacz, E. S., Lawrence, J. M., Contreras, R., Nager, C. W., & Lubner, K. M. (2006). Parity, Mode of Delivery, and Pelvic Floor Disorders LEVEL OF EVIDENCE: II-2. *Obstet Gynecol*, 107(6), 1253–1260.
- Memon, H. U., & Handa, V. L. (2013). Vaginal childbirth and pelvic floor disorders. *Women's Health*, 9(3), 265–277. <https://doi.org/10.2217/whe.13.17>
- Micussi, M. T., Freitas, R. P., Angelo, P. H., Soares, E. M., Lemos, T. M., & Maranhão, T. M. (2015). Evaluation of the relationship between the pelvic floor muscles and insulin resistance. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, 8, 409–413. <https://doi.org/10.2147/DMSO.S85816>
- Nygaard, I., Bradley, C., & Brandt, D. (2004). Pelvic organ prolapse in older women: Prevalence and risk factors. *Obstetrics and Gynecology*, 104(3), 489–497. <https://doi.org/10.1097/01.AOG.0000136100.10818.d8>
- Pangastuti, N., Iman, S. B., Denny, A., & Emilia, O. (2020). Validation test of Indonesian pelvic floor distress inventory-20 (Indonesian PFDI-20). *Journal of Thee Medical Sciences (Berkala Ilmu Kedokteran)*, 52(02). <https://doi.org/10.19106/jmedsci005202202004>
- Persu, C., Chapple, C. R., Cauni, V., Gutue, S., & Geavlete, P. (2011). Pelvic Organ Prolapse Quantification System (POP-Q) - a new era in pelvic prolapse staging. *Journal of Medicine and Life*, 4(1), 75–81.
- Politi, M. C., Schleinitz, M. D., & Col, N. F. (2008). Revisiting the duration of vasomotor symptoms of menopause: A meta-analysis. *Journal of General Internal Medicine*, 23(9), 1507–1513. <https://doi.org/10.1007/s11606-008-0655-4>
- Purwara, B. H. (2014). Faktor Risiko Penderita Prolapsus Organ Panggul terhadap Hiatus Genitalis, Panjang Total Vagina, dan Perineal Body. *Majalah Kedokteran Bandung*, 46(1), 57–60. <https://doi.org/10.15395/mkb.v46n1.229>
- Putri, D. I., Wati, D. M., & Ariyanto, Y. (2014). *QoL Menopausal Women*. 2(1).
- Rahmatyah, R., Hardiyanto, G., & Susanti, D. (2018). Correlation between Pelvic Organ Prolapse Symptoms and Age of Menopause. *Majalah Obstetri & Ginekologi*, 25(1), 30. <https://doi.org/10.20473/mog.v25i12017.30-32>
- Shobeiri, F., Jenabi, E., Hazavehei, S. M. M., & Roshanaei, G. (2016). Quality of Life in Postmenopausal Women in Iran: A Population-based Study. *Journal of Menopausal Medicine*, 22(1), 31. <https://doi.org/10.6118/jmm.2016.22.1.31>

- Sung, V. W., & Hampton, B. S. (2009). Epidemiology of Pelvic Floor Dysfunction. *Obstetrics and Gynecology Clinics of North America*, 36(3), 421–443. <https://doi.org/10.1016/j.ogc.2009.08.002>
- Sydora, B. C., Fast, H., Campbell, S., Yuksel, N., Lewis, J. E., & Ross, S. (2016). Use of the Menopause-Specific Quality of Life (MENQOL) questionnaire in research and clinical practice: A comprehensive scoping review. *Menopause*, 23(9), 1038–1051. <https://doi.org/10.1097/GME.0000000000000636>
- Thomas, H. N., Neal-Perry, G. S., & Hess, R. (2018). Female Sexual Function at Midlife and Beyond. *Obstetrics and Gynecology Clinics of North America*, 45(4), 709–722. <https://doi.org/10.1016/j.ogc.2018.07.013>
- Walters, M.D dan Karram, M.M. 2007. Urogynecology and Reconstructive Pelvic Surgery. Elsevier Inc.
- Weidner, A. C., Jamison, M. G., Branham, V., South, M. M., Borawski, K. M., & Romero, A. A. (2006). Neuropathic injury to the levator ani occurs in 1 in 4 primiparous women. *American Journal of Obstetrics and Gynecology*, 195(6), 1851–1856. <https://doi.org/10.1016/j.ajog.2006.06.062>
- World Health Organization. (1998). Whoqol user manual: Programme on mental health. *World Health Organization*, 1–19. https://doi.org/10.1007/SpringerReference_28001