

DAFTAR PUSTAKA

1. Wijayanto T, Masloman N. Sindrom Lennox-Gastaut. *Maj Kedokt Indones.* 2005;55:36–41.
2. Shneker BF, Fountain NB. *Epilepsy. Disease-a-Month.* 2003;5029(July).
3. Hageman JR. Pediatric seizures. *Pediatr Ann.* 2013;42(12):494.
4. Perucca P, Scheffer IE, Kiley M. The management of epilepsy in children and adults. *Med J Aust.* 2018;208(5):226–33.
5. Kementerian Kesehatan Republik Indonesia. Pedoman Nasional Pelayanan Kedokteran Tata Laksana Epilepsi pada Anak. Menteri Kesehatan Republik Indones [Internet]. 2017;1(1):1–79. Available from: <http://e-journal.usd.ac.id/index.php/LLT%0Ahttp://jurnal.untan.ac.id/index.php/jpdpb/article/viewFile/11345/10753%0Ahttp://dx.doi.org/10.1016/j.sbspro.2015.04.758%0Awww.iosrjournals.org>
6. Suwarba IGNM. Insidens dan Karakteristik Klinis Epilepsi pada Anak. *Sari Pediatr.* 2016;13(2):123.
7. Moreno A, Peel M. Posttraumatic seizures in survivors of torture: manifestations, diagnosis, and treatment. *J Immigr Health.* 2004;6(4):179–86.
8. Laxer KD, Trinkka E, Hirsch LJ, Cendes F, Langfitt J, Delanty N, et al. The consequences of refractory epilepsy and its treatment. *Epilepsy Behav.* 2014;37:59–70.
9. Martin-McGill KJ, Jackson CF, Bresnahan R, Levy RG, Cooper PN. Ketogenic diets for drug-resistant epilepsy. *Cochrane Database Syst Rev.* 2018;2018(11).
10. Zhu D, Wang M, Wang J, Yuan J, Niu G, Zhang G, et al. Ketogenic diet effects on neurobehavioral development of children with intractable epilepsy: A prospective study. *Epilepsy Behav.* 2016;55(June 2013):87–91.
11. Caraballo RH, Fortini S, Fresler S, Armeno M, Ariela A, Cresta A, et al. Ketogenic diet in patients with Lennox-Gastaut syndrome. *Seizure [Internet].* 2014;23(9):751–5. Available from: <http://dx.doi.org/10.1016/j.seizure.2014.06.005>
12. Williams TJ, Cervenka MC. The role for ketogenic diets in epilepsy and status epilepticus in adults. Vol. 2, *Clinical Neurophysiology Practice.* 2017. p. 154–60.
13. Sampath a, Kossoff EH, Furth SL, Pyzik PL, Vining EP. Kidney stones and the ketogenic diet: risk factors and prevention. *J Child Neurol.* 2007;22(4):375–8.
14. Kwiterovich PO, Vining EPG, Pyzik P, Skolasky R, Freeman JM. Effect of a High-Fat Ketogenic Diet on Plasma Levels of Lipids, Lipoproteins, and Apolipoproteins in Children. *J Am Med Assoc.* 2003;290(7):912–20.
15. Dashti HM, Mathew TC, Hussein T, Asfar SK, Behbahani A, Khourshed M a., et al. Long-term effects of a ketogenic diet in obese patients. *Exp Clin Cardiol.* 2004;9(3):200–5.

16. Mangunatmadja I, Mulyani DI, Pardede SO, Tridjadja B, Wulandari HF. Faktor Risiko Epilepsi Intraktabel pada Anak dengan Epilepsi Umum. 2019;59–65.
17. Ostendorf AP, Ng YT. Treatment-resistant Lennox-Gastaut syndrome: Therapeutic trends, challenges and future directions. *Neuropsychiatr Dis Treat*. 2017;13:1131–40.
18. Novita Novita; Imam Rusdi CT. Terapi sindrom Lennox-Gastaut. *Berk Neurosains*. 2018;17(3):162–8.
19. Qu S, Catron M, Zhou C, Janve V, Shen W, Howe RK, et al. GABAA receptor $\beta 3$ subunit mutation D120N causes Lennox–Gastaut syndrome in knock-in mice. *Brain Commun*. 2020;2(1):1–20.
20. Saleh TA, Stephen L. Lennox gastaut syndrome, review of the literature and a case report. *Head Face Med*. 2008;4(1):1–7.
21. Misiewicz Runyon A, So T-Y. The Use of Ketogenic Diet in Pediatric Patients with Epilepsy. *ISRN Pediatr*. 2012;2012:1–10.
22. Oguni H, Hayashi K, Osawa M. Long-term prognosis of Lennox-Gastaut syndrome. *Epilepsia*. 1996;37(SUPPL. 3):44–7.
23. Rantala H, Putkonen T. Occurrence, outcome, and prognostic factors of infantile spasms and Lennox-Gastaut syndrome. *Epilepsia*. 1999;40(3):286–9.
24. Fisher RS, Cross JH, French JA, Higurashi N, Hirsch E, Jansen FE, et al. Operational classification of seizure types by the International League Against Epilepsy: Position Paper of the ILAE Commission for Classification and Terminology. *Epilepsia*. 2017;58(4):522–30.
25. Kovac S, Diehl B. Atonic phenomena in focal seizures : Nomenclature , clinical findings and pathophysiological concepts. *Seizure Eur J Epilepsy*. 2012;21(8):561–7.
26. Ngurah G, Wisnu P, Berawi N, Wahyudo R, Berawi KN. Diet Ketogenik: Terapi Nonfarmakologis yang Menjanjikan untuk Epilepsi. *Majority*. 2017;7(1):118–22.
27. Vera R, Dewi M, Nursiah N. Sindrom Epilepsi Pada Anak. *Maj Kedokt Sriwij*. 2014;46(1):72–6.
28. Lefevre F, Aronson N. Ketogenic diet for the treatment of refractory epilepsy in children: A systematic review of efficacy. *Pediatrics*. 2000;105(4).
29. Raju KNV, Gulati S, Kabra M, Agarwala A, Sharma S, Pandey RM, et al. Efficacy of 4:1 (classic) versus 2.5:1 ketogenic ratio diets in refractory epilepsy in young children: A randomized open labeled study. *Epilepsy Res [Internet]*. 2011;96(1–2):96–100. Available from: <http://dx.doi.org/10.1016/j.epilepsyres.2011.05.005>
30. Kwan P, Arzimanoglou A, Berg AT, Brodie MJ, Hauser WA, Mathern G, et al. Definition of drug resistant epilepsy: Consensus proposal by the ad hoc Task Force of the ILAE Commission on Therapeutic Strategies. *Epilepsia*. 2010;51(6):1069–77.
31. Hancock E, Cross H. Treatment of Lennox-Gastaut syndrome. *Cochrane database Syst Rev [Internet]*. 2003;(3):CD003277. Available from:

- <http://www.ncbi.nlm.nih.gov/pubmed/12917958>
32. Misiewicz Runyon A, So T-Y. The Use of Ketogenic Diet in Pediatric Patients with Epilepsy. *ISRN Pediatr*. 2012;2012:1–10.
 33. Bostock ECS, Kirkby KC, Taylor BVM. The current status of the ketogenic diet in psychiatry. *Front Psychiatry*. 2017;8(MAR):1–10.
 34. Wieser HG, Blume WT, Fish D, Goldensohn E, Hufnagel A, King D, et al. Proposal for a New Classification of Outcome with Respect to Epileptic Seizures Following Epilepsy Surgery. 2001;42(2):282–6.
 35. Wu Q, Wang H, Fan YY, Zhang JM, Liu XY, Fang XY, et al. Ketogenic diet effects on 52 children with pharmaco-resistant epileptic encephalopathy: A clinical prospective study. *Brain Behav*. 2018;8(5):1–8.
 36. Kim EH, Ko TS. Cognitive impairment in childhood onset epilepsy: Up-to-date information about its causes. *Korean J Pediatr*. 2016;59(4):155–64.
 37. Vining EPG. Tonic and atonic seizures: Medical therapy and ketogenic diet. *Epilepsia*. 2009;50(SUPPL. 8):21–4.
 38. Rini RIRS. Mengenali gejala dan penyebab dari conduct disorder. *J Psycho Idea Tahun 8 No 1*. 2010;1–17.
 39. Dunn DW. Neuropsychiatric aspects of epilepsy in children. *Epilepsy Behav*. 2003;4(2):101–6.
 40. Searight HR, Rottnek F, Abby SL. Conduct disorder: Diagnosis and treatment in primary care. *Prim Care Companion J Clin Psychiatry*. 2001;3(2):89.
 41. Freeman JM, Kossoff EH, Hartman AL. The ketogenic diet: One decade later. *Pediatrics*. 2007;119(3):535–43.
 42. Coppola G, Natale F, Torino A, Capasso R, D’Aniello A, Pironti E, et al. The impact of the ketogenic diet on arterial morphology and endothelial function in children and young adults with epilepsy: A case-control study. *Seizure*. 2014;23(4):260–5.
 43. Cotter DG, Ercal B, Huang X, Leid JM, D’Avignon DA, Graham MJ, et al. Ketogenesis prevents diet-induced fatty liver injury and hyperglycemia. *J Clin Invest*. 2014;124(12):5175–90.
 44. Groesbeck DK, Bluml RM, Kossoff EH. Long-term use of the ketogenic diet in the treatment of epilepsy. *Dev Med Child Neurol*. 2006;48(12):978–81.
 45. Partsalaki I, Karvela A, Spiliotis BE. Metabolic impact of a ketogenic diet compared to a hypocaloric diet in obese children and adolescents. *J Pediatr Endocrinol Metab*. 2012;25(7–8):697–704.
 46. Sri S. Nasar, Asosiasi Dietisien Indonesia (AsDI), Universitas Indonesia Fakultas Kedokteran, Ikatan Dokter Anak Indonesia (IDAI) PAGI (PERSAGI). *Penuntun Diet Anak*. 3rd ed. Nasar SS, editor. Jakarta: Fakultas Kedokteran Universitas Indonesia; 2014. 164–172 p.
 47. Frick PJ. Effective Interventions for Children and Adolescents with Conduct Disorder. *Can J Psychiatry*. 2001;46:597–607.
 48. Casetta I, Granierí E, Desiderò M, Monetti VC, Tola MR, Paolino E, et al. Phenytoin-induced gingival overgrowth: A community-based cross-sectional

- study in ferrara, italy. *Neuroepidemiology*. 1997;16(6):296–303.
49. Utomo T, Husni A, Hanum F. Pemberian Fenitoin Oral dan Timbulnya Hiperplasia Ginggiva pada Pasien Epilepsi. *Media Med Indones*. 2011;45(3):200-205–205.
 50. Ng YT, Conry J, Mitchell WG, Buchhalter J, Isojarvi J, Lee D, et al. Clobazam is equally safe and efficacious for seizures associated with Lennox-Gastaut syndrome across different age groups: Post hoc analyses of short- and long-term clinical trial results. *Epilepsy Behav* [Internet]. 2015;46:221–6. Available from: <http://dx.doi.org/10.1016/j.yebeh.2015.01.037>
 51. Autry AR, Trevathan E, Van Naarden Braun K, Yeargin-Allsopp M. Increased risk of death among children with lennox-gastaut syndrome and infantile spasms. *J Child Neurol*. 2010;25(4):441–7.
 52. Ibrahim Z, Provinsi D. Hubungan Kepatuhan Pengobatan Terhadap Kejadian Kejang Pada Pasien Epilepsi Yang Bebas Kejang Selama Minimal 1 Tahun Pengobatan Di Poli Neurologi RSUD Dr . a . Dadi Tjokrodipo Bandar Lampung Tahun 2014. *J Fak Kedokt*. 2015;2(4):183–7.
 53. Glaser BA, Kronsoble KM, Forkner CBW. Parents and teachers as raters of children's problem behaviors. *Child Fam Behav Ther*. 1997;19(4):1–13.
 54. Lowe LA. Using the Child Behavior Checklist in Assessing Conduct Disorder : Issues of Reliability and Validity. 1998;8(3):286–301.
 55. Özdemir R, Güzel O, Küçük M, Karadeniz C, Yılmaz Ü, Calik T, et al. The Impact of 3:1 Ketogenic Diet on Cardiac Repolarization Changes in Children with Refractory Seizures: A Prospective Follow-Up Study. *Neuropediatrics*. 2016;47(3):157–61.
 56. Groesbeck DK, Bluml RM, Kossoff EH. Long-term use of the ketogenic diet in the treatment of epilepsy. *Dev Med Child Neurol*. 2006;48(12):978–81.