

- Abayli, B., Gençdal, G., Değirmencioglu, S. 2018. Correlation Between Neutrophil/Lymphocyte Ratio and Ranson Score in Acute Pancreatitis. *Journal of Clinical Laboratory Analysis*. Vol.32: 1-4.
- Acar, R. D., Sahin, M. and Kirma, C. (2013) ‘One of the most urgent vascular circumstances: Acute limb ischemia.’, *SAGE open medicine*, 1, p. 2050312113516110. doi: 10.1177/2050312113516110.
- Alkharizmi (2017) *HUBUNGAN ANTARA STATUS GIZI, DIABETES MELITUS, DURASI OPERASI DENGAN KEJADIAN INFEKSI LUKA OPERASI PADA PASIEN APENDISITIS DI RUMAH SAKIT UMUM KABUPATEN TANGERANG TAHUN 2015 SKRIPSI*.
- Carmona, C. R. *et al.* (2019) ‘The Neutrophil-to-Lymphocyte Ratio is a Prognostic Biomarker of Amputation and Mortality Among Patients Suffering an Acute Limb Ischemia Event’, *European Journal of Vascular and Endovascular Surgery*, 58(6), p. e576. doi: 10.1016/j.ejvs.2019.09.045.
- Darwis, I. *et al.* (2020) ‘Pengaruh Nilai Neutrophil Lymphocyte Ratio (NLR) sebagai Prediktor Mortalitas pada Pasien Ulkus Diabetikum’, *Jurnal Kedokteran Universitas Lampung*, 4(2), pp. 128–134. Available at: <http://juke.kedokteran.unila.ac.id/index.php/JK/article/view/2875>.
- Fahmi, I. (2016) *Hubungan diabetes mellitus dan lama operasi terhadap infeksi luka operasi orthopaedi kriteria bersih di rsup haji adam malik medan*.
- Freisinger, E. *et al.* (2017) ‘Impact of diabetes on outcome in critical limb ischemia with tissue loss: a large-scaled routine data analysis.’, *Cardiovascular diabetology*, 16(1), p. 41. doi: 10.1186/s12933-017-0524-8.
- Fukuda, I. *et al.* (2015) ‘Acute limb ischemia: contemporary approach.’, *General thoracic and cardiovascular surgery*, 63(10), pp. 540–548. doi: 10.1007/s11748-015-0574-3.
- Gary, T., Pichler, M., Belaj, K., Hafner, F., Gerger, A., Froehlich, H., Eller, P., Pilger, E., Brodmann, M. 2013. Neutrophil-to-Lymphocyte Ratio and Its Association with Critical Limb Ischemia in PAOD Patients. *PLOS ONE*. Vol.8(2):e56745-e56745.
- Gunawan, H., Isnanta, R., Syafri, Z., Hasan, R. 2017. Iskemia Tungkai Akut. *Indonesian Journal Chest & Critical Care Medicine*. Vol.4(2): 14-22.
- Günay, A. E. and Ekici, M. (2021) ‘Relationship Between Neutrophil/Albumin Ratio and Early Mortality After Major Lower Extremity Amputation.’, *Cureus*, 13(9), p. e17733. doi: 10.7759/cureus.17733.
- Hess, C., Huang, Z., Patel, M., Baumgartener, I., Berger, J., Blomster, J. 2019. Acute Limb Ischemia in Peripheral Artery Disease. *Circulation*. 140:556–560.
- Howard, D., Banerjee, A., Fairhead, J., Hands, L., Silver, L., Rothwell, P. 2015. Population-Based Study of Incidence, Risk Factors, Outcome, and Prognosis of Ischemic Peripheral Arterial Events. *Circulation*. 132:1805-1815
- Howard, D. P. J. *et al.* (2015) ‘Population-Based Study of Incidence, Risk Factors, Outcome, and Prognosis of Ischemic Peripheral Arterial Events: Implications for Prevention.’, *Circulation*, 132(19), pp. 1805–1815. doi: 10.1161/CIRCULATIONAHA.115.016424.



Jaffery, Z., Thornton, S., White, C. 2011. Acute Limb Ischemia. *The American Journal of the Medical Sciences*. 342(3):226-234.

Kulezic, A. and Acosta, S. (2022) 'Epidemiology and Prognostic Factors in Acute Lower Limb Ischaemia: A Population Based Study', *European Journal of Vascular and Endovascular Surgery*, 63(2), pp. 296–303. doi: 10.1016/j.ejvs.2021.10.044.

Laksono, G.A.K., Erwin, F., Tahalele, P.L. 2020. Acute Limb Ischemia (ALI): An Overview of Clinical Diagnosis and Treatment. *Journal of Widya Medika Junior*. Vol. 2(2): 138-150.

Liao, K.-C. *et al.* (2013) 'The amputation and mortality rates of diabetic patients with critical limb ischemia: A nationwide population-based follow-up study in Taiwan', *Formosan Journal of Surgery*, 46(3), pp. 79–86. doi: <https://doi.org/10.1016/j.fjs.2013.05.002>.

Lim, S. *et al.* (2018) 'Epidemiology, treatment, and outcomes of acute limb ischemia in the pediatric population', *Journal of Vascular Surgery*, 68(1), pp. 182–188. doi: 10.1016/j.jvs.2017.11.064.

Luo, M. *et al.* (2015) 'Relationship between neutrophil-lymphocyte ratio and insulin resistance in newly diagnosed type 2 diabetes mellitus patients', *BMC Endocrine Disorders*, 15(1), p. 9. doi: 10.1186/s12902-015-0002-9.

Obara, H., Matsubara, K., Kitagawa, Y. 2018. Acute Limb Ischemia. *Ann Vasc Dis*. 11(4):443-448)

Olinic, D.-M. *et al.* (2019) 'Acute Limb Ischemia: An Update on Diagnosis and Management.', *Journal of clinical medicine*, 8(8). doi: 10.3390/jcm8081215.

Pierre-Louis, W. S., Bath, J., Mikkilineni, S., Scott, M. C., Harlander-Locke, M., Rasor, Z., & Smeds, M. (2019). Neutrophil to Lymphocyte Ratio as a Predictor of Outcomes after Amputation. *Annals of vascular surgery*, 54, 84–91. <https://doi.org/10.1016/j.avsg.2018.10.002>

Quedarusman, H., Lasut, P. 2019. Critical Limb Ischemia: Laporan Kasus. *Medical Scope Journal (MSJ)*. Vol.1(1):8-15.

Rutherford RB, Baker JD, Ernst C, *et al.* Recommended standards for reports dealing with lower extremity ischemia: revised version. *J Vasc Surg* 1997;26:517–38.

Santistevan, J. R. 2017. Acute Limb Ischemia: An Emergency Medicine Approach. *Emerg. Med. Clin. North Am.* 35, 889–909.

Simon, F., Oberhuber, A., Floros N., Busch, A., Wagenhauser, M., Schelzig, H., *et al.* 2018. Acute Limb Ischemia—Much More Than Just a Lack of Oxygen. *International Journal of Molecular Science*. 19:1-16

Shiny, A. *et al.* (2014) 'Association of neutrophil-lymphocyte ratio with glucose intolerance: an indicator of systemic inflammation in patients with type 2 diabetes.', *Diabetes technology & therapeutics*, 16(8), pp. 524–530. doi: 10.1089/dia.2013.0264.

Tasoglu, I., Sert, D., Colak, N., Uzun, A., Songur, M., Ecevit, A. 2014. Neutrophil–Lymphocyte Ratio and the Platelet–Lymphocyte Ratio Predict the Limb Survival in Critical Limb Ischemia. *Clinical and Applied Thrombosis/Hemostasis 2014*. Vol. 20(6): 645-600

Vatankhah, N. *et al.* (2017) 'Predictive value of neutrophil-to-lymphocyte ratio in diabetic wound healing', *Journal of Vascular Surgery*, 60(2), pp. 478–483. doi: 10.1016/j.jvs.2016.08.108.



Hubungan Antara Rasio Netrofil Limfosit (NLR), Diabetes Mellitus (DM) Dan Usia Dengan Lama Rawat (LOS), Infeksi Luka Operasi (ILO) Dan Mortalitas Pasien Acute Limb Ischemic (ALI) Paska Amputasi di RSUO Sardjito Tahun 2020

UNIVERSITAS
GADJAH MADA

KIKI RIZKIA, dr. R. Haryo Aribowo, Sp.B, Sp.BTKV (K) ; dr. Sumadi Lukman Anwar, M.Sc, Ph.D, Sp.B(K)Onk

Universitas Gadjah Mada, 2020. Diunduh dari <http://eip.repository.ugm.ac.id/>

Wang, Q. *et al.* (2017). 'Neutrophil-to-lymphocyte ratio is effective prognostic indicator for post-amputation patients with critical limb ischemia.', *Saudi medical journal*, 38(1), pp. 24–29. doi: 10.15537/smj.2017.1.15936.

Yesil, S., Akinci, B., Yener, S., Bayraktar, F., Karabay, O., Havitcioglu, H., Yapar, N., Atabey, A., Kucukyavas, Y., Comlekci, A., & Eraslan, S. (2009). Predictors of amputation in diabetics with foot ulcer: single center experience in a large Turkish cohort. *Hormones* (Athens, Greece), 8(4), 286–295. <https://doi.org/10.14310/horm.2002.1245>