

## DAFTAR PUSTAKA

Abbas, J., & Koni, P. (2005). The role of VCAM-1 in self-reactive T cell fate. *Clin. Immunol.*

Adamis, A. P., & Berman, A. J. (2008). Immunological mechanisms in the pathogenesis of diabeticretinopathy. *Seminars in immunopathology.*

Antonetti, D. A., Barber, A. J., Bronson, S. K., Freeman, W. M., Gardner, T. W., Jefferson, L. S., Kester, M., Kimball, S. R., Krady, J. K., & LaNoue, K. F. (2006). Diabetic retinopathy: seeing beyond glucose-induced microvascular disease. *Diabetes*, 55(9), 2401-2411.

Antonetti, D. A., Klein, R., & Gardner, T. W. (2012). Mechanisms of Disease Diabetic Retinopathy. *N Engl J Med*, 366, 1227-1239.

Armulik, A., Abramsson, A., & Betsholtz, C. (2005). Endothelial/pericyte interactions. *Circ. Res.*, 97(6), 512-523.

Atlas, I. D. (2020). 9th. In.

Azmy, R., Dawood, A., Kilany, A., El-Ghobashy, Y., Ellakwa, A. F., & El-Daly, M. (2012). Association analysis of genetic variations of eNOS and  $\alpha 2\beta 1$  integrin genes with type 2 diabetic retinopathy. *The appl. of Clin. Genet.*, 5, 55.

Banks, E., Brozinick Jr, J., Yaspelkis III, B., Kang, H., & Ivy, J. (1992). Muscle glucose transport, GLUT-4 content, and degree of exercise training in obese Zucker rats. *Am. J. Physiol. Endocrinol. Metab.*, 263(5), E1015-E1020.

Bird, S. R., & Hawley, J. A. (2017). Update on the effects of physical activity on insulin sensitivity in humans. *BMJ open sport & Exerc. Med.*, 2(1), e000143.

Broderick, T. L., Jankowski, M., & Gutkowska, J. (2017). The effects of exercise training and caloric restriction on the cardiac oxytocin natriuretic peptide system in the diabetic mouse. *Diabetes, metab. syndr. & obes.: targets and ther.*, 10, 27.

Cannata, F., Vadalà, G., Russo, F., Papalia, R., Napoli, N., & Pozzilli, P. (2020). Beneficial effects of physical activity in diabetic patients. *J. of Funct. Morphol. & Kinesiol.*, 5(3), 70.

Capitão, M., & Soares, R. (2016). Angiogenesis and inflammation crosstalk in diabetic retinopathy. *J. Cell. Biochem.*, 117(11), 2443-2453.

Cheung, A. K., Fung, M. K., Lo, A. C., Lam, T. T., So, K. F., Chung, S. S., & Chung, S. K. (2005). Aldose reductase deficiency prevents diabetes-induced blood-retinal barrier breakdown, apoptosis, and glial reactivation in the retina of db/db mice. *Diabetes*, 54(11), 3119-3125.

Chua, J., Lim, C. X. Y., Wong, T. Y., & Sabanayagam, C. (2018). Diabetic retinopathy in the Asia-Pacific. *The Asia-Pac. J. Ophthalmol.*, 7(1), 3-16.

Cortez, M., Torgan, C., Brozinick Jr, J., & Ivy, J. (1991). Insulin resistance of obese Zucker rats exercise trained at two different intensities. *Am. J. Physiol. Endocrinol. Metab.*, 261(5), E613-E619.

Dharmastuti, D., Agni, A., Widyaputri, F., Pawirotanu, S., Sofro, Z., Wardhana, F., Haryanto, S., Widayanti, T., Kotha, S., & Gupta, P. (2018). Associations of physical activity and sedentary behaviour with vision-threatening diabetic retinopathy in Indonesian population with type 2 diabetes mellitus: Jogjakarta Eye Diabetic Study in the Community (JOGED. COM). *Ophthalmic Epidemiol.*, 25(2), 113-119.

Ding, Y.-H., Young, C. N., Luan, X., Li, J., Rafols, J. A., Clark, J. C., McAllister, J. P., & Ding, Y. (2005). Exercise preconditioning ameliorates inflammatory injury in ischemic rats during reperfusion. *Acta Neuropathol. (Berl.)*, 109(3), 237-246.

Farsi, S., Azarbayjani, M., Hosseini, S., & Pirei, P. (2016). Response of Serum Levels of ICAM-1 VCAM-1 and CRP to High and Moderate Endurance Trainings in Sprague Dawley Male Diabetic Rats. *Armaghane danesh*, 21(8), 757-771.

Förstermann, U., & Li, H. (2011). Therapeutic effect of enhancing endothelial nitric oxide synthase (eNOS) expression and preventing eNOS uncoupling. *Br. J. Pharmacol.*, 164(2), 213-223.

Francois, M. E., & Little, J. P. (2015). Effectiveness and safety of high-intensity interval training in patients with type 2 diabetes. *Diabetes Spectr.*, 28(1), 39-44.

Gao, J., Pan, X., Li, G., Chatterjee, E., & Xiao, J. (2021). Physical exercise protects against endothelial dysfunction in cardiovascular and metabolic diseases. *J. Cardiovasc. Transl. Res.*, 1-17.

Gardner, T. W., Antonetti, D. A., Barber, A. J., LaNoue, K. F., Levison, S. W., & Group, P. S. R. R. (2002). Diabetic retinopathy: more than meets the eye. *Surv. Ophthalmol.*, 47, S253- S262.

Ghiasi, R., Soufi, F. G., hossein Somi, M., Mohaddes, G., Babil, F. M., Naderi, R., & Alipour, M. R. (2015). Swim training improves HOMA-IR in type 2 diabetes induced by high fat diet and low dose of streptozotocin in male rats. *Advanced pharmaceutical bulletin*, 5(3), 379.

Ghorbanzadeh, V., Mohammadi, M., Dariushnejad, H., Chodari, L., & Mohaddes, G. (2016). Effects of crocin and voluntary exercise, alone or combined, on heart VEGF-A and HOMA-IR of HFD/STZ induced type 2 diabetic rats. *J. Endocrinol. Invest.*, 39(10), 1179-1186.

Gologorsky, D., Thanos, A., & Vavvas, D. (2012). Therapeutic interventions against inflammatory and angiogenic mediators in proliferative diabetic retinopathy. *Mediators Inflamm.*, 2012.

Grijalva, J., Hicks, S., Zhao, X., Medikayala, S., Kaminski, P. M., Wolin, M. S., & Edwards, J. G. (2008). Exercise training enhanced myocardial endothelial nitric oxide synthase (eNOS) function in diabetic Goto-Kakizaki (GK) rats. *Cardiovasc. Diabetol.*, 7(1), 1-12.

Groves, J. T., & Wang, C. C. (2000). Nitric oxide synthase: models and mechanisms. *Curr. Opin. Chem. Biol. Curr Opin Chem Biol*, 4(6), 687-695.

Gu, H. F., Ma, J., Gu, K. T., & Brismar, K. (2013). Association of intercellular adhesion molecule 1 (ICAM1) with diabetes and diabetic nephropathy. *Front. Endocrinol. (Lausanne)*, 3, 179.

Gustavsson, C., Agardh, C.-D., Zetterqvist, A. V., Nilsson, J., Agardh, E., & Gomez, M. F. (2010). Vascular cellular adhesion molecule-1 (VCAM-1) expression in mice retinal vessels is affected by both hyperglycemia and hyperlipidemia. *PLoS one*, 5(9), e12699.

Guthrie, R. A., & Guthrie, D. W. (2004). Pathophysiology of diabetes mellitus. *Crit. Care Nurs. Q.*, 27(2), 113-125.

Hammes, H.-P., Feng, Y., Pfister, F., & Brownlee, M. (2011). Diabetic retinopathy: targeting vasoregression. *Diabetes*, 60(1), 9-16.

Hansen, D., Dendale, P., Jonkers, R., Beelen, M., Manders, R., Corluy, L., Mullens, A., Berger, J., Meeusen, R., & Van Loon, L. (2009). Continuous low-to moderate-intensity exercise training is as effective as moderate-to high-intensity exercise training at lowering blood HbA 1c in obese type 2 diabetes patients. *Diabetologia*, 52(9), 1789-1797.

Harreiter, J., & Roden, M. (2019). Diabetes mellitus—definition, klassifikation, diagnose, screening und prävention (update 2019). *Wien. Klin. Wochenschr.*, 131(1), 6-15.

Hejazi, S. M., Abrishami, L. H., Mohammad Khani, J., & Boghrabadi, V. (2013). The effects of 8-week aerobic exercises on serum levels of cell adhesion molecules among middle-aged women. *Adv. Stud. Biol.*, 5(6), 279-289.

Horstman, L. L., Jy, W., Jimenez, J. J., & Ahn, Y. S. (2004). Endothelial microparticles as markers of endothelial dysfunction. *Frontiers in Bioscience-Landmark*, 9(2), 1118-1135.

Iliaki, E., Poulaki, V., Mitsiades, N., Mitsiades, C. S., Miller, J. W., & Gragoudas, E. S. (2009). Role of  $\alpha 4$  integrin (CD49d) in the pathogenesis of diabetic retinopathy. *Invest. Ophthalmol. Vis. Sci.*, 50(10), 4898-4904.

Joussen, A. M., Poulaki, V., Mitsiades, N., Cai, W. y., Suzuma, I., Pak, J., Ju, S. T., Rook, S. L., Esser, P., & Mitsiades, C. (2003). Suppression of Fas-FasL-induced endothelial cell apoptosis prevents diabetic blood-retinal barrier breakdown in a model of streptozotocin-induced diabetes. *The FASEB Journal*, 17(1), 76-78.

Kargarfard, M., Lam, E. T., Shariat, A., Asle Mohammadi, M., Afrasiabi, S., Shaw, I., & Shaw, B. S. (2016). Effects of endurance and high intensity training on ICAM-1 and VCAM-1 levels and arterial pressure in obese and normal weight adolescents. *Phys. Sportsmed.*, 44(3), 208-216.

Kaul, K., Tarr, J. M., Ahmad, S. I., Kohner, E. M., & Chibber, R. (2013). Introduction to diabetes mellitus. *Diabetes*, 1-11.

Khademi, Y., Azarbayejani, M., & Hossini, S. (2016). The impact of high intensity aerobic interval training (HIIT) and flaxseed oil on ICAM-1 gene expression in heart tissue in male Wistar rats. *Armaghane danesh*, 21(9), 873-886.

Khalifaoui, T., Lizard, G., & Ouertani-Meddeb, A. (2008). Adhesion molecules (ICAM-1 and VCAM-1) and diabetic retinopathy in type 2 diabetes. *J. Mol. Histol.*, 39(2), 243-249.

Killingley, J. (2010). Diabetic Retinopathy: Examining Pericyte-Endothelial Ratios Using Zebrafish and Rat Retinas. *Submitted to the Coll. of Arts & Sci. Am. Univ.*

Kim, S. Y., Johnson, M. A., McLeod, D. S., Alexander, T., Hansen, B. C., & Luty, G. A. (2005). Neutrophils are associated with capillary closure in spontaneously diabetic monkey retinas. *Diabetes*, 54(5), 1534-1542.

Koch, A. E., Halloran, M. M., Haskell, C. J., Shah, M. R., & Polverini, P. J. (1995). Angiogenesis mediated by soluble forms of E-selectin and vascular cell adhesion molecule-1. *Nat.*, 376(6540), 517-519.

Koh, Y., & Park, J. (2018). Cell adhesion molecules and exercise. *J. inflamm. Res.*, 11, 297.

Kowluru, R. A. (2020). Retinopathy in a diet-induced type 2 diabetic rat model and role of epigenetic modifications. *Diabetes*, 69(4), 689-698.

Koya, D., & King, G. L. (1998). Protein kinase C activation and the development of diabetic complications. *Diabetes*, 47(6), 859-866.

Liu, L., & Kubes, P. (2006). Chemokines in leukocyte transendothelial migration. In *Chemokine Biol.—Basic Res. & Clin. Appl.* (pp. 109-122). Springer.

Mastej, K., & Adamiec, R. (2008). Neutrophil surface expression of CD11b and CD62L in diabetic microangiopathy. *Acta diabetologica*, 45(3), 183-190.

Mayhan, W. G., Arrick, D. M., Patel, K. P., & Sun, H. (2011). Exercise training normalizes impaired NOS-dependent responses of cerebral arterioles in type 1 diabetic rats. *Am. J. Physiol. Heart Circ. Physiol.*, 300(3), H1013-H1020.

McLeod, D. S., Lefer, D. J., Merges, C., & Luty, G. A. (1995). Enhanced expression of intracellular adhesion molecule-1 and P-selectin in the diabetic human retina and choroid. *Am. J. Pathol.*, 147(3), 642.

Miyamoto, K., Khosrof, S., Bursell, S.-E., Rohan, R., Murata, T., Clermont, A. C., Aiello, L. P., Ogura, Y., & Adamis, A. P. (1999). Prevention of leukostasis and vascular leakage in streptozotocin-induced diabetic retinopathy via intercellular adhesion molecule-1 inhibition. *Proc. Natl. Acad. Sci.*, 96(19), 10836-10841.

Mohammad, P., Esfandiar, K. Z., Abbas, S., & Ahoora, R. (2019). Effects of moderate-intensity continuous training and high-intensity interval training on serum levels of resistin, chemerin and liver enzymes in streptozotocin-nicotinamide induced type-2 diabetic rats. *J. diab. & metab. disord.*, 18(2), 379-387.

Mrugacz, M., Bryl, A., & Zorena, K. (2021). Retinal Vascular Endothelial Cell Dysfunction and Neuroretinal Degeneration in Diabetic Patients. *J. Clin. Med.*, 10(3), 458.

Muniyappa, R., Iantorno, M., & Quon, M. J. (2008). An integrated view of insulin resistance and endothelial dysfunction. *Endocrinol. Metab. Clin. N. Am.*, 37(3), 685-711.

Nagareddy, P. R., Xia, Z., McNeill, J. H., & MacLeod, K. M. (2005). Increased expression of iNOS is associated with endothelial dysfunction and impaired pressor responsiveness in streptozotocin-induced diabetes. *Am. J. Physiol. Heart Circ. Physiol.*, 289(5), H2144- H2152.

Noda, K., Nakao, S., Ishida, S., & Ishibashi, T. (2012). Leukocyte adhesion molecules in diabetic retinopathy. *J. Ophthalmol.*, 2012.

Olivares, A. M., Althoff, K., Chen, G. F., Wu, S., Morrisson, M. A., DeAngelis, M. M., & Haider, N. (2017). Animal models of diabetic retinopathy. *Curr. Diab. Rep.*, 17(10), 1-17.

Opatrikova, R., Kubatka, P., Caprnda, M., Büsselberg, D., Krasnik, V., Vesely, P., Saxena, S., Ruia, S., Mozos, I., & Rodrigo, L. (2018). Nitric oxide in the pathophysiology of retinopathy: evidences from preclinical and clinical researches. *Acta Ophthalmol.*, 96(3), 222-231.

Pescatello, L. S., Franklin, B. A., Fagard, R., Farquhar, W. B., Kelley, G. A., & Ray, C. A. (2004). American College of Sports Medicine position stand. Exercise and hypertension. *Med. Sci. Sports Exerc.* 36(3), 533-553.

Pieper, G. (1999). Enhanced, unaltered and impaired nitric oxide-mediated endothelium-dependent relaxation in experimental diabetes mellitus: importance of disease duration. *Diabetologia*, 42(2), 204-213.

Pouranfar, S., Azarbayjani, M. A., & Abedi, B. (2019). The Effects of Resistance Training on VCAM-1, ICAM-1 and CRP in Diabetic Rats. *Iran. J. Diabetes Obes.*, 11(1), 38-45.

Praidou, A., Harris, M., Niakas, D., & Labiris, G. (2017). Physical activity and its correlation to diabetic retinopathy. *J. Diab. Complicat.*, 31(2), 456-461.

Qian, Y., Li, S., Ye, S., Chen, Y., Zhai, Z., Chen, K., & Yang, G. (2008). Renoprotective effect of rosiglitazone through the suppression of renal intercellular adhesion molecule-1 expression in streptozotocin-induced diabetic rats. *J. Endocrinol. Invest.*, 31(12), 1069-1074.

Rask-Madsen, C., & King, G. L. (2013). Vascular complications of diabetes: mechanisms of injury and protective factors. *Cell Metab.*, 17(1), 20-33.

Ren, C., Liu, W., Li, J., Cao, Y., Xu, J., & Lu, P. (2019). Physical activity and risk of diabetic retinopathy: a systematic review and meta-analysis. *Acta Diabetol.*, 56(8), 823-837.

Richter, E. A., & Hargreaves, M. (2013). Exercise, GLUT4, and skeletal muscle glucose uptake. *Physiol. Rev.*

Roberts, C. K., Little, J. P., & Thyfault, J. P. (2013). Modification of insulin sensitivity and glycemic control by activity and exercise. *Med Sci Sports Exerc.*, 45(10), 1868-1877.

Rocha, B., Rodrigues, A., Tomada, I., Martins, M., Guimaraes, J., Gouveia, A., Almeida, H., & Neves, D. (2018). Energy restriction, exercise and atorvastatin treatment improve endothelial dysfunction and inhibit miRNA-155 in the erectile tissue of the aged rat. *Nutr. Metab.*, 15(1), 1-12.



Sasongko, M. B., Wardhana, F. S., Febryanto, G. A., Agni, A. N., Supanji, S., Indrayanti, S. R., Widayanti, T. W., Widyaputri, F., Widhasari, I. A., & Lestari, Y. D. (2020). The estimated healthcare cost of diabetic retinopathy in Indonesia and its projection for 2025. *Br. J. Ophthalmol.* 104(4), 487-492.

Sasongko, M. B., Widyaputri, F., Agni, A. N., Wardhana, F. S., Kotha, S., Gupta, P., Widayanti, T. W., Haryanto, S., Widyaningrum, R., & Wong, T. Y. (2017). Prevalence of diabetic retinopathy and blindness in Indonesian adults with type 2 diabetes. *Am. J. Ophthalmol.*, 181, 79-87.

Savoia, C., & Schiffrin, E. L. (2007). Vascular inflammation in hypertension and diabetes: molecular mechanisms and therapeutic interventions. *Clin. Sci.*, 112(7), 375-384.

Schalkwijk, C. G., & Stehouwer, C. D. (2005). Vascular complications in diabetes mellitus: the role of endothelial dysfunction. *Clin. Sci.*, 109(2), 143-159.

Soori, R., Ranjbar, K., Salehian, O., & Eslam Doost, M. (2013). The effect of high interval intensity training on Plasma Soluble Intercellular Adhesion Molecule-1 (sICAM-1) in sedentary obese male. *J. Sport Biosci.*, 5(3), 91-102.

Srinivasan, K., Viswanad, B., Asrat, L., Kaul, C., & Ramarao, P. (2005). Combination of high-fat diet-fed and low-dose streptozotocin-treated rat: a model for type 2 diabetes and pharmacological screening. *Pharmacol. Res.*, 52(4), 313-320.

Stevens, H. Y., Melchior, B., Bell, K. S., Yun, S., Yeh, J.-C., & Frangos, J. A. (2008). PECAM-1 is a critical mediator of atherosclerosis. *Dis. Model. Mech.*, 1(2-3), 175-181.

Sun, Y., & Smith, L. E. (2018). [WITHDRAWN] Retinal Vasculature in Development and Diseases. *Annu. Rev. Vis. Sci.*, 4, 101-122.

Tarr, J. M., Kaul, K., Wolanska, K., Kohner, E. M., & Chibber, R. (2013). Retinopathy in diabetes. *Diabetes*, 88-106.

Terada, S., Yokozeki, T., Kawanaka, K., Ogawa, K., Higuchi, M., Ezaki, O., & Tabata, I. (2001). Effects of high-intensity swimming training on GLUT-4 and glucose transport activity in rat skeletal muscle. *J. Appl. Physiol.*, 90(6), 2019-2024.

Ugurlu, N., Gerceker, S., Yülek, F., Ugurlu, B., Sari, C., Baran, P., & Çagil, N. (2013). The levels of the circulating cellular adhesion molecules ICAM-1, VCAM-1 and endothelin-1 and the flow-mediated vasodilatation values in patients with type 1 diabetes mellitus with early-stage diabetic retinopathy. *Intern. Med.*, 52(19), 2173-2178.

Van Hecke, M., Dekker, J., Nijpels, G., Moll, A., Heine, R., Bouter, L., Polak, B., & Stehouwer, C. (2005). Inflammation and endothelial dysfunction are associated with retinopathy: the Hoorn Study. *Diabetologia*, 48(7), 1300-1306.

Verma, A., Han, P.-Y., Nakagawa, T., Johnson, R., Grant, M., Campbell-Thompson, M., Atkinson, M., Segal, M., Hauswirth, W., & Li, Q. (2009). Diabetic eNOS Knockout Mice Develop Accelerated Retinopathy. *Invest. Ophthalmol. Vis. Sci.*, 50(13), 5908-5908.



Victor, A. A., Gondhowiardjo, T. D., Dharma, R., Jusman, S. W., Yandri, V. R., Yuneta, R., & Yuriza, R. E. (2019). Intercellular adhesive molecule-1 (ICAM-1) in proliferative diabeticretinopathy. *Asian J. Ophthalmol.*, 17(1), 120-127.

Wahyu, T., & Syumarti, M. The Epidemiology of Diabetic Retinopathy.

Wang, J., Polaki, V., Chen, S., & Bihl, J. C. (2020). Exercise improves endothelial function associated with alleviated inflammation and oxidative stress of perivascular adipose tissuein type 2 diabetic mice. *Oxid. Med. Cell. Longev.*, 2020.

Wang, S., Li, J., Zhang, C., Xu, G., Tang, Z., Zhang, Z., Liu, Y., & Wang, Z. (2019). Effects of aerobic exercise on the expressions and activities of nitric oxide synthases in the blood vessel endothelium in prediabetes mellitus. *Exp. Ther. Med.*, 17(5), 4205-4212.

Widlansky, M. E., Gokce, N., Keaney, J. F., & Vita, J. A. (2003). The clinical implications of endothelial dysfunction. *J. Am. Coll. Cardiol.*, 42(7), 1149-1160.

Xu, H., Chen, M., & Forrester, J. V. (2009). Para-inflammation in the aging retina. *Prog. Retin. EyeRes.*, 28(5), 348-368.

Yoo, T. K., Oh, B. K., Lee, M. Y., & Sung, K.-C. (2022). Association between physical activity andinsulin resistance using the homeostatic model assessment for insulin resistance independent of waist circumference. *Sci. Rep.*, 12(1), 1-9.