

DAFTAR PUSTAKA

- Abdelgaleil, S.A.M., Doe, M., Morimoto, Y., Nakatani, M. (2006). Rings B,D-seco limonoids from the leaves of *Swietenia mahogani*. *Phytochemistry*, 67(5), 452–458.
- Acamovic, T., Brooker, J.D. (2005). Biochemistry of plant secondary metabolites and their effects in animals. *Proceedings of the Nutrition Society*, 64, 403–412.
- Aguilera, A., Zamora, R. (2009). Surface roughness in sapwood and heartwood of Blackwood (*Acacia melanoxylon* R. Br.) machined in 90-0 direction. *European Journal of Wood and Wood Product*, 67, 297–301. DOI: 10.1007/s00107-009-0308-2.
- Alasalvar, C., Grigor, J.M., Zhang, D.L., Quantick, P.C., Shahidi, F. (2001). Comparison of volatiles, phenolics, sugars, antioxidant vitamins, and sensory quality of different colored carrot varieties. *Journal of Agricultural and Food Chemistry*, 49, 1410–1416.
- Ali, H.M., EL-Mahrouk, E.S.M., Hassan, F.A., EL-Tarawy, M.A. (2011). Usage of sewage effluent in irrigation of some woody tree seedlings. Part 3: *Swietenia mahagoni* (L.) Jacq. *Saudi Journal of Biological Sciences*, 18, 201–207. DOI: 10.1016/j.sjbs.2010.08.001.
- Anderson, A.B., Riffer, R., Wong, A. (1968). Chemistry of the genus *Sequoia*. VI. On the cyclitols present in heartwood of *Sequoia sempervirens*. *Phytochemistry*, 7, 1867–1870. DOI: 10.1016/S0031-9422(00)86661-3.
- Anish, M.C., Anoop, E.V., Vishnu, R., Sreejith, B., Jijeesh, C.M. (2015). Effect of growth rate on wood quality of teak (*Tectona grandis* L. f.): a comparative study of teak grown under differing site quality conditions. *Journal of the Indian Academy of Wood Science*, 12, 81-88. DOI: 10.1007/s13196-015-0147-1.
- Anjos, O., Pereira, H., Rosa, M.E. (2010). Tensile properties of cork in the tangential direction: variation with quality, porosity, density and radial position in the cork plank. *Materials & Design*, 4(31), 2085–2090.
- Anjos, O., Rodrigues, C., Morais, J., Pereira, H. (2014). Effect of density on the compression behaviour of cork. *Materials & Design*, 53, 1089–1096.
- Anon. (1988). Dry kiln operator's manual. USDA Forest Service, Forest Products Lab., Hardwood Research Council, Memphis, TN, USA, 6.
- Anoop, E.V., Jijeesh, C.M., Sindhumathi, C.R., Jayasree, C.E. (2014). Wood physical, anatomical and mechanical properties of Big Leaf Mahogany (*Swietenia macrophylla* Roxb) a potential exotic for South India. *Research Journal of Agriculture and Forestry Sciences*, 2(8), 7-13.
- Antony, F., Schimleck, L.R., Daniels, R.F. (2012). Identification of representative sampling heights for specific gravity and moisture content in plantation-grown loblolly pine (*Pinus taeda*). *Canadian Journal of Forest Research*, 42, 574– 584. DOI:10.1139/x2012-009.

- Aragao, V.P.M., Navarro, B.V., Passamani, L.Z., Macedo, A.F., Floh, E.S., Silveira, V., Santa-Catarina, C. (2015). Free amino acids, polyamines, soluble sugars and proteins during seed germination and early seedling growth of *Cedrela fissilis* Vellozo (Meliaceae), an endangered hardwood species from the Atlantic Forest in Brazil. *Theoretical and Experimental Plant Physiology*, 27, 157–169. DOI: 10.1007/s40626-015-0041-7.
- Arisandi, R., Ashitani, T., Takahashi, K., Marsoem, S.N., Lukmandaru, G. (2020). Lipophilic extractives of the wood and bark from *Eucalyptus pellita* F. Muell grown in Merauke, Indonesia. *Journal of Wood Chemistry and Technology*, 40(2), 146–154.
- Arisandi, R., Marsoem, S.N., Lukmandaru, G., Ashitani, T., Takahashi, K. (2019a). The contents of phenolics and cell wall component of *Eucalyptus pellita* F. Muell stemwood and bark. *Wood Research*, 64(3), 411–422.
- Arisandi, R., Masendra, Purba, B.A.V., Wati, F.Z., Ihda, F.V., Sumantri, F., Lukmandaru, G. (2019b). Lipophilic extractives of mahogany (*Swietenia macrophylla* King) barks. – Proceedings of 9th International Symposium of Indonesian Wood Research Society, Indonesia, Sept. 2017. Denpasar, 192–201.
- Ayobi, E., Kiaei, M., Bakhshi, R. (2011). Heartwood and sapwood properties of *Quercus castaneaefolia* in the Iranian forests. *Middle East Journal Scientific Research*, 8(3), 669-673.
- Babbar, N., Oberoi, H.S., Sandhu, S.K., Bhargav, V.K. (2014) Influence of different solvents in extraction of phenolic compounds from vegetable residues and their evaluation as natural sources of antioxidants. *Journal of Food Science and Technology*, 51, 2568–2575.
- Baeza, J., Freer, J. (2001). Chemical characterization of wood and its components. Dalam : Wood and Cellulosic Chemistry .D. N. S. Hon and N. Shiraishi (Ed). New York: Marcel Dekker.
- Bajpai, P. (2018). Hardwood anatomy. In: Biermann's Handbook of Pulp and Paper: raw material and pulp making 3 Eds. Doi: 10.1016/B978-0-12-814240-0.00005-7.
- Bamber, R.K. (1976). Heartwood, its function and formation. *Wood Science and Tecnology*, 10, 1-8. DOI: 10.1007/BF00376379.
- Bamber, R.K., Fukazawa, K. (1985). Sapwood and heartwood: A review. *Forestry Abstract*, 46, 567-580.
- Baqui, S.A., Shah, J.J. (1985). Histoenzymatic studies in wood of *Acacia auriculiformis* during heartwood formation. *Holzforschung*, 39(6), 311-320.
- Basri, E.V., Jijeesh, C.M., Sindhumathi, C.R., Jayasree, C.E. (2014). Wood physical, anatomical and mechanical properties of Big Leaf Mahogany (*Swietenia macrophylla* Roxb) a potential exotic for South India. *Research Journal of Agriculture and Forestry Science*, 2, 7-13.
- Basri, E., Wahyudi, I. (2012). Wood basic properties of Jati Plus Perhutani from different ages and their relationship to drying properties and qualities. *Journal of Forest Product Research* 31:93-102.

- Batubara, I., Darusman, L.K., Mitsunaga, T., Rahminiwati, M., Djauhari, E. (2010). Potency of Indonesian medicinal plants as tyrosinase inhibitor and antioxidant agent. *Journal of Biological Sciences*, 10(2), 138–144.
- Batubara, I., Kotsuka, S., Yamauchi, K., Kuspradini, H., Mitsunaga, T., Darusman, L.K. (2012). TNF- α production inhibitory activity, phenolic, flavonoid, and tannin contents of selected Indonesian medicinal plants. *Research Journal of Medicinal Plants*, 6(6), 406–415.
- Batubara, I., Mitsunaga, T., Ohashi, H. (2009). Screening antiacne potency of Indonesian medicinal plants: antibacterial, lipase inhibition, and antioxidant activities. *Journal of Wood Science*, 55, 230–235.
- Beauchamp, K. (2011) The biology of heartwood formation in Sitka spruce and Scots pine. Dissertation, The University of Edinburgh.
- Belt, T., Altgen, M., Mäkelä, M., Hänninen, T., Rautkari, L. 2019. Cellular level chemical changes in Scots pine heartwood during incipient brown rot decay. *Scientific Report*, 9, 1-10. DOI: 10.1038/s41598-019-41735-8.
- Bergström, B. (2003). Chemical and structural changes during heartwood formation in *Pinus sylvestris*. *Forestry*, 76, 45–53. DOI: 10.1093/forestry/76.1.45.
- Berrocal, A., Baeza, J., Rodríguez, J., Espinosa, M., Freer, J. (2004). Effect of tree age on variation of *Pinus radiata* D. Don chemical composition. *Journal of the Chilean Chemical Society*, 49(3), 251–256.
- Berrocal, A., Gaitan-Alvarez, J., Moya, R., Fernandez-Solis, D., Ortiz-Malavassi, E. (2020). Development of heartwood, sapwood, bark, pith and specific gravity of teak (*Tectona grandis*) in fast-growing plantations in Costa Rica. *Journal of Forestry Research*, 36, 667-676.
- Bhat, K.M., Bhat, K.V., Dhamodaran, T.K. (1987). Effect of age and location on pulpwood quality of *Eucalyptus grandis*. Kerala Forest Research Institute Peechi, Thrissur, India.
- Bhat, K.M., Florence, E.J.M. (2003). Natural decay resistance of juvenile teak wood grown in high input plantations. *Holzforschung*, 57, 453-455.
- Bikovens, O., Roze, L., Pranovich, A., Reunanen, M., Telysheva, G. (2013). Chemical composition of lipophilic extractives from grey alder (*Alnus incana*). *BioResources*, 8(1), 350–357.
- Björklund, L. (1999). Identifying heartwood-rich stands or stems of *Pinus sylvestris* by using inventory data. *Silva Fennica*, 33(2), 119–129.
- BPS. (2019). Statistik produksi kehutanan. Badan Pusat Statistik, Jakarta.
- BPS. (2022). Badan pusat statistik kabupaten Gunungkidul. <https://gunungkidulkab.bps.go.id/indicator/151/55/1/curah-hujan.html> (Akses 1 Juli 2022).
- Bradbury, G.J. (2005). A comparison of timber quality of blackwood grown in young swamp forest, fenced regeneration, and a plantation. *Tasforests* 16:95–110.
- Bradbury, G.J., Potts, B.M., Beadle, C.J. (2011). Genetic and environmental variation in wood properties of *Acacia melanoxylon*. *Annals of Forest Science*, 68, 1363–1373.

- Brix, H., Mitchell, A.K. (1983). Thinning and nitrogen fertilization effects of sapwood development and relationships of foliage quantity of sapwood area and basal area in Douglas-fir. *Canadian Journal of Forest Research*, 13, 384–389.
- Briscoe, C.B., Harris, J.B., Wyckoff F, (1963). Variation of specific gravity in plantation-grown trees of big-leaf mahogany. *Caribbean Forestry*, 24, 67-74.
- Burgert, I., Cabane, E., Zollfrank, C., Berglund, L. (2015). Bio-inspired functional wood-based materials – hybrids and replicates. *International Materials Reviews*, 60(8), 431-450.
- Burtin, P., Jay-Allemand, C., Charpentier, J.-P., Janin, G. (1998). Natural wood colouring process in *Juglans* sp. (*J. nigra*, *J. regia* and hybrid *J. nigra* 23 × *J. regia*) depends on native phenolic compounds accumulated in the transition zone between sapwood and heartwood. *Trees*, 12, 258–264.
- Bush, D., McCarthy, K., Meder, R. (2011). Genetic variation of natural durability traits in *Eucalyptus cladocalyx* (sugar gum). *Annals of Forest Science*, 68, 1057–1066
- Cambie, R.C., Ser, N.G.A., Kokubun, T. (1997). Heartwood constituents of *Planchonella vitiensis*. *Biochemical Systematics and Ecology*, 25, 677–678.
- Caron, A., Altaner C.M., Gardiner, B., Jarvis, M.C. (2013). Distribution of extractives in Sitka spruce (*Picea sitchensis*) grown in the northern UK. *European Journal of Wood and Wood Products*, 71(6), 697–704.
- Carrodus BB (1971) Carbon dioxide and the formation of heartwood. *New Phytol.* 70, 939-943.
- Celedon, J.M., Chiang, A., Yuen, M.M.S., Diaz-Chavez, M.L., Madilao, L.L., Finnegan, P.M., Barbour, E.L., Bohlmann, J. (2016). Heartwood-specific transcriptome and metabolite signatures of tropical sandalwood (*Santalum album*) reveal the final step of (Z)- santalol fragrance biosynthesis. *The Plant Journal*, 86, 289-299. DOI: 10.1111/tpj.13162.
- Chattaway, Y.M.M. (1949). The development of tyloses and secretion of gums in heartwood formation. *Australian Journal of Scientific Research B2*, 227-240.
- Chave, J., D. Coomes, S. Jansen, S. L. Lewis, N. G. Swenson, and A. E. Zanne. (2009). Towards a worldwide wood economics spectrum. *Ecology Letters* 12, 351–366.
- Chen, Y.-Y., Wang, X.-N., Fan, C.-Q., Yin, S., Yue, J.-M. (2007). Swiemahogins A and B, two novel limonoids from *Swietenia mahagoni*. *Tetrahedron Letters*, 48, 7480–7484.
- Chitra, V., Senthil, V.K.N., Rahul, V.A., Usha, S., Jeeva, G.R. A review on veppampattai chooranam – A siddha single herbal formulation. *World Journal of Pharmaceutical Research*, 6, 366-372.
- Chiou, T.J., Bush, D.R. (1998). Sucrose is a signal molecule in assimilate partitioning. *Proceedings of the National Academy of Sciences of the United States of America*, 95, 4784–4788.

- Chowdhury, M.Q., Ishiguri, F., Iizuka, K., Taka-segarhima, Y., Matsumoto, K., Hiraiwa, T., Ishido, M., Sanpe, H., Yokota, S., Yoshizawa, N. (2009) Radial variations of wood properties in *Casuarina equisetifolia* growing in Bangladesh. *Journal of Wood Science*, 55, 139-143. DOI: 10.1007/s10086-008-1004-2.
- Climent, J., Chambel, M.R., Gil, L., Pardos, J. (2003). Vertical heartwood variation patterns and prediction of heartwood volume in *Pinus canariensis* Sm. *Forest Ecology and Management*, 174, 203–211.
- Climent, J., Gil, L., Pardos, J. (1993). Heartwood and sapwood development and its relationship to growth and environment in *Pinus canariensis* Chr. Sm. ex DC. *Forest Ecology and Management*, 59, 165–174.
- Climent, J., Chambel, M.R., Perez, E., Gil, L., Pardos, J. (2002). Relationship between heartwood radius and early radial growth, tree age, and climate in *Pinus canariensis*. *Canadian Journal of Forest Research*, 32, 103-111.
- Cown, D.J. (1981). Wood density of *Pinus caribaea* Var. *Hondurensis* grown in Fiji. *New Zealand Journal of Forestry Science*, 11(3), 244-53.
- Crespo, R., Jimenez, E., Suatunace, P., Law, G. Sanchez, C. (2008). Comparative analysis of physicalmechanical properties of teak (*Tectona grandis* L.F.) from Quevedo and Baltazar. *Ciencia y Tecnologia*, 1, 55–63 (in Spanish).
- Cui, Z., Li, X., Xu, D., Yang, Z. (2020). Changes in non-structural carbohydrates, wood properties and essential oil during chemically-induced heartwood formation in *Dalbergia odorifera*. *Frontiers in Plant Science*, 11, 1-13. DOI: 10.3389/fpls.2020.01161.
- Cuvilas, C.A. (2009). Characterization of available and potential sources of wood fuels in Mozambique. MSc thesis, Swedish University of Agricultural Sciences, Uppsala.
- da Silva, A.R., Pastore, T.C.M., Braga, J.W.B., Davrieux, F., Okino, E.Y.A., Coradin, V.T.R., Camargos, J.A.A., do Prado, A.G.S. (2013). Assessment of total phenols and extractives of mahogany wood by near infrared spectroscopy (NIRS). *Holzforschung*, 67, 1–8.
- Datta, S.K., Kumar, A. (1987). Histochemical studies of the transition from sapwood to heartwood in *Tectona grandis*. *IAWA Bulletin* ns, 8, 363-368. DOI: 10.1163/22941932-90000456.
- Debell, J.D., Lachenbruch, B. (2009). Heartwood/sapwood variation of western redcedar as influenced by cultural treatments and position in tree. *Forest Ecology and Management*, 258, 2026-2032. DOI: 10.1016/j.foreco.2009.07.054.
- Déjardin, A.C., Rochat, S., Maugenest, J.P., Boutin. (1997). Purification, characterisation and physiological role of sucrose synthase in the pee seed coat (*Pisum sativum* L.). *Planta*, 201, 128-137.
- De Schepper, V., De Swaef, T., Bauweraerts, I., Steppe, K. (2013). Phloem transport: a review of mechanisms and controls. *Journal of Experimental Botany*, 64, 4839–4850. DOI: 10.1093/jxb/ert302.
- Dietrichs, H.H. (1965). Das Verhalten von Kohlenhydraten bei der Holzverkernung. *Holzforschung*, 18, 14–24. DOI: 10.1515/hfsg.1964.18.1-2.14.

- DLHK. (2017). Budidaya mahoni (*Swietenia macrophylla*). Balai Pengelolaan Hutan Wilayah Lebak dan Tangerang. Dinas Lingkungan Hidup dan Kehutanan Provinsi Banten, Banten.
- Downes, G.M., Hudson, I.L., Raymond, C.A., Dean, G.H., Michell, A.J., Schimleck, L.S., Evans, R., Muneri, A. (1997). Sampling plantation eucalypts for wood and fibre properties. CSIRO Publishing, Melbourne.
- DuBois, M., Gilles, K.A., Hamilton, J.K., Rebers, P.A., Smith, F. (1956). Colorimetric method for determination of sugars and related substances. *Analytical Chemistry*, 28(3), 350–356.
- Dulsalam., Tinambunan, D., Sumantri, I., Sinaga, M. (2000). Peningkatan efisiensi pemungutan kayu sebagai bahan baku industri. Prosiding Lokakarya Penelitian Hasil Hutan tanggal 7 Desember 2000. Pusat Penelitian dan Pengembangan Hasil Hutan, Bogor.
- Dünisch, O., Richter, H.-G., Koch, G. (2010). Wood properties of juvenile and mature heartwood in *Robinia pseudoacacia* L. *Wood Science and Technology*, 44, 301–313.
- Edreva, A., Velikova, V., Tsonev, T., Dagnon, S., Gürel, A.L., Aktas, L. (2008) Stress-protective role of secondary metabolites: Diversity of functions and mechanisms. *General and Applied Plant Physiology*, 34, 67–78.
- Ekimoto, H., Irie, Y., Araki, Y., Han, G.Q., Kadota, S., Kikuchi, T. (1991). Platelet aggregation inhibitors from the seeds of *Swietenia mahagoni*: inhibition of in vitro and in vivo platelet-activating factor-induced effects of tetranortriterpenoids related to swietenine and swietenolide. *Planta Medica*, 57(1), 56–58.
- Ericsson, T., Fries, A. (1999). High heritability for heartwood in north Swedish Scots pine. *Theoretical and Applied Genetics*, 98, 732–735.
- Falah, S., Suzuki, T., Katayama, T. (2008). Chemical constituents from *Swietenia macrophylla* bark and their antioxidant activity. *Pakistan Journal of Biological Sciences*, 11(16), 2007–2012.
- Fengel, D., Wegener, G. (1989). Wood: Chemistry, Ultrastructure, Reactions. Berlin, Walter de Gruyter.
- Fernández-Sólis, D., Berrocal, A., Moya, R. (2018). Heartwood formation and prediction of heartwood parameters in *Tectona grandis* L.f. trees growing in forest plantations in Costa Rica. *Bois et Forêts des Tropiques*, 335, 25–37. DOI: 10.19182/bft2018.335.a31499.
- Francis, J.K. (2003). Wood densities of mahagonies in Puerto Rican plantation. *Ecological Studies*, 159, 358–361.
- Freire, C.S.R., Silvestre, J.D., Neto, C.P. (2005) Lipophilic extractives in Eucalyptus globulus kraft pulps. Behaviour during ECF bleaching. *Journal of Wood Chemistry and Technology*, 25, 67–80.
- Freitas, T.P., Oliveira, J.T.S., Paes, J.B., Vidaurre, GB, Lima, JL. (2019). Environmental effect on growth and characteristics of *Eucalyptus* wood. *Floresta e Ambiente*, 26(4), 1–9.
- Fuji, T., Marsoem, S.N., Fujiwara, T. (2000). Annual growth rings in Mahogany (*Swietenia macrophylla*) growing in Central Java. In: Kim YS (ed) New horizon in wood anatomy. Chonnam National University Press, Kwangju.

- Fukazawa, K. (1984). Juvenile wood of hardwoods judged by density variation. *IAWA Bulletin*, 5, 65–73. DOI:10.1163/22941932-90000861.
- Fukazawa, K., Yamamoto, K., Ishida, S. (1980). The season of heartwood formation in genus *Pinus*. Mitt. Bundesforschungsanst. *Forst- und Holzwirtschaft*, 131, 113-131.
- Furze, M.E., Trumbore, S., Hartmann, H. (2018). Detours on the phloem sugar highway: stem carbon storage and remobilization. *Current Opinion in Plant Biology*, 43, 89–95. DOI: 10.1016/j.pbi.2018.02.005
- Gala, S., Sumarno, S., Mahfud, M. (2020). Comparison of microwave and conventional extraction methods for natural dyes in wood waste of mahogany (*Swietenia mahagoni*). *Journal of Applied Engineering Science*, 18(4), 618–623.
- Gominho, J., Pereira, H. (2000). Variability of heartwood content in plantation grown *Eucalyptus globulus* Labill. *Wood and Fiber Science*, 32, 189-195.
- Gominho, J., Figueira, J., Rodríguez, J.C., Pereira, H. (2001). Within-tree variation of heartwood, extractives and wood density in the eucalypt hybrid *urograndis* (*Eucalyptus grandis* × *E. urophylla*). *Wood and Fiber Science*, 33, 3-8.
- Gominho, J., Pereira, H. (2005) The influence of tree spacing in heartwood content in *Eucalyptus globulus* labill. *Wood and Fiber Science*, 37, 582 – 590.
- Gominho, J., Lourenço, A., Miranda, I., Pereira, H. (2015). Radial and axial variation of heartwood properties and extractives in mature trees of *Eucalyptus globulus*. *BioResources*, 10(1), 721–731.
- Gryc, V., Vavrcík, K., Horn, K. (2011). Density of juvenile and mature wood of selected coniferous species. *Journal of Forest Science*, 57, 123–130.
- Guilley, E. (2000). La densité du bois de chêne sessile (*Quercus petraea* Liebl.) Elaboration d'un modèle pour l'analyse des variabilités intraet interarbres. Origine et Evaluation nondestructive de l'effet "arbre", Interprétation anatomique du modèle proposé. Ecole Nationale du Genie Rural des Eaux et Forêts, Nancy, France.
- Gutierrez, A., del Rio, J.C., Gonza'lez-Vila, F. (1998). Variation in the composition of wood extractives from *Eucalyptus globulus* during seasoning. *Journal of Wood Chemistry and Technology*, 18(4), 439–446.
- Hamdani, F.A., Darusman, D., Tiryana, T. (2015). Evaluasi praktek tebang butuh di hutan rakyat Kabupaten Ciamis Provinsi Jawa Barat. *Risalah Kebijakan Pertanian dan Lingkungan*, 2, 33–41.
- Hamidah, S., Burhanudin, V., Istikowati, W. T. (2009). Kajian sifat-sifat dasar kayu manis sebagai pertimbangan pemanfaatan limbah pemanenan kulit kayu manis (*Cinnamomum burmanii* Blume). *Jurnal Hutan Tropis Borneo*, 10(26), 210–223.
- Han, J.S., Rowell, J.S. (1997). Chemical composition of fibres. – Rowell, R.M., Young, R.A., Rowell, J.K. (eds.). - Paper and composites from agro-based resources. New York, Lewis Publishers, 83–134.

- Hansen, J.R., Türk, G., Vogg, R., Heim, E.B. (1997). Conifer carbohydrates physiology: updating classical views. In *Trees -Contributions to Modern Tree Physiology*, Rennenberg, H., Eschrich, W., Ziegler, H. Eds. Backhuys Publishers Leiden: The Netherlands, pp. 97-108.
- Hardjanto. (2017). *Pengelolaan hutan rakyat*. IPB Press, Bogor.
- Hasan, M.A. (2017). *Budidaya mahoni (Swietenia macrophylla King.)*. Balai Pengelolaan Hutan Wilayah Lebak dan Tangerang. Tangerang: Dinas Lingkungan Hidup dan Kehutanan Provinsi Banten. 1-29.
- Haslett, A.N., Young, G.D. (1990). Plantation grown tropical timbers 1.Wood property and processing evaluation procedures to improve usage. *Journal of Tropical Forest Science*, 3, 131-139.
- Haslett, A.N., Young, G.D., Britton, R.A.J. (1991) Plantation grown tropical timbers 2. Properties, processing and uses. *Journal of Tropical Forest Science*, 3, 229-237.
- Hauch, S., Magel, E. (1998). Extractable activities and protein content of sucrose-phosphate synthase, sucrose synthase and neutral invertase in trunk tissues of *Robinia pseudoacacia* are related to cambial wood production and heartwood formation. *Planta*, 207, 266-274.
- Hazenberg, G., Yang, K.C. (1991). Sapwood/heartwood width relationships with tree age in balsam fir. *IAWA Bulletin n.s.*, 12(1), 95-99.
- He, T., Marco, J., Soares, R., Yin, Y., Wiedenhoef, A.C. (2020). Machine learning models with quantitative wood anatomy data can discriminate between *Swietenia macrophylla* and *Swietenia mahagoni*. *Forest*, 11, 1-13. DOI: doi.org/10.3390/f11010036.
- Hillinger, C., Holl, W., Ziegler, H. (1996). Lipids and lipolytic enzymes in the trunkwood of *Robinia pseudiacacia* during heartwood formation. *Trees* 10, 376-381
- Hillis, W.E. (1987). *Heartwood and tree exudates*. Springer-Verlag, Berlin.
- Hillis, W.E. (1971). Distribution, properties and formation of some wood extractives. *Wood Science and Technology*, 5, 272-289.
- Hillis, W.E. (1999). The formation of heartwood and its extractives. In: John T. Romeo (ed) *Phytochemicals in Human Health Protection, Nutrition, and Plant Defense*. Springer US, New York, pp 215-253.
- Hillis, W.E., Hasegawa, M. (1963). The formation of polyphenols in trees I. Administration of c glucose and subsequent distribution of radioactivity. *Phytochemistry*, 2(2), 195-199.
- Hilsenbeck, C. E., Johnson, A.F., Beckman, J. (2020). NatureServe Conservation Status Factors. https://explorer.natureserve.org/Taxon/ELEMENT_GLOBAL.2.150600/Swietenia_mahagoni. (Akses 8 April 2022).
- Hsu, E.W. (1997). Wood quality requirements for panel products. In: Zhang SY, Gosselin R, Chauret G (eds) *Timber Management Toward Wood Quality and End-Product Value*. Proceedings of the ATIA/IUFRO International Wood Quality Workshop, Quebec City, Part I, pp 7-10.

- IAWA committee. (1964). Multilingual glossary of terms used in wood anatomy. Committee on Nomenclature, International Association of Wood Anatomists (IAWA). Verlagsanstalt Buchdruckerei Konkordia Winterthur, Zürich.
- Igartúa, D.V., Monteoliva, S. (2009). Densidad básica de la madera de *Acacia melanoxylon* R. Br. en relación con la altura de muestreo, el árbol y el sitio. *Investigación Agraria: Sistemas y Recursos Forestales*, 18(1), 101–10.
- Iglesias, C., Santos, A.J.A., Martínez, J., Pereira, H., Anjos, O. (2017). Influence of heartwood on wood density and pulp properties explained by machine learning techniques. *Forest*, 8, 1-10. DOI: 10.3390/f8010020.
- Ikonen, V-P., Peltola, H., Wilhelmsson, L., Kilpeläinen, A., Väisänen, H., Nuutinen, T., Kellomäki, S. (2008). Modelling the distribution of wood properties along the stems of Scots pine (*Pinus sylvestris* L.) and Norway spruce (*Picea abies* (L.) Karst.) as affected by silvicultural management. *Forest Ecology and Management*, 256, 1356–1371. Doi:10.1016/j.foreco.2008.06.039.
- Imamura, H. (1989). Contribution of extractives to wood characteristics. In: Rowe J.W. (eds) *Natural Products of Woody Plants*. Springer Series in Wood Science. Springer, Berlin, Heidelberg. pp. 865-857.
- Iswanto, A.H., Siregar, Y.S., Susilowati, A., Darwis, A., Hartono, R., Wirjosentono, B., Rachmat, H.H., Hidayat, A., Fatriasari, W. (2019). Variation in chemical constituent of *Styrax sumatrana* wood growing at different cultivation site in North Sumatra, Indonesia. *Biodiversitas*, 20(2), 448–452.
- ITIS .(2021). *Swietenia mahagoni* (L.) Jacq. Taxonomic Serial No.: 29026. https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=29026#null (Akses 19 Februari 2021).
- Jahan, M.S., Mun, S.P. (2005). Effect of tree age on the cellulose structure of Nalita wood (*Trema orientalis*). *Wood Science and Technology*, 39, 367–373. DOI: 10.1007/s00226-005-0291-7.
- Jankowsky, I.P., Galvão, A.P.M. (1979). Influence of extractives content on the equilibrium moisture content of wood. (Influência do teor de extrativos na umidade de equilíbrio da madeira). *IPEF*, 18, 1–33.
- Johnson, L. (1942). Studies on the relation of growth rate to wood quality in *Populus* hybrids. *Canadian Journal of Research*, 20, 28-40. DOI: 10.1139/cjr42c-003.
- Jøker, D. (2000). *Swietenia mahagoni*. Seed Leaflet, (18). Danida Forest Seed Centre, University of Copenhagen, Denmark.
- Jorgensen, E., Balsille, D. (1969). Formation of heartwood phenols in callus tissue cultures of red pine. *Canadian Journal of Research*, 47, 1015-1016.
- Kampe, A., Magel, E. (2013). New insights into heartwood and heartwood formation. In *Cellular aspects of wood formation*. Berlin, Heidelberg, Springer, pp. 71-95.
- Kasuga, J., Arakawa, K., Fujikawa, S. (2007). High accumulation of soluble sugars in deep supercooling Japanese white birch xylem parenchyma cells. *New Phytologist*, 174, 569-579.

- Kiaei, M., Farsi, M. (2016). Vertical variation of density, flexural strength and stiffness of Persian silk wood. *Madera y Bosques*, 22(1), 169-175.
- Knapic S, Pereira H. (2005). Within-tree variation of heartwood and ring width in maritime pine (*Pinus pinaster* Ait.). *Forest Ecology and Management*, 210, 81–89.
- Knapic, S., Tavares, F., Pereira, H. (2006). Heartwood variation in *Acacia melanoxylon* R. Br. trees. *Forestry (Oxf.)*. 79,371– 380.
- Kord, B., Kialashaki, A., Kord, B. (2010). The within-tree variation in wood density and shrinkage, and their relationship in *Populus euramericana*. *Turkish Journal of Agriculture and Forestry*, 34, 121- 126. DOI: 10.3906/tar-0903-14.
- Kort, I. (1993). Relationships between sapwood amount, latewood percentage, moisture content and crown vitality of *Douglas-fir*, *Pseudotsuga menziesii*. *IAWA Journal*, 14, 413–427. DOI: 10.1163/22941932-90000596.
- Kozłowicz, K., Różyło, R., Gładyszewska, B., Matwijczuk, A., Gładyszewski, G., Chocyk, D., Samborska, K., Piekut, J., Smolewska, M. (2020). Identification of sugars and phenolic compounds in honey powders with the use of GC–MS, FTIR spectroscopy, and X-ray diffraction. *Scientific Report*, 10, 1-10. DOI: 10.1038/s41598-020-73306-7.
- Krisdianto. (2004). Perubahan struktur anatomi kayu akibat pemanasan dengan microwave. *Journal Ilmu dan Teknologi Kayu Tropis*, 2(2), 73-78.
- Krisnawati H, Kallio M, Kanninen M (2011) *Swietenia macrophylla* King: ecology, silviculture and productivity. CIFOR, Bogor, Indonesia.
- Kuroda, K., Yamashita, K., Fujiwara, T. (2009). Cellular level observation of water loss and the refilling of tracheids in the xylem of *Cryptomeria japonica* during heartwood formation. *Trees*, 23, 1163–1172. DOI: 10.1007/s00468-009-0356-6.
- Langum, C.E., Yadama, V., Lowell, E.C. (2009). Physical and mechanical properties of young growth Douglas-fir and western hemlock from western Washington. *Forest Product Journal*, 59, 37-47.
- Lachowicz, H., Wróblewska, H., Wojtan, R., Sajdak, M. (2019). The effect of tree age on the chemical composition of the wood of silver birch (*Betula pendula* Roth.) in Poland. *Wood Science and Technology*, 53, 1135–1155. DOI: 10.1007/s00226-019-01121-z.
- Larcher, W. (1995). *Physiological Plant Ecology*, 3rd ed., Springer, Berlin, Germany.
- Lebourgeois, F. (2000). Climatic signals in earlywood, latewood and total ring width of Corsican pine from western France. *Annals of Forest Science*, 57, 155–164.
- Lehnebach, R., Morel, H., Bossu, J., Moguedec, G.L., Amusant, N., Beauchene, J., Nicolini, E. (2016). Heartwood/sapwood profile and the tradeoff between trunk and crown increment in a natural forest: the case study of a tropical tree (*Dicorynia guianensis* Amsh., Fabaceae). *Trees*. DOI: 10.1007/s00468-016-1473-7.

- Lestari, E., Pramasari, D.A., Amin, Y., Adi, D.S., Bahanawan, A., Dwianto, W. 2016. The chemical components changes of platinum teak wood. – Proceedings of the 6th International Symposium for Sustainable Humanosphere, Indonesia, Nov. 2016. Bogor, Humanosphere Science School, 165–171.
- Lin, B.-D., Zhang, C.-R., Yang, S.-P., Wu, Y., Yue, J.-M. (2011). Phragmalin-type limonoid orthoesters from the twigs of *Swietenia macrophylla*. *Chemical and Pharmaceutical Bulletin*, 59(4), 458–465.
- Lindberg, B., Selleby, L. (1958). Birch wood constituents I. Carbohydrates of low molecular weight. *Acta Chemica Scandinavica*, 12, 1512–1515. DOI: 10.3891/acta.chem.scand.12-1512.
- Lindström, H. (1996). Basic density in Norway spruce. Part III. Development from pith outwards. *Wood Fiber Science*, 28, 391–405.
- Li, X., Wu, H.X., Southerton, S.G. (2011). Transcriptome profiling of wood maturation in *Pinus radiata* identifies differentially expressed genes with implications in juvenile and mature wood variation. *Gene*, 487, 62–71.
- Longuetaud, F., Mothe, F., Santenoise, P., Diop, N., Dlouha, J., Fournier, M., Deleuze, C. (2017). Patterns of within-stem variations in wood specific gravity and water content for five temperate tree species. *Annals of Forest Science*, 74, 1-19.
- Longwood, F.R. (1962) Present and potential commercial timber of the caribbean with special reference to the west indies, the guianas, and british honduras. Northeastern forest experiment station. United States Department of Agriculture, Washington D.C.
- Lourenço, A., Baptista, I., Gominho, J., Pereira, H. (2008). The influence of heartwood on the pulping properties of *Acacia melanoxylon* wood. *Journal of Wood Science*, 54, 464–469.
- Lundgren, C. (2004). Microfibril angle and density patterns of fertilized and irrigated Norway spruce. *Silva Fennica*, 38, 107–117.
- Lukmandaru, G. (2009). Chemical and colour properties of teak heartwood from three different ages. *Journal of Tropical Wood Science and Technology*, 7(1), 1–7.
- Lukmandaru, G. (2011). Variability in the natural termite resistance of plantation teak wood and its relations with wood extractive content and color properties. *Journal of Forestry Research*, 8(1), 17–31.
- Lukmandaru, G., Falaah, A.N., Listyanto, T., Rodiana, D. (2021). Extractive content and colour properties of 11-year-old superior teak wood. *Wood Research Journal*, 12(1), 10–17.
- Lukmandaru, G., Susanti, D., Widyorini, R. (2018). Sifat kimia kayu mahoni yang dimodifikasi dengan perlakuan panas. *Jurnal Penelitian Kehutanan Wallacea*, 7(1), 37-46.
- Magel, E., Abdel-Latif, A., Hampp, R. (2001). Non-structural carbohydrates and catalytic activities of sucrose metabolizing enzymes in trunks of two *Juglans* species and their role in heartwood formation. *Holzforschung*, 55(2), 135–145.

- Magel, E. A., Drouet, A., Claudot, A. C., Ziegler, H. (1991). Formation of heartwood substances in the stem of *Robinia pseudoacacia* L. *Trees*, 5, 203–207. DOI: 10.1007/bf00227526.
- Magel, E., Einig, W., Hampp, R. (2000). Carbohydrates in trees. *Developments in Crop Science*, 26, 317–336.
- Magel, E., Hillinger, C., Höll, W., Ziegler, H. (1997). Biochemistry and physiology of heartwood formation: Role of reserve substances. – Rennenberg, H., Eschrich, W., Ziegler, H. (eds.). *Trees - Contributions to Modern Tree Physiology*. Leiden, Backhuys Publishers, pp. 477– 506.
- Magel, E.A., Holl, W. (1993). Storage carbohydrates and adenine nucleotides in trunks of *Fagus sylvatica* L. in relation to discolored wood. *Holzforschung*, 47, 19–24. DOI: 10.1515/hfsg.1993.47.1.19.
- Magel, E., Jay-Allemand, C., Ziegler, H. (1994). Formation of heartwood substances in the stemwood of *Robinia pseudoacacia* L. II. Distribution of nonstructural carbohydrates and wood extractives across the trunk. *Trees*, 8, 165–171.
- Maiti, A., Dewanjee, S., Mandal, S.C., Annadurai, S. (2007). Exploration of antimicrobial potential of methanol and water extract of seeds of *Swietenia macrophylla* (family: Meliaceae), to substantiate folklore claim. *Iranian Journal of Pharmacology and Therapeutics*, 6, 99–102.
- Marsoem, S.N., Prasetyo, V.E., Sulisty, J., Sudaryono, S., Lukmandaru, G. (2014) Study of teak wood quality in Gunungkidul community forest III. Physical wood properties. *Jurnal Ilmu Kehutanan* 8, 75–88.
- Martawijaya, A., Kartasujana, I., Kadir, K., Prawira, S.A. (2005). Atlas kayu Indonesia Jilid 1. Departemen Kehutanan, Badan Penelitian dan Pengembangan Kehutanan. Bogor, Indonesia.
- Martinez-Sotres, C., Lopez-Albarran, P., Cruz-de-Leon, J., Garcia-Moreno, T., Rutiaiga-Quinones, J.G., Vazquez-Marrufo, G., Tamariz-Mascarua, J., Herrera-Bucio, R., (2012). Medicarpin, an antifungal compound identified in hexane extract of *Dalbergia congestiflora* Pittier heartwood. *International Biodeterioration & Biodegradation*, 69, 38–40.
- Masendra, Arisandi, R., Purba, B.A.V., Sumantri, F., Ihda, F.V., Wati, F.Z., Lukmandaru, G. (2020). Extractives contributing to the color of *Swietenia macrophylla*'s bark. *Wood Research Journal*, 11(1), 20–26.
- Mayer, I., Koch, G., and Puls, J. (2006). Topochemical investigations of wood extractives and their influence on colour changes in American black cherry (*Prunus serotina* Borkh.). *Holzforschung*, 60, 589–594. doi: 10.1515/HF.2006.100.
- Metsa-Kortelainen, S., Viitanen, H. (2009). Decay resistance of sapwood and heartwood of untreated and thermally modified Scots pine and Norway spruce compared with some other wood species. *Wood Material Science and Engineering*, 3, 105–114. DOI: 10.1080/17480270903326140.
- Meszaros, E., Jakab, E., Varhegyi, G. (2007). TG/MS, Py-GC/MS and THM-GC/MS study of the composition and thermal behavior of extractive components of *Robinia pseudoacacia*. *Journal of Analytical and Applied Pyrolysis*, 79, 61–70.

- Miranda, I., Gominho, J., Araujo, C., Pereira, H. (2014). Family effects in heartwood content of *Eucalyptus globulus* L. *European Journal of Forest Research*, 133, 81–87.
- Miranda, I., Gominho, J., Lourenço, A., Pereira, H. (2006). The influence of irrigation and fertilization on heartwood and sapwood contents in 18-years-old *Eucalyptus globulus* trees. *Canadian Journal of Forest Research*, 36, 2675–2683.
- Miranda, I., Pereira, H. (2002). The variation of chemical composition and pulping yield with age and growth factors in young *Eucalyptus globulus*. *Wood and Fiber Science*, 34, 140–145.
- Miranda, I., Sousa, V., Ferreira, J., Pereira, H. (2017). Chemical characterization and extractives composition of heartwood and sapwood from *Quercus faginea*. *PLOS ONE*, 12(6), e0179268. DOI: 10.1371/journal.pone.0179268.
- Mishra, G., Collings, D.A., Altaner, C.M. (2016). Physiological changes during heartwood formation in young *Eucalyptus bosistoana* trees. *IAWA Journal*, 39(4), 382–394.
- Miyamoto, N., Iizuka, K., Nasu, J., Yamada, H. (2016). Genetic effects on heartwood color variation in *Cryptomeria japonica*. *Silvae Genetica*, 65(2), 80–87.
- Moghadamtousi, S.Z., Goh, B.H., Chan, C.K., Shabab, T., Kadir, H.A. (2013). Review biological activities and phytochemicals of *Swietenia macrophylla* King. *Molecules*, 18, 10465–10483.
- Monties B (1991) Plant cell walls as fibrous lignocellulosic composites: relations with lignin structure and function. *Animal Feed Science and Technology*, 32, 159–175. DOI: 10.1016/0377-8401(91)90019-O
- Morais, M.C., Pereira, H. (2007). Heartwood and sapwood variation in *Eucalyptus globulus* Labill. Trees at the end of rotation for pulpwood production. *Annals of Forest Science*, 64, 665–671. DOI: 10.1051/forest:2007045.
- Morais, M.C., Pereira, H. (2012). Variation of extractives content in heartwood and sapwood of *Eucalyptus globulus* trees. *Wood Science and Technology*, 46, 709–719.
- Moya, R., Perez, D. (2008). Effect of physical and chemical soil properties on physical wood characteristics of *Tectona grandis* plantations in Costa Rica. *Journal of Tropical Forest Science*, 20, 147–155.
- Moya, R., Berrocal, A. (2010). Wood colour variation in sapwood and heartwood of young trees of *Tectona grandis* and its relationship with plantation characteristics, site, and decay resistance. *Annals and Forest Science*, 67, 109–120.
- Moya, R., Munoz, F. (2010). Physical and mechanical properties of eight fast-growing plantation species in Costa Rica. *Journal of Tropical Forest Science*, 22, 317–328.

- Moya, R., Bond, B., Quesada, H. (2014). A review of heartwood properties of *Tectona grandis* trees from fast-growth plantations. *Wood Science and Technology*, 48, 411-433. DOI: 10.1007/s00226-014-0618-3
- Moya, R., Salas, C., Berrocal, A., Valverde, J.C. (2015). Evaluation of chemical compositions, air-dry, preservation and workability of eight fastgrowing plantation species in Costa Rica. *Madera Bosques*, 21, 31-47.
- Mulholland, D.A., Parel, B., Coombes, P.H. (2000). The Chemistry of the Meliaceae and Ptaeroxylaceae of southern and eastern Africa and Madagascar. *Current Organic Chemistry*, 4, 1011-1054. DOI: 10.2174/1385272003375941.
- Murmanis, L. (1975). Formation of tylosis in felled *Querus rubra* L. *Wood Science and Technology*, 9, 3-14.
- Nahed, G.A.A., Hashish, K.I., Mahgoub, M.H., Mazhar, A.A.M. (2017). Ability of *Swietenia mahagoni* seedlings to remove of wastewater pollutants and bio-fertilization effect on growth and chemical constituents. *Agricultural Engineering International: CIGR Journal*, Special issue, 227–233.
- Nair, N.M.B. (1988). Wood anatomy and heartwood formation in Neem (*Azadirachta indica* A. Juss.). *Botanical Journal of the Linnean Society*, 97, 79-90.
- Nair, N.M.B., Shah, J.J. (1983). Histochemistry of paraquatred wood in *Azadirachta indica* A. Juss. *IAWA Bulletin n.s.*, 4(4), 249-254.
- Nakaba, S., Yamagishi, Y., Sano, Y., Funada, R. (2012). Temporally and spatially controlled death of parenchyma cells is involved in heartwood formation in pith regions of branches of *Robinia pseudoacacia* var. *inermis*. *Jornal of Wood Science*, 58, 69–76. DOI: 10.1007/s10086-011-1221-y.
- Nakada, R., Fukatsu, E. (2012). Seasonal variation of heartwood formation in *Larix kaempferi*. *Tree Physiology*, 32, 1497–1508. DOI: 10.1093/treephys/tps108.
- Nascimento, M.S., Santana, A.L.B.D., Maranhão, C.A., Oliveira, L.S., Bieber, L. 2013. Phenolic extractives and natural resistance of wood. *Intech.Open*. [WWW document]. – URL <http://dx.doi.org/10.5772/56358>. [Accessed 14 June 2021].
- Nelson, N.D. (1977). Xylem ethylene, phenol-oxidizing enzymes and nitrogen and heartwood formation in walnut and cherry. *Canadian Journal of Botany*, 56, 626-634.
- Neverova, N.A., Levchuk, A.A., Ostroukhova, L.A., Medvedeva, E.N., Onuchina, N.A., Babkin, V.A. (2013). Distribution of extractive substances in wood of the Siberian larch (*Larix sibirica* Ledeb.). *Russian Journal of Bioorganic Chemistry*, 39(7), 712–719.
- Niamke, F.B., Amusant, N., Kokutse, A.D., Chaix, G., Charpentier, J.P., Adima, A.A., Kati-koulibaly, S., C. Jay-allemant, C. (2010). Radial distribution of non-structural carbohydrates in Malaysian teak. *International Journal of Biological and Chemical Sciences*, 4, 710-720.

- Niamké, F.B., Amusant, N., Charpentier, J.P., Chaix, G., Baissac, Y., Boutahar, N., Adima, A.A., Kati-Coulibaly, S., Jay-Allemand, C. (2011). Relationships between biochemical attributes (non-structural carbohydrates and phenolics) and natural durability against fungi in dry teak wood (*Tectona grandis* L. f.). *Annals of Forest Science*, 68, 201– 211. DOI: 10.1007/s13595-011-0021-2.
- Nicholas, I.D., Young, G.D., Gifford, H.H. (1994). Wood properties of *Acacia melanoxylon*: variation within and between four seedlots. In: Faces of farm forestry, proceedings of the Australian forest growers conference, Launceston, Tasmania, Australia. Australian Forest Growers, Braddon, ACT, Australia, pp. 105–110.
- Nicholas, I.D., Dungey, H., Gifford, H.H., Cox, J., Hodgkiss, P., Jones, T. (2007). Preliminary analysis of the wood properties of *Acacia melanoxylon* thinnings from five North Island regime trials in New Zealand. In: Beadle C. L. and Brown A. G. (eds) *Acacia utilisation and management: adding value—3rd Blackwood Industry Group (BIG) workshop*, Marysville, VIC, 26–29 April 2006. RIRDC publication no. 07/095, Canberra, Australia. pp. 14–20.
- Nobuchi, T., Harada, H. (1983). Physiological features of the “white zonw” of Sugi (*Cryptomeria japonica* D. Don) cytological structure and moisture content. *Mokuzai Gakkaishi*, 29(12), 824-832.
- Nobuchi, T., Janmahasatian, S., Sakai, M. (1996). Seasonal changes of wood formation and some characteristics of heartwood formation in teak (*Tectona grandis* L.) plantation. *Kasegaretsart Journal: Natural Science*, 30(2), 254–263.
- Nobuchi, T., Okada, N., Nishida, M., Siripatanadilok, S., Veenin, T., Tobing, T.L., Sahri, M.H. (2005). Some characteristics of wood formation in Teak (*Tectona grandis*) with special reference to water conditions. In *Quality timber products of teak from sustainable forest management.*, Bhat, K.M., Nair, K.K.N., Bhat, K.V., Muralidharan, E.M., Sharma, J.K. Eds. Proceedings of international conference on quality timber products of teak from sustainable forest management. India, pp. 495–499.
- Nobuchi, T., Sato, T., Iwata, R., Harada, H. (1984). Season of heartwood formation and related cytological structure of ray parenchyma cells in *Robinia pseudoacacia*. *Mokuzai Gakkaishi*, 30(8), 628-636.
- Nobuchi, T., Takahara, S., Harada, H. (1979). Studies on the survival rate of ray parenchyma cells with aging process in coniferous secondary xylem. *Bulletin Kyoto University Forest*, 51, 239-246.
- Normand, D., Sallenave, P. (1955). Caracteristiques et proprietes des acajous (*Swietenia et Khaya*). *Bois et Forets des Tropiques*, 59, 43-52.
- Novitskaya, L.L., Kushnir, F.V. (2006). The Role of sucrose in regulation of trunk tissue development in *Betula pendula* Roth. *Journal of Plant Growth Regulation*, 25, 18-29.
- Ofori, J., Brentuo, B. (2003). Green moisture content, basic density, shrinkage and drying characteristics of the wood of *Cedrela odorata* grown in Ghana. *Journal of Tropical Forest Science*. 17(2), 211-223.

- Okkonen, E.A., Wahlgren, H.E., Maeglin, R.R. (1972). Relationships of specific gravity to tree height in commercially important species. *Forest Product Journal*, 22, 37-41.
- Oliva, A.G., Baonza-Merino, V., Fernández-Golfín, Seco, J.I., Conde-García, M., Hermoso- Prieto, E. (2006). Effect of growth conditions on wood density of Spanish *Pinus nigra*. *Wood Science and Technology*, 40, 190–204. Doi: 10.1007/s00226- 005-0014-0.
- Orwa C, Mutua A., Kindt, Jamnadass R, Anthony, S. (2009). Agroforestry Database:a tree reference and selection guide version 4.0 (<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>).
- Panshin, A.J. (1933) Comparative anatomy of the woods of the *Meliaceae*, Sub-Family *Swietenioideae*. *American Journal of Botany*, 20, 638-668. DOI: 10.2307/2436127.
- Peichl, M., Arain, M.A. (2006). Above and below ground ecosystem biomass and carbon pools in an age-sequence of temperate pine plantation forests. *Agricultural and Forest Meteorology*, 140, 51- 63. DOI: 10.1016/j.agrformet.2006.08.004.
- Pennington, T.D. (1981). *Meliaceae*. Flora Neotropica Monograph 28. [Swietenioideae by B.T. Styles and Chemotaxonomy by D.A.H. Taylor].
- Pereira, H. (1988). Variability in the chemical composition of plantation eucalypts (*Eucalyptus globulus* Labill.). *Wood and Fiber Science*, 20, 82–89.
- Perez, D., Kanninen, M. (2003). Heartwood, sapwood and bark content, and wood dry density of young and mature teak (*Tectona grandis*) trees grown in Costa Rica. *Silva Fennica*, 37, 45–54. DOI: 10.14214/sf.511.
- Piispanen, R., Saranpää, P. (2001). Variation of non-structural carbohydrates in silver birch (*Betula pendula* Roth) wood. *Trees*, 15, 444–451.
- Pillai, P.K.C., Pandalai, R.C., Dhamodaran, T.K., Sankaran, KV. (2013). Wood density and heartwood proportion in *Eucalyptus* trees from intensively-managed short rotation plantations in Kerala, India. *Journal of Tropical Forest Science*, 25, 220-227.
- Pinto, I., Pereira, H., Usenius, A. (2003). Analysis of log shape and internal knots in twenty maritime pine (*Pinus pinaster* Ait.) stems based on visual scanning and computer aided reconstruction. *Annals of Forest Science*, 60, 137–144.
- Pinto, I., Pereira, H., Usenius, A. (2004). Heartwood and sapwood development within maritime pine (*Pinus pinaster* Ait.) stems. *Trees*, 18, 284-294. DOI: 10.1007/s00468-003-0305-8.
- Piqueras, S., Füchtner, S., Oliveira, RR., Gómez-Sánchez, A., Jelavic, S., Keplinger, T, Juan, A., Thygesen, LG. (2020). Understanding the formation of heartwood in Larch using synchrotron infrared imaging combined with multivariate analysis and atomic force microscope infrared spectroscopy. *Frontiers in Plant Science*, 10. DOI: 10.3389/fpls.2019.01701.
- Pratama, A.R., Yuwono, S.B., Hilmanto, R. (2015). Pengelolaan hutan rakyat oleh kelompok pemilik hutan rakyat di desa bandar dalam Kecamatan Sidomulyo Kabupaten Lampung Selatan. *Jurnal Sylva Lestari* 3, 99-112.

- Pratt, R.B., Jacobsen, A.L., Ewers, F.W., Davis, S.D. (2007). Relationships among xylem transport, biomechanics and storage in stems and roots of nine Rhamnaceae species of the California chaparral. *New Phytologist*, 174, 787–798.
- Prawirohatmodjo, S. (2004). Sifat-sifat Kimia Kayu. Yogyakarta: Yayasan Pembina Fakultas Kehutanan. Universitas Gadjah Mada. (tidak dipublikasikan).
- Prayitno, T.A. (1992). Sifat kimia kayu salam (*Zyzigium poliantha* Wight). Buletin, Fakultas Kehutanan, Universitas Gadjah Mada, 22, 53–65.
- Rababah, T., Ereifej, K., Esh, R., Al-u'datt, M., Alrababah, M., Yang, W. (2011). Antioxidant activities, total phenolics and HPLC analyses of the phenolic compounds of extracts from common Mediterranean plants. *Natural Product Research*, 25, 596–605. DOI: 10.1080/14786419.2010.488232.
- Randhir R, Lin YT, Shetty K (2004) Stimulation of phenolics, antioxidant and antimicrobial activities in dark germinated mung bean sprouts in response to peptide and phytochemical elicitors. *Process Biochemistry*, 39 (5), 637–646.
- Rastogi, R.P., Mehrotra, B.N. (1993). Compendium of Indian Medicinal Plants, Vol. 3. New Delhi, Publications and Information Directorate, CSIR. pp. 615-616.
- Ratnasingam, J., Ramasamy, G., Loras, F. (2016). The influence of log felling season on the extent of discoloration in Rubberwood sawn timber during the kiln drying process. *BioResources*, 11, 5003-5012.
- Raymond, C.A., Joe, B. (2007). Patterns of basic density variation for *Pinus radiata* grown in south-west slopes region of NSW, Australia. *New Zealand Journal of Forestry Science*, 37, 81–95.
- Rencoret, J., Gutiérrez, A., Nieto, L., Jiménez-Barbero, J., Faulds, C.B., Kim, H., Ralph, J., Martínez, Á.T., del Río, J.C. (2011). Lignin composition and structure in young versus adult *Eucalyptus globulus* plants. *Plant Physiology*, 155(2), 667–682.
- Rink, G., Phelps, J.E. (1989). Variation in heartwood and sapwood properties among 10-year-old black walnut trees. *Wood Science and Technology*, 21, 177–182.
- Rust, S. (1999). Comparison of three methods for determining the conductive xylem area of Scots pine (*Pinus sylvestris*). *Forestry*, 72, 103-108.
- Rutiaga Quiñones, J.G., Windeisen, E., Strobel, C. (1998a). Extractive content of *Swietenia macrophylla* King. (Extraktstoffgehalt von *Swietenia macrophylla* King). *European Journal of Wood and Wood Product*, 56(4), 228.
- Rutiaga Quiñones, J.G., Windeisen, E., Strobel, C. (1998b). Polysaccharide von *Swietenia macrophylla* King. *European Journal of Wood and Wood Product*, 56(4), 234.
- Saka, S. (2001). Chemical composition and distribution. In *Wood and Cellulosic Chemistry*; Hon, D.N.S., and Shiraishi, N., Eds. Marcel Dekker: New York, pp. 51-82.

- Santana, W.M.S., Calegario, N., Arantes, M.D.C., Trugilho, P.F. (2012). Effect of age and diameter class on the properties of wood from clonal *Eucalyptus. Cerne*, 18(1), 1–8.
- Saranpaa, P., Holl W. (1989). Soluble carbohydrates of *Pinus sylvestris* L. sapwood and heartwood. *Trees*, 3, 138-143. DOI: 10.1007/BF00226648.
- Sellin, A. (1994). Sapwood–heartwood proportion related to tree diameter, age, and growth rate in *Picea abies*. *Canadian Journal of Forest Research*, 24, 1022–1028.
- Sellin, A. (1996). Sapwood amount in *Picea abies* (L.) Karst. Determined by tree age and radial growth rate. *Holzforschung*, 50, 291–296.
- Sarhaan, A.M.Z., Arafa, A.M.S., Habba, E.E., El- Aziz, N.G.A., Mazhar, A.A., Youssef, N.M. (2015). Effect of growing media and microbien on growth and chemical composition of *Swietenia mahagoni* (L.) Jacq. *Plants. Journal of Horticultural Science & Ornamental Plants*, 7, 141-146.
- Savidge, R.A. (2003). Tree growth and wood quality. In: Barnett JR, Jeronimidis G (eds) Wood quality and its biological basis. Blackwell, Oxford, pp. 1–29.
- Skaar, C. (1988). Wood-water relations. Springer-Verlag, Berlin, Germany.
- Setyawan, C., Susanto, S., Lee, C.Y. (2019) Spatial modelling of watershed health assessment by using GIS. *IOP Conf. Series: Earth and Environmental Science* 355. DOI: 10.1088/1755-1315/355/1/012018.
- Shain, L., Hillis, W.E. (1973). Ethylene production in xylem of *Pinus radiata* in relation to heartwood formation. *Canadian Journal of Botany*, 51, 1331-1335
- Shain, L., Mackay, J.P.G. (1973) Seasonal fluctuation in respiration if aging xylem in relation to heartwood formation in *Pinus radiata*. *Canadian Journal of Botany*, 51, 737-741.
- Shigo, A.L., Hillis, W.E. (1973). Heartwood, discolored wood, and microorganisms in living trees. *Annual Review of Phytopathology*, 11, 197-222.
- Shmulsky, R., Jones, P.D. (2011). Forest Products and Wood Science: An Introduction, Sixth Edition. John Wiley & Sons, Inc. DOI: 10.1002/9780470960035.
- Singleton, R., Debell, D.S., Gartner, B.L. (2003). Effect of extraction on wood density of western hemlock (*Tsuga heterophylla* (Raf.) Sarg.). *Wood and Fiber Science*, 35, 363-369.
- Singleton, V.L., Orthofer, R., Lamuela-Raventós, R.M. (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymology*, 299, 152–178.
- Simpson, W., Tenwolde, A. (2013). Physical properties and moisture relations of wood. International Plant Protection Convention Physical properties and moisture relations of wood. TPFQ_2013_Jun_36, pp. 1-25.
- Sjökqvist T, Blom A, Wålinder MEP (2019) The influence of heartwood, sapwood and density on moisture fluctuations and crack formations of coated Norway spruce in outdoor exposure. *Journal of Wood Science*, 65, 1-9.

- Sjostrom E (1993) Kimia Kayu: Dasar-dasar dan Penggunaan. Edisi 2. Sastrohamidjojo H, penerjemah. Jogjakarta : Gadjah Mada University Press. Terjemahan dari: Wood Chemistry: Fundamentals and Applications.
- Solorzano, S., Moya, R., Chauhan, S. (2012a). Early genetic evaluation of morphology and some wood properties of *Tectona grandis* L clones. *Silva Genetica*, 61, 58–65.
- Solorzano S, Moya R, Murillo O. (2012b). Early prediction of basic density, shrinking, presence of growth stress, and dynamic elastic modulus based on the morphological tree parameters of *Tectona grandis*. *Journal of Wood Science*, 58, 290–299.
- Sorz, J., Hietz, P. (2008). Is oxygen involved in beech (*Fagus sylvatica*) red heartwood formation? *Trees*, 22, 175–185.
- Sotelo Montes, C., Hernández, R., Beaulieu, J., Weber, J. (2008). Genetic variation in wood color and its correlations with tree growth and wood density of *Calycophyllum spruceanum* at an early age in the Peruvian Amazon. *New Forest*, 35, 57–73.
- Stokes, A., Berthier, S. (2000). Irregular heartwood formation in *Pinus pinaster* Ait. is related to eccentric, radial, stem growth. *Forest Ecology and Management*, 135, 115–121.
- Stolarski, M.J., Szczukowski, S., Tworowski, J., Wróblewska, H., Krzyżaniak, M. (2011). Short rotation willow coppice biomass as an industrial and energy feedstock. *Industrial Crops and Products*, 33, 217–223.
- Streit, W., Fengel, D. (1994). On the changes of the extractive composition during heartwood formation in Quebracho Colorado (*Schinopsis balansae* Engl.). *Holzforschung*, 48, 15-20. DOI: 10.1515/hfsg.1994.48.s1.15.
- Sukadaryati, Yuniawati, Dulsalam. (2018). Timber harvesting in community forest (case study in Ciamis, West Java). *Jurnal Ilmu Kehutanan*, 12, 142-155.
- Sulaiman, A. (1994). Plant resources of South-East Asia. No. 5(1). In: Soerianegara, I & Lemmens, RHMJ (eds). Timber trees: major commercial timbers. Prosea, Bogor. pp 442-447.
- Szczepkowski, A., Nicewicz, D., Koczoń, P. (2007). The relationship between tree health and chemical composition of beech (*Fagus sylvatica* L.) and oak (*Quercus robur* L.) wood of Polish provenances. *Acta Scientiarum Polonorum Silvarum Colendarum Ratio et Industria Lignaria*, 6(3), 77–88.
- Taylor, A.M., Gartner, B.L., Morrell, J.J. (2002). Heartwood formation and natural durability-a review. *Wood and Fiber Science*, 34, 587-611.
- Thulasidas, P.K., Bhat, K.M., Okuma, T., (2006). Heartwood colour variation in home garden teak (*Tectona grandis*) from wet and dry localities of Kerala, India. *Journal of Tropical Forest Science*, 18, 51–54.
- Tomczak, A., Tomczak, K., Smarul, N., Rutkowski, K., Wenda, M., Jelonek, T. (2018). The gradient of wood moisture within-stem of sessile oak (*Quercus petaea* (Matt.) Liebl.) in summer. *Wood Research*, 63, 809-820.
- Topaloglu, E., Erisir, E. (2018). Longitudinal variation in selected wood properties of oriental beech and caucasian fir. *Maderas: Ciencia y Tecnología*, 20, 403 – 416.

- Trockenbrodt, M., Josue, J. (1999). Wood properties and utilization potential of plantation teak (*Tectona grandis*) Malaysia—a critical review. *Journal of Tropical Forest Science*, 5, 58–70.
- Tsoumis, G. (1991). Science and Technology of Wood: Structure, Properties, utilization. Von Nostrand Reinhold. New York.
- Tsuruta, K., Yoshida, Y., Kusumoto, N., Sekine, N., Ashitani, T., Takahashi, K. (2011). Inhibition activity of essential oils obtained from Japanese trees against *Skeletonema costatum*. *Journal of Wood Science*, 57, 520–525.
- Uetimane, Jr.E., Ali, A.C. (2011). Relationship between mechanical properties and selected anatomical features of ntholo (*Pseudolachnostylis maprounaefolia*). *Journal of Tropical Forest Science*, 23(2), 166–176.
- Uetimane, jr.E., Terziev, N., Dananiel, G. (2009). Wood anatomy of three lesser known species from Mozambique. *IAWA Journal*, 30, 277–291.
- Umezawa, T. 2001. Chemistry of extractives. – Hon, D.N.-S., Shiraishi, N. (eds.). Wood and Cellulosic Chemistry, 2nd Edition. New York, Marcel Dekker, pp. 213–242.
- Uprichard, J.M., Lloyd, J.A. (1980). Influence of tree age on the chemical composition of Radiata pine. *New Zealand Journal of Forestry Science*, 10, 551–557.
- Usda Forest Product Laboratory. (1999). Wood handbook-Wood as an engineering material. Tech. Rep. No. FPL GTR-I 13. USDA Forest Serv. Forest Product Lab. Madison WI. 466 pp.
- van Putten, R.-J., Winkelman, J.G.M., Keihan, F., van der Waal, J.C., de Jong, E., Heeres, H.J. (2014). Experimental and modelling studies on the solubility of D-arabinose, D-fructose, D-glucose, D-mannose, sucrose and D-xylose in methanol and methanol-water mixtures. *Industrial and Engineering Chemistry Research*, 53(19), 8285–8290.
- Velderrain-Rodríguez, G.R., Palafox-Carlos, H., Wall-Medrano, A., AyalaZavala, J.F., Chen, C.-Y.O., Robles-Sanchez, M., Astiazaran-García, H., Alvarez-Parrilla, E., González-Aguilar, G.A. (2014) Phenolic compounds: Their journey after intake. *Food & Function journal*, 5, 189–197.
- Wadsworth, F.H., Gonzalez, E. (2008) Sustained mahogany (*Swietenia macrophylla*) plantation heartwood increment. *Forest Ecology and Management*, 255, 320–323. DOI: 10.1016/j.foreco.2007.09.053.
- Wahyudi, I., Priadi, T., Rahayu, I.S. (2014). Characteristics and basic properties of 4 and 5 year-old of superior Teakwoods from West Java. *Jurnal Ilmu Pertanian Indonesia* 19, 50–56.
- Wanneng, P.X., Ozarska, B., Daian, M.S. (2014). Physical properties of *Tectona grandis* grown in Laos. *Journal of Tropical Forest Science*, 26, 389–396.
- Westoby, M., Wright, I.J. (2006). Land-plant ecology on the basis of functional traits. *Trends in Ecology & Evolution*, 21, 261–268.
- Wiemann, M.C., Williamson, GB. (1989). Radial gradients in the specific gravity of wood in some tropical and temperate trees. *Forest Science*, 35, 197–210. DOI: 10.1093/forestscience/35.1.197.

- Wilkes, J. (1984) The influence of rate of growth on the density and heartwood extractives content of eucalypt species. *Wood Science and Technology*, 18, 113–120.
- Wilkes, J. (1991). Heartwood development and its relationship to growth in *Pinus radiata*. *Wood Science and Technology*, 25, 85-90. DOI: 10.1007/BF00226808.
- Wilkins, A.P. (1991). Sapwood, heartwood and bark thickness of silviculturally treated *Eucalyptus grandis*. *Wood Science and Technology*, 25, 415–423. DOI: 10.1007/BF00225234.
- Windeisen, E., Klassen, A., Wegener, G. (2003). On the chemical characterization of plantation teakwood from Panama. *Holz als Roh und Werkstoff*, 61(6), 416-418.
- Woodcock, D., Shier, A.D. (2002). Wood specific gravity and its radial variations: the many ways to make a tree. *Trees*, 16, 437-443.
- Woeste, K. E. (2002). Heartwood production in a 35-year-old black walnut progeny test. *Canadian Journal of Forest Research*, 32, 177- 181.
- Wu, H.X., Eldridge, K.G., Matheson, A.C., Powell, M.B., McRae, T.A., Butcher, T.B., Johnson, I.G. (2007). Achievements in forest tree improvement in Australia and New Zealand 8. Successful introduction and breeding of radiata pine in Australia. *Australian Forestry*, 70, 215–225.
- Yamamoto, K. (1982). Yearly and seasonal process of maturation of ray parenchyma cells in *Pinus species*. *Research Bulletins of the College Experiment Forests - Hokkaido University*, 39, 245-296.
- Yamamoto, K., Sulaiman, O., Kitingan, C., Choon, L.W., Nhan, N.T. (2013). Moisture distribution in stems of *Acacia mangium*, *A. auriculiformis* and *Hybrid Acacia Trees*. *JARQ*, 37(3), 207 – 212.
- Yang, B., Jia, H., Zhao, Z., Pang, S., Cai, D. (2020): Horizontal and vertical distributions of heartwood for teak plantation. *Forests*, 11(225), 1-12.
- Yanchuk, A.D., Dancik, B.P., Micko, M.M. (1984). Variation and heritability of wood density and fiber length of trembling aspen in Alberta, Canada. *Silvae Genetica*, 33, 11-16.
- Yang, K.C., Hazenberg, G. (1991). Relationship between tree age and sapwood/heartwood width in *Populus tremuloides*. *Wood and Fiber Science*, 23, 247-252.
- Yang, K.C., Murchison, H.G. (1992) Sapwood thickness in *Pinus contorta* var. *latifolia*. *Canadian Journal of Forest Research*, 22, 2004–2006.
- Yao, J. (1970). Influence of growth rate on specific gravity and other selected properties of loblolly pine. *Wood Science and Technology*, 4, 163–175. Doi:10.1007/BF00571851.
- Yazaki, Y. (2015). Wood colors and their coloring matters: a review. *Natural Product Communications*, 10(3), 505–512.
- Yeh, T.-F., Braun, J.L., Goldfarb, B., Chang, H.-M., Kadla, J.F. (2006). Morphological and chemical variations between juvenile wood, mature wood, and compression wood of loblolly pine (*Pinus taeda* L.). *Holzforschung*, 60(1), 1–8.

- Yuniarti, T. (2008). *Ensiklopedia Tanaman Obat Tradisional*. Cetakan Pertama. Media Pressindo, Yogyakarta.
- Zhang, L.Y., Deng, X.W., Lei, X.D., Xiang, W.H., Peng, C.H., Lei, P.F., Yan, W.D. (2012). Determining stem biomass of *Pinus massoniana* L. Through variations in basic density. *Forestry*, 85, 601–609. Doi:10.1093/forestry/cps069.
- Zimmerman, M.H. (1983). Failure and "senescence" of xylem function. Pages 98-106 in M. H. Zimmermann, ed. *Xylem structure and the ascent of sap*. Springer-Verlag, Berlin, Germany. 143 pp.
- Zobel, B.J., Van Buijtenen, J.P. (1989). *Wood Variation, Its Causes and Control*. Springer-Verlag, Berlin.
- Zule, J., Može, A. (2003). GC analysis of extractive compounds in beech wood. *Journal of Separation Science*, 26, 1292-1294. DOI: 10.1002/jssc.200301416.
- Zulkahfi., Irawati, D., Listyanto, T., Rodiana, D., Lukmandaru, G. (2020). Extractives content and colour properties of 11-year-old Perhutani superior teakwood from Ngawi Forest Management Unit. *Jurnal Ilmu Kehutanan*, 14(2), 213–227.