



DAFTAR PUSTAKA

- Aguiar, L., T., Martins, J., C., Nadeau, S., Britto, R., R., Teixeira-Salmela, L., F., Faria C, D. 2017. Efficacy of interventions to improve physical activity levels in individuals with stroke: a systematic review protocol. *BMJ Open*. 2017 Jan 5;7(1):e012479.
- Ahmadizad, S., Avansar, A., S., Ebrahim, K., Avandi, M., Ghasemikaram, M. 2015. The effects of short-term high-intensity interval training vs. moderate-intensity continuous training on plasma levels of nesfatin-1 and inflammatory markers. *Horm Mol Biol Clin Invest*. 2015; 21(3): 165–173
- Aiken, M. 2019. An Updated Evaluation of Google Translate Accuracy. *Studies in Linguistics and Literature* ISSN 2573-6434 (Print) ISSN 2573-6426 (Online) Vol. 3, No. 3, 2019. University of Mississippi, Oxford, MS, USA
- Amanat, S., Sinaei, E., Panji, M., MohammadporHodki, R., Bagheri-Hosseinabadi, Z., Asadimehr H., et al. 2020. A Randomized Controlled Trial on the Effects of 12 Weeks of Aerobic, Resistance, and Combined Exercises Training on the Serum Levels of Nesfatin-1, Irisin-1 and HOMA-IR. *Front. Physiol*. 11:562895
- American College of Sports Medicine. (2022). ACSM's guidelines for exercise testing and prescription Eleventh Edition. Philadelphia :Lippincott Williams & Wilkins
- Ammann, B., C., Knols, R., H., Baschung, P., de Bie, R., A., de Bruin, E., D. 2014. Application of principles of exercise training in sub-acute and chronic stroke survivors: a systematic review. *BMC Neurol*. Aug 22;14:167.
- Arikan, Sukran. (2020). Effects of Acute and Chronic Exercises on Plasma Nesfatin-1 Levels in Young Adults. *Cyprus J Med Sci*. 2020; 5(1): 77-80.
- Asad, M., R., Ghadiri, M., Barzegari, A.2019. Effect of sequence of combined strength and endurance training on the levels of nesfatin-1 and some metabolic risk factors of overweight womenISSN: 2538-5887; Pathobiology Research. 2019;22(3):129-134
- Ayada, C., Toru, Ü., Korkut, Y. 2015.Nesfatin-1 and its effects on different systems. *Hippokratia*. Jan-Mar;19(1):4-10. PMID: 26435639; PMCID: PMC4574585.
- Aydin, S. 2013. Multi-functional peptide hormone NUCB2/nesfatin-1. *Endocrine*. Oct;44(2):312-25



- Bajer, B., Vlcek, M., Galusova, A., Imrich, R., Penesova, A. 2015. Exercise associated hormonal signals as powerful determinants of an effective fat mass loss. *Endocr Regul.* Jul;49(3):151-63.
- Baravati, S, A, P., Mogharnasi, M., Badiakhar, H., Rahimi. E. 2016. Response And Compatibility Of Plasma Levels Of Nesfatin-1, Glucose And Insulin Resistance Index To Circuit Resistance Training In Obese Disabled Men. Special ISSUE (ARBS).ISSN: 0976-3104
- Bilski, J., Mazur-Bialy, A. I., Surmiak, M., Hubalewska-Mazgaj, M., Pokorski, J., Nitecki, J., et al. 2020. Effect of Acute Sprint Exercise on Myokines and Food Intake Hormones in Young Healthy Men. *Int. J. Mol. Sci.* 2020, 21, 8848.
- Ciolac, Emmanuel, Guimaraes, Guilherme. 2004. Physical Exercise and Metabolic Syndrome. *Revista Brasileira de Medicina do Esporte.* July 2004.10. 325-330.
- Cramond. F, O'Mara-Eves. A, Doran-Constant. L, Rice. A. S, Macleod. M, Thomas. J. 2019. The development and evaluation of an online application to assist in the extraction of data from graphs for use in systematic reviews. *Wellcome Open Res.* 2019 Mar 7;3:157.
- Dissemination, University. (2001). Undertaking systematic reviews of research on effectiveness: CRD's guidance for carrying out or commissioning reviews. CRD Report Number 4 (2nd Edition). MARCH 2001. NHS Center for Reviews and Dissemination, University of York.
- Fedewa, M,V., Hathaway, E,D., Ward-Ritacco, C, L., Williams, T, D., Dobbs, W,C. 2018. The Effect of Chronic Exercise Training on Leptin: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Sports Med.* 2018 Jun;48(6):1437-1450.
- Feijóo-Bandín, S., Rodríguez-Penas, D., García-Rúa, V., Mosquera-Leal, A., González-Juanatey, J, R., Lago, F. 2016. Nesfatin-1: a new energy-regulating peptide with pleiotropic functions. Implications at cardiovascular level. *Endocrine.* Apr;52(1):11-29.
- Frigo, L, M., Chowen, J, A.2015. Hypothalamic Leptin and Ghrelin Signaling as Targets for Improvement in Metabolic Control. *Curr Pharm Des.* 2015;21(25):3596-605.
- García-Galiano, D., Navarro, V, M., Gaytan, F., Tena-Sempere, M. 2010. Expanding roles of NUCB2/nestin-1 in neuroendocrine regulation. *J Mol Endocrinol.* Nov;45(5):281-90.



- Haff, G. G., Triplett, N. T. 2016. Essentials of strength training and conditioning / National Strength and Conditioning Association Fourth edition. United States. Human Kinetics, 2016.
- Higgins, J. P. T., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M. J., Welch, V. A. 2022. *Cochrane Handbook for Systematic Review of Interventions version 6.3* (diperbarui Februari 2022). Cochrane, 2022. Available from www.training.cochrane.org/handbook.
- Jabar, B., Farhad,G., Adel, R., Vahid, T. 2012. Effect Of Single Bout Of Aerobic Exercise On Serum Nesfatin-1 Levels In Non-Athlete Elderly Men. *Medical Journal Of Tabriz University Of Medical Sciences*. Okt. – Nov. 2012 , Vol.34 , Number 4; Page(s) 25 To 30
- Jafari, M., Mogharnas, M. 2015. The Protective Effect of Different Methods of Exercise Training on Plasma Levels of Nesfatin-1, Cardiorespiratory Endurance and Body Composition in Overweight and Obese Females. *Mod Care J* 2015 April;12(2):61-67
- Mogharnasi, M., Chadorneshin, H, T., Baravati,S, A, P., Teymuri, A. 2019 (1). Effects Of Upper-Body Resistance Exercise Training On Serum Nesfatin-1 Level, Insulin Resistance, And Body Composition In Obese Paraplegic Men, *Disability and Health Journal*, Volume 12, Issue 1,2019, Pages 29-34.
- Mogharnasi, M., Chadorneshin, H, T., Abbasi-Deloei, N. 2019 (2). Effect of exercise training type on plasma levels of vaspin, nesfatin-1, and high-sensitivity C-reactive protein in overweight and obese women, *Obesity Medicine*, Volume 13,2019, Pages 34-38,
- Mogharnasi,M., TajiTabas, A., Tashakorizadeh, M., Nayebifar, S, H. 2018. The Effects of Resistance and Endurance Training on Levels of Nesfatin-1, HSP70, Insulin Resistance and Body Composition in Women with Type 2 Diabetes Mellitus, *Science & Sports* ,Volume 34, Issue 1,2019 ,Pages e15-e23,
- Murray, B., Kenney, W. L. 2021. Practical Guide to Exercise Physiology: The Science of Exercise Training and Performance Nutrition Second Edition. United States of America. Human Kinetics.
- Nazar Ali, P., Fathi, R., Imeri, B, B.S. 2014. The effect of 8 weeks resistance training on plasma Nesfatin-1 levels in overweight women. (Autumn 2013 and Winter 2014), Vol. 3, No. 2
- Oh-I, S., Shimizu, H., Satoh, T., Okada, S., Adachi, S., Inoue, K., et al. 2006. Identification of nesfatin-1 as a satiety molecule in the hypothalamus. *Nature*. 2006 Oct 12;443(7112):709-12.



- Özdenk, S., Kurudirek, I. 2021. Effects of Moderate Exercise Intensity on Serum Nesfatin-1 Levels in Young Females. *Kastamonu Med J.* 2021; 1(3): 67-70
- Öztürk Özkan G. 2020. Effects of Nesfatin-1 on Food Intake and Hyperglycemia. *J Am Coll Nutr.* 2020 May-Jun;39(4):345-351
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., et al. (2021) The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *PLoS Med* 18(3): e1003583.
- Plowman, S. H., Smith, D. L. 2017. Exercise Physiology For Health, Fitness, and Performance Fifth Edition. Wolters Kluwer. Philadelphia.
- Prinz, P., Stengel, A. 2016. Nesfatin-1: current status as a peripheral hormone and future prospects. *Curr Opin Pharmacol.* Dec;31:19-24..
- Rahimi, G.R., Hejazi, K., Hofmeister, M. (2022). The Effect Of Exercise Interventions On Irisin Level: A Systematic Review And Meta-Analysis Of Randomized Controlled Trials. *EXCLI Journal.* 2022;21:524-539
- Review Manager (RevMan) [Computer program]. Version 5.4. 2020. The Cochrane
- Riebe, D., Ehrman, J. K., Liguori, G., Magal, M. 2018. American College of Sports Medicine's guidelines for exercise testing and prescription Tenth edition. Wolters Kluwer. Philadelphia
- Feijóo-Bandín, S., Rodríguez-Penas, D., García-Rúa, V., Mosquera-Leal, A., Otero, M. F., Pereira, E., et al. 2013. Nesfatin-1 in human and murine cardiomyocytes: synthesis, secretion, and mobilization of GLUT-4. *Endocrinology.* Dec;154(12):4757-67.
- Riva, M., Nitert, M. D., Voss, U., Sathanoori, R., Lindqvist, A., Ling, C., et al. 2011. Nesfatin-1 stimulates glucagon and insulin secretion and beta cell NUCB2 is reduced in human type 2 diabetic subjects. *Cell Tissue Res.* Dec;346(3):393-405.
- Schalla, M. A., Stengel, A. 2018. Current Understanding of the Role of Nesfatin-1. *J Endocr Soc.* Sep 10;2(10):1188-1206
- Shea, B. J., Reeves, B. C., Wells, G., Thuku, M., Hamel, C., Moran, J., et al. 2017. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ.* 2017 Sep 21;358:j4008.
- Shimizu, H., Ohsaki, A., Oh-I, S., Okada, S., Mori, M. 2009. A new anorexigenic protein, nesfatin-1. *Peptides.* May;30(5):995-8.
- Soori, R., Mahmoodi, F., Ranjbar, K., Ramezankhani, A. 2015. Effects of regular physical activity on levels of nesfatin-1, neuropeptideY and cortisol in obese men. *Winter 2017, 19 (1):64-74 Plasma Levels Of Nesfatin-1,*



Glucose And Insulin Resistance Index To Circuit Resistance Training
In Obese Disabled Men. Special Issue (ARBS) papoli. ISSN: 0976-3104

- Stengel, A. 2015. Nesfatin-1 - More than a food intake regulatory peptide. *Peptides*. Oct;72:175-83.
- Stengel, A., Mori, M., Taché, Y. 2013. The Role of Nesfatin-1 in The Regulation of Food Intake and Body Weight – Recent Developments and Future Endeavors. *Obes Rev.* November ; 14(11)
- Sterne, J. A., Hernán, M. A., Reeves, B. C., Savović, J., Berkman, N. D., Viswanathan, M., et al. 2016. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*. 2016 Oct 12;355:i4919.
- Szlachcic, A., Sliwowski, Z., Krzysiek-Maczka, G., Majka, J., Surmiak, M., Pajdo, R., et al. 2013. New satiety hormone nesfatin-1 protects gastric mucosa against stress-induced injury: mechanistic roles of prostaglandins, nitric oxide, sensory nerves and vanilloid receptors. *Peptides* 49 (2013) 9-20.
- Tahmasebi, W., Alizadeh, R., Nourshahi, M., Salehpour, M. 2019. Plasma Nesfatin-1 and Insulin Resistance Changes Immediately After Circuit-Resistance Exercise and During Recovery Periods in Healthy Men. *J Kermanshah UnivMed Sci.* In Press(In Press):e96932.
- Taji Tabas, A., Mogharnasi, M. 2015. The Effect Of 10 Week Resistance Exercise Training On Serum Levels Of Nesfatin-1 And Insulin Resistance Index In Women With Type 2 Diabetes. *Iranian journal of Diabetes and Metabolism*; Vol.14, No3, 2015
- Tavassoli H, Tofighi A, Hossein panah F, Hedaytai M. 2014. Appetite and Exercise Influence of 12 Weeks of Circuit Resistance Training on the Nesfatin-1 to Acylated Ghrelin Ratio of Plasma in Overweight Adolescents. *Iranian Journal of Endocrinology and Metabolism*. 2014; 15 (6) :519-526
- Tekin, T.,Cicek, B., Konyaligil, N. 2019. Regulatory Peptide Nesfatin-1 and its Relationship with Metabolic Syndrome. *Eurasian J Med*.2019;51(3): 280-4
- Tofighi, A., Mehrabani, J., Khadivi, S, M.2014. The Effect Of 8 Week Aerobic Exercise On Nesfatin-1 And Acylated Ghrelin In Young Obese Men. *Medical Journal of Mashhad University of Medical Sciences*. Vol. 57, No. 3 P: 562-570. Jul-Aug 2014
- WHO, 2021. Obesity and Overweight. Unduh tanggal 20 Juni 2022 di <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
- Xu, Y., Zai, Z., Zhang, T., Wang, L., Qian, X., Xu, D., et al. 2022. Nesfatin-1 exerts protective effects on acidosis-stimulated chondrocytes and rats with



adjuvant-induced arthritis by inhibiting ASIC1a expression. *Lab Invest.*
2022 Mar 15.