

## DAFTAR PUSTAKA

- Adinugroho, W. C., Suryadiputra, I. N. N., Saharjo, B. H., & Siboro, L. (2005). *Panduan Pengendalian Kebakaran Hutan*. Bogor: Wetlands International – Indonesia Programme dan Wildlife Habitat Canada.
- Apriyanti, Nur Ridha. 2015. *Pengolahan Citra Digital Landsat 8 dengan Algoritma K-Means Clustering (Studi Kasus: Banjarbaru, Kalimantan Selatan)*. Program S-1 Ilmu Komputer, Universitas Lambung Mangkur Banjarbaru.
- Badarinath, K. V. S., Sharma, A. R., & Kharol, S. K. (2011). Forest fire monitoring and burnt area mapping using satellite data: A study over the forest region of Kerala state, India. *International Journal of Remote Sensing*, 32(1), 85–102. <https://doi.org/10.1080/01431160903439890>
- Badarinath, K. V. S., Sharma, A. R., & Kharol, S. K. (2011). Forest fire monitoring and burnt area mapping using satellite data: A study over the forest region of Kerala state, India. *International Journal of Remote Sensing*, 32(1), 85–102. <https://doi.org/10.1080/01431160903439890>
- Bahadir Kurnaz, Caglar Bayik, Saygin Abdikan. 2020 . Forest Fire Area Detection by Using Landsat-8 and Sentinel-2 Satellite Images: A Case Study in Mugla, Turkey.
- Balogun, A.-L.; Marks, D.; Sharma, R.; Shekhar, H.; Balmes, C.; Maheng, D.; Arshad, A.; Salehi, P. Assessing the Potentials of Digitalization as a Tool for Climate Change Adaptation and Sustainable Development in Urban Centres. *Sustain. Cities Soc.* 2020, 53, 101888.
- Bar, S., Parida, B. R., & Pandey, A. C. (2020). Landsat-8 and Sentinel-2 based Forest fire burn area mapping using machine learning algorithms on GEE cloud platform over Uttarakhand, Western Himalaya. *Remote Sensing Applications: Society and Environment*, 18. <https://doi.org/10.1016/j.rsase.2020.100324>
- Berita Satu. 2019. 2019, Karhutla di Kalbar Terbesar dalam 5 Tahun Terakhir. [Online] URL: [2019, Karhutla di Kalbar Terbesar dalam 5 Tahun Terakhir \(beritasatu.com\)](https://www.beritasatu.com/2019/karhutla-di-kalbar-terbesar-dalam-5-tahun-terakhir)
- Bishop, C.M., 2014. Pattern recognition and machine learning, Antimicrobial agents and chemotherapy. <https://doi.org/10.1128/AAC.03728-14>.
- BMKG. 2019. Prakiraan Musim Hujan Tahun 2018-2019 di Indonesia. [Online] URL: [Prakiraan Musim Hujan Tahun 2018-2019 di Indonesia | BMKG](https://www.bmkg.go.id/prakiraan-musim-hujan-tahun-2018-2019-di-indonesia).
- BNPB. 2019. 99% Penyebab Kebakaran Hutan dan Lahan adalah Ulah Manusia. [Online] URL: <https://bnpb.go.id/berita/99-penyebab-kebakaran-hutan-dan-lahan-adalah-ulah-manusia>
- Bobbe, T.; Finco, M.V.; Quayle, B.; Lannom, K.; Sohlberg, R.; Parsons, A. Field Measurements for the Training and Validation of Burn Severity Maps from Spaceborne, Remotely Sensed Imagery. *USDI Joint. Fire Sci. Program Final Proj. Rep. JFSP RFP 2001*, 2, 1–15.
- Boschetti, L., Roy, D.P., Justice, C.O. and Giglio, L., 2010. Global assessment of the temporal reporting accuracy and precision of the MODIS burned area product. *International Journal of Wildland Fire*, 19(6), pp.705-709.

- Boschetti, M., Stroppiana, D., & Brivio, P. A. (2010). Mapping burned areas in a Mediterranean environment using soft integration of spectral indices from high-resolution satellite images. *Earth Interactions*, 14(17), 1–20. <https://doi.org/10.1175/2010EI349.1>
- Breiman, L. (2001). *Random forests*. *Mach. Learn.* 45, 5–32. doi: 10.1023/A:1010933404324.
- Brivio, P. A., Maggi, E., Binaghi, and I. Gallo, 2003: Mapping burned surfaces in Sub-Saharan Africa based on multi-temporal neural classification. *Int. J. Remote Sens.*, 24, 4003–4018.
- Campbell, J.B. and Wynne, R.H. 2011. *Introduction to Remote Sensing Fifth Edition*. The Guilford Press, New York.
- Chang, Ni-Bin dan Bai, Kaixu. 2018. *Multisensor Data Fusion and Machine Learning for Environmental Remote Sensing*. CRC Press Taylor and Francis Group.
- Chongo, D., R. Nagasawa, A. O. C. Ahmed, and F. Perveen, 2007: Fire monitoring in savana ecosystems using MODIS data: A case study of Kruger National Park, South Africa. *Landscape Ecol. Eng.*, 3, 79–88.
- Cochran, W. G. (1977). *Sampling techniques (3rd ed.)*. New York: John Wiley & Sons.
- Congalton, R. and K. Green. 2009. *Assessing the Accuracy of Remotely Sensed Data: Principles and Practices, 2nd edn*. CRC/Taylor & Francis, Boca Raton, FL, 183pp.
- Congalton, R. G. 1988b. A comparison of sampling schemes used in generating error matrices for assessing the accuracy of maps generated from remotely sensed data. *Photogrammetric Engineering and Remote Sensing*, 54(5), 593–600.
- Crutzen, P.J., Heidt, L.E., Krasnec, J.P., Pollock, W.H., Seiler, W., 1979. Biomass burning as a source of atmospheric gases CO, H<sub>2</sub>, N<sub>2</sub>O, NO, CH<sub>3</sub>Cl and COS. *Nature* 282, 253–256. <https://doi.org/10.1038/282253a0>.
- Danoedoro, P. 2012. *Pengantar Penginderaan Jauh Digital*. Yogyakarta: Penerbit ANDI.
- Eitrich, T., Lang, B., 2006. Efficient optimization of support vector machine learning parameters for unbalanced datasets. *J. Comput. Appl. Math.* 196, 425–436.
- Escuin, S., Navarro, R., Fernandez, P., 2008. Fire severity assessment by using NBR (normalized burn ratio) and NDVI (normalized difference vegetation index) derived from LANDSAT TM/ETM images. *Int. J. Rem. Sens.* 29, 1053–1073. <https://doi.org/10.1080/01431160701281072>.
- Filipponi, F. 2018. BAIS22: Burned Area Index for Sentinel-2. *Proceedings 2018*, 2,364; doi:10.3390/ecrs-2-05177
- Forgy, E. W., 1965. Cluster analysis of multivariate data: Efficiency versus interpretability of classifications. *Biometrics*, 21, 768–769.
- Gaveau, D. L. A., Descals, A., Salim, M. A., Sheil, D., & Sloan, S. (2021). Refined burned-area mapping protocol using Sentinel-2 data increases estimate of 2019 Indonesian burning. *Earth System Science Data*, 13(11), 5353–5368. <https://doi.org/10.5194/essd-13-5353-2021>

- Giglio, L., Boschetti, L., Roy, D.P., Humber, M.L., Justice, C.O., 2018. The Collection 6 MODIS burned area mapping algorithm and product. *Remote Sens. Environ.* <https://doi.org/10.1016/j.rse.2018.08.005>.
- Giglio, L., Randerson, J.T., Van Der Werf, G.R., 2013. Analysis of daily, monthly, and annual burned area using the fourth-generation global fire emissions database (GFED4). *J. Geophys. Res. Biogeosciences*. <https://doi.org/10.1002/jgrg.20042>.
- Giorgos Mountrakis, Jungho Im, Caesar Ogole. 2011. Support vector machines in remote sensing: A review. *ISPRS Journal of Photogrammetry and Remote Sensing*, Volume 66, Issue 3, 2011, Pages 247-259.
- Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R. (2017). Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment*, 202, 18–27. <https://doi.org/10.1016/j.rse.2017.06.031>
- Huntington, J.L.; Hegewisch, K.C.; Daudert, B.; Morton, C.G.; Abatzoglou, J.T.; McEvoy, D.J.; Erickson, T. Climate Engine: Cloud Computing and Visualization of Climate and Remote Sensing Data for Advanced Natural Resource Monitoring and Process Understanding. *Bull. Am. Meteorol. Soc.* 2017, 98, 2397–2410.
- Indah, Lestari & Lalita, Luhurkinanti & Indah, Fitriasari & Harwahyu, Ruki & Sari, Riri. (2020). Machine learning approaches for burned area identification using Sentinel-2 in Central Kalimantan. *Journal of Applied Engineering Science*. 18. 207-215. [10.5937/jaes18-25495](https://doi.org/10.5937/jaes18-25495).
- KLHK. (2019). Rekapitulasi Luas Kebakaran Hutan dan Lahan (Ha) Per Provinsi Di Indonesia Tahun 2014-2019. In SiPongi - Karhutla Monitoring Sistem. Jakarta.
- Kontoes, C. C., H. Poilve', G. Florsch, I. Keramitsoglou, and S. Paralikidis, 2009: A comparative analysis of a fixed thresholding vs. a classification tree approach for operational burn scar detection and mapping. *Int. J. Appl. Earth Obs. Geoinformation*, 11, 299–316.
- Koutsias, N., and M. Karteris, 2000: Burned area mapping using logistic regression modeling of a single post-fire Landsat-5 Thematic Mapper image. *Int. J. Remote Sens.*, 21, 673–687.
- Kurnaz, B., Bayik, C., & Abdikan, S. (2020). *Forest Fire Area Detection by Using Landsat-8 and Sentinel-2 Satellite Images: A Case Study in Mugla, Turkey*. <https://doi.org/10.21203/rs.3.rs-26787/v1>
- Landgrebe, D.A. 2003. *Signal Theory Methods in Multispectral Remote Sensing*. John Wiley and Sons, Hoboken. <http://dx.doi.org/10.1002/0471723800>.
- LAPAN. (2015). Pedoman Pemanfaatan Data LANDSAT-8 untuk Deteksi Daerah Terbakar (Burned Area). Jakarta: Pusat Pemanfaatan Penginderaan Jauh.
- Leeuwen, B. V., Tobal, Z., Kovács, F. 2020. Machine Learning Techniques For Land Use/Land Cover Classification Of Medium Resolution Optical Satellite Imagery Focusing On Temporary Inundated Areas. *Journal of Environmental Geography* 13 (1–2), 43–52.
- Lentile, Leigh & Holden, Zachary & Smith, Alistair & Falkowski, Michael & Hudak, AT & Morgan, Penelope & Lewis, Sarah & Gessler, Paul & Benson, Nate. .2006. Remote sensing techniques to assess active fire characteristics and post-fire effects. *International Journal of Wildland Fire*. 15. [10.1071/WF05097](https://doi.org/10.1071/WF05097).

- Lillesand TM, Kiefer RW. 1997. Penginderaan Jauh dan Interpretasi Citra; diterjemahkan oleh Dulbahri *et al.* Yogyakarta (ID): Gadjah Mada University Press.
- Lillesand, T.M., Kiefer, R.W., dan Chipman, J.W. 2008. Remote Sensing And Image Interpretation. New York : John Willey & Sons Inc.
- Lizundia-Loiola, J., Otón, G., Ramo, R., & Chuvieco, E. (2020). A spatio-temporal active-fire clustering approach for global burned area mapping at 250 m from MODIS data. *Remote Sensing of Environment*, 236. <https://doi.org/10.1016/j.rse.2019.111493>
- M. Ataei, and M. Osanloo. Using a combination of genetic algorithm and the grid search method to determine optimum cutoff grades of multiple metal deposits. *International Journal of Surface Mining, Reclamation and Environment* 18.1 (2004): 60-78.
- Mantero, P., Moser, G., Serpico, S.B., 2005. Partially supervised classification of remote sensing images through SVM-based probability density estimation. *IEEE Transactions on Geoscience and Remote Sensing* 43 (3), 559–570.
- Mariana Belgiu, Lucian Drăguț. 2016. Random forest in remote sensing: A review of applications and future directions. *ISPRS Journal of Photogrammetry and Remote Sensing* Volume 114, Pages 24-31, ISSN 0924-2716. <https://doi.org/10.1016/j.isprsjprs.2016.01.011>.
- McHugh ML. Interrater reliability: the kappa statistic. *Biochem Med (Zagreb)*. 2012;22(3):276-82. PMID: 23092060; PMCID: PMC3900052.
- Mell, P dan Grance, T. 2011. The NIST Definition of Cloud Computing. National Institute of Standards and Technology, Special Publication 800-145.
- Mohri, M., 2012. *Foundations of Machine Learning*. The MIT Press, Massachusetts, USA.
- Morgan, P., Keane, R.E., Dillon, G.K., Jain, T.B., Hudak, A.T., Karau, E.C., Sikkink, P.G., Holden, Z.A., Strand, E.K., 2014. Challenges of assessing fire and burn severity using field measures, remote sensing and modelling. *Int. J. Wildland Fire*. <https://doi.org/10.1071/WF13058>.
- Nugroho, G., Rarasati, A., dan Kushardono. 2019. Penyediaan Informasi Geospasial Berbasis Cloud Computing Data Penginderaan Jauh. *Majalah Inderaja* Vol. X, No. 12, November 2019.
- Olofsson, P., Foody, G.M., Stehman, S.V., & Woodcock, C.E. (2013). Making better use of accuracy data in land change studies: Estimating accuracy and area and quantifying uncertainty using stratified estimation. *Remote Sensing of Environment*, 129, 122–131.
- Parker, B.M.; Lewis, T.; Srivastava, S.K. Estimation and Evaluation of Multi-Decadal Fire Severity Patterns Using Landsat Sensors. *Remote Sens. Environ.* 2015, 170, 340–349.
- Patterson, M. W., and S. R. Yool, 1998: Mapping fire-induced vegetation mortality using Landsat Thematic Mapper data: A comparison of linear transformation techniques. *Remote Sens. Environ.*, 65, 132–142.
- Pereira, José. (2003). Remote sensing of burned areas in tropical savanas. *International Journal of Wildland Fire*. 12. 259-270. 10.1071/WF03028.

- Rahmi, Khalifah & Febrianti, Nur. 2020. Pemanfaatan Data Sentinel-2 untuk Analisis Indeks Area Terbakar (*Burned Area*). *Jurnal Penginderaan Jauh Indonesia Februari 2020, Volume 02, No 01*.
- Ramo, R, García, M, Rodríguez, D, Chuvieco, E, 2018. A data mining approach for global burned area mapping. *International Journal of Applied Earth Observation and Geoinformation*. Volume 73, 2018, Pages 39-51.
- Riswanto E. 2009. Evaluasi akurasi klasifikasi penutupan lahan menggunakan citra Alos Palsar resolusi rendah, Studi kasus di Pulau Kalimantan [Skripsi]. Bogor (ID): Institut Pertanian Bogor.
- Roy, P. S., & Dun, D. (2003). Forest Fire and Degradation Assessment Using Satellite Remote Sensing and Geographic Information System. 361–400.
- Sap, M.N.M., and Kohram, M., (2008): Spectral Angle Based Kernels for the Classification of Hyperspectral Images Using Support Vector Machines. *Second Asia International Conference on Modelling & Simulation, Kuala Lumpur, Malaysia 13-15 May*, pp.559-563.
- Saputra, E., Khaira, U., dan Abidin, Z., 2021. Implementasi Algoritme Support Vector Machines untuk Klasifikasi Area Terbakar di Lahan Gambut. *Jurnal Nasional Teknik Elektro dan Teknologi Informasi*. Vol. 10. No. 1 Februari 2021.
- Scholes, R.J., Kendall, J., Justice, C.O., 1996. The quantity of biomass burned in southern Africa. *J. Geophys. Res. Atmos.*
- Silva De Carvalho, N., José, I., Ferreira, M., Körting, T. S., Oliveira, L. E., de Aragão, C., & Anderson, L. O. (n.d.). *Random Forest And Support Vector Machine Applied For Mapping Burned Areas In Amazon*. <http://earthexplorer.usgs.gov>
- Silva, J.M.N., Sa, A.C.L., Pereira, J.M.C., 2005. Comparison of burned area estimates derived from SPOT-VEGETATION and Landsat ETM<sub>p</sub> data in Africa: influence of spatial pattern and vegetation type. *Remote Sens. Environ.* 96, 188–201. <https://doi.org/10.1016/j.rse.2005.02.004>.  
silvaPress.
- Smith, A. M. S., M. J. Wooster, N. A. Drake, F. M. Dipotso, M. J. Falkowski, and A. T. Hudak, 2005: Testing the potential of multi-spectral remote sensing for retrospectively estimating fire severity in African savana environments. *Remote Sens. Environ.*, 97, 92–115.
- Sumaryati, Andarini, D. F., Cholianawati N., Indrawati, A., 2021. Forest Fire in East Nusa Tenggara during 2015-2019: Comparison to Forest Fire in Kalimantan and Sumatera. *IOP Conf. Ser.: Earth Environ. Sci.* 893 012010.
- Sutanto. 1987. Penginderaan Jauh Dasar II. Yogyakarta : Gadjah Mada University
- Sutanto. 1994. Penginderaan Jauh Jilid I. Yogyakarta: Gadjah Mada University Press.
- Suwarsono. (2012). Daerah Bekas Kebakaran Hutan dan Lahan (Burned Area) di Kalimantan. Universitas Indonesia.
- Talukdar, S., Singha, P., Mahato, S., Shahfahad., Pal, S., Liou, Y., Rahman, A. 2020. Land-Use Land-Cover Classification by Machine Learning Classifiers for Satellite Observations—A Review. *Remote Sens.* 2020, 12, 1135.

- Tamiminia, H.; Salehi, B.; Mahdianpari, M.; Quackenbush, L.; Adeli, S.; Brisco, B. Google Earth Engine for Geo-Big Data Applications: A Meta-Analysis and Systematic Review. *ISPRS J. Photogramm. Remote Sens.* 2020, 164, 152–170.
- Thoha, Achmad & SAHARJO, BAMBANG & Boer, Rizaldi & ARDIANSYAH, MUHAMMAD. (2019). Characteristics and causes of forest and land fires in Kapuas District, Central Kalimantan Province, Indonesia. *Biodiversitas.* 20. 110-117. 10.13057/biodiv/d200113.
- Tian, S. Z. (2016). Random forest classification of wetland landcovers from multisensor data in the arid region of Xinjiang. China. *Xinjiang. China: RemotenSens.* 8 (11), 954.
- TRIGG, S. & Flasse, Stéphane. (2001). An evaluation of different bi-spectral spaces for discriminating burned shrub-savanah. *International Journal of Remote Sensing.* 22. 10.1080/01431160110053185.
- Tucker C (1979) Red and photographic infrared linear combinations for monitoring vegetation. *Remote Sensing of Environment* 8: 127–150.
- Vianna, C. M. Girão, I. Rocha, J. 2019. Training samples from open data for satellite imagery classification: using K-means clustering algorithm. *AGILE 2019 – Limassol, June 17-20, 2019.*
- Warner, T. A., Skowronski, N. S., & Gallagher, M. R. (2017). High spatial resolution burn severity mapping of the New Jersey Pine Barrens with WorldView-3 near-infrared and shortwave infrared imagery. *International Journal of Remote Sensing*, 38(2), 598–616. <https://doi.org/10.1080/01431161.2016.1268739>
- Widodo, Saptono; Hidayatno, Achmad; Isnanto, R Rizal. 2011. Segmentasi Citra Menggunakan Teknik Pemetaan Warna (Color Mapping) dengan Bahasa Pemrograman Delphi. Program Sarjana Teknik Elektro Universitas Diponegoro, Semarang.
- World Bank. 2016. *Kerugian dari Kebakaran Hutan Analisa Dampak Ekonomi dari Krisis Kebakaran Hutan 2015.*
- Xulu, Sifiso, Nkanyiso Mbatha, and Kabir Peerbhay. 2021. "Burned Area Mapping over the Southern Cape Forestry Region, South Africa Using Sentinel Data within GEE Cloud Platform" *ISPRS International Journal of Geo-Information* 10, no. 8: 511. <https://doi.org/10.3390/ijgi10080511>
- Zhu, G., Blumberg, D.G., 2002. Classification using ASTER data and SVM algorithms; The case study of Beer Sheva, Israel. *Remote Sensing of Environment* 80 (2), 233–240.