

DAFTAR PUSTAKA

- Abe, M., Fujii, H., Funakoshi, S., Satoh, A., Kawazoe, M., Maeda, T., Tada, K., Yokota, S., Yamanokuchi, T., Yoshimura, C., Mimata, R., Takahashi, K., Ito, K., Yasuno, T., Kuga, T., Mukoubara, S., Akiyoshi, K., Kawanami, D., Masutani, K., & Arima, H. (2021). Comparison of Body Mass Index and Waist Circumference in the Prediction of Diabetes: A Retrospective Longitudinal Study. *Diabetes Therapy*, 12(10), 2663–2676. <https://doi.org/10.1007/s13300-021-01138-3>
- Anand, S. S., Tarnopolsky, M. A., Rashid, S., Schulze, K. M., Desai, D., Mente, A., Rao, S., Yusuf, S., Gerstein, H. C., & Sharma, A. M. (2011). Adipocyte hypertrophy, fatty liver and metabolic risk factors in south asians: The molecular study of health and risk in ethnic groups (mol-SHARE). *PLoS ONE*, 6(7), 1–8. <https://doi.org/10.1371/journal.pone.0022112>
- Appleton, S. L., Seaborn, C. J., Visvanathan, R., Hill, C. L., Gill, T. K., Taylor, A. W., & Adams, R. J. (2013). Diabetes and cardiovascular disease outcomes in the metabolically healthy obese phenotype: A cohort study. *Diabetes Care*, 36(8), 2388–2394. <https://doi.org/10.2337/dc12-1971>
- Aynalem, S. B., & Zeleke, A. J. (2018). Prevalence of Diabetes Mellitus and Its Risk Factors among Individuals Aged 15 Years and above in Mizan-Aman Town, Southwest Ethiopia, 2016: A Cross Sectional Study. *International Journal of Endocrinology*, 2018. <https://doi.org/10.1155/2018/9317987>
- Bazzano, L. A., He, J., Ogden, L. G., Loria, C. M., Vupputuri, S., Myers, L., & Whelton, P. K. (2002). Fruit and vegetable intake and risk of cardiovascular disease in US adults: The first National Health and Nutrition Examination Survey Epidemiologic Follow-up Study. *American Journal of Clinical Nutrition*, 76(1), 93–99. <https://doi.org/10.1093/ajcn/76.1.93>
- Bazzano, L. A., Li, T. Y., Joshipura, K. J., & Hu, F. B. (2008). Intake of fruit, vegetables, and fruit juices and risk of diabetes in women. *Diabetes Care*, 31(7), 1311–1317. <https://doi.org/10.2337/dc08-0080>
- Bell, J. ., Kivimaki, M., & Hamer, M. (2014). *Metabolically healthy obesity and risk of incident type 2 diabetes a meta-analysis of prospective cohort studies* (pp. 504–515). *Obesity Reviews*. <https://doi.org/10.1111/obr.12157>
- Beydoun, M. A., Fanelli Kuczmarski, M. T., Wang, Y., Mason, M. A., Evans, M. K., & Zonderman, A. B. (2011). Receiver-operating characteristics of adiposity for metabolic syndrome: The Healthy Aging in Neighborhoods of Diversity across the Life Span (HANDLS) study. *Public Health Nutrition*, 14(1), 77–92. <https://doi.org/10.1017/S1368980010002648>
- Bodicoat, D. H., Gray, L. J., Henson, J., Webb, D., Guru, A., Misra, A., Gupta, R., Vikram, N., Sattar, N., Davies, M. J., & Khunti, K. (2014). Body mass index

and waist circumference cut-points in multi-ethnic populations from the UK and India: The ADDITION-Leicester, Jaipur Heart Watch and New Delhi cross-sectional studies. *PLoS ONE*, 9(3). <https://doi.org/10.1371/journal.pone.0090813>

Boeing, H., Bechthold, A., Bub, A., Ellinger, S., Haller, D., Kroke, A., Leschik-Bonnet, E., Müller, M. J., Oberritter, H., Schulze, M., Stehle, P., & Watzl, B. (2012). Critical review: Vegetables and fruit in the prevention of chronic diseases. *European Journal of Nutrition*, 51(6), 637–663. <https://doi.org/10.1007/s00394-012-0380-y>

Breen, C., Ryan, M., McNulty, B., Gibney, M. J., Canavan, R., & O'Shea, D. (2014). High saturated-fat and low-fibre intake: A comparative analysis of nutrient intake in individuals with and without type 2 diabetes. *Nutrition and Diabetes*, 4(2), 1–10. <https://doi.org/10.1038/nutd.2014.2>

Brü, J. C., Dodson Michael, # M, Winnay, J. N., Hayashi, T., Hö, D., Accili, D., Goodyear, L. J., & Kahn, R. (1998). A Muscle-Specific Insulin Receptor Knockout Exhibits Features of the Metabolic Syndrome of NIDDM without Altering Glucose Tolerance (IR) itself are rare, many studies have demonstrated de-creases in insulin receptor number and kinase activity in muscle and other tissues of rodents and humans with early NIDDM (Kahn et al. In *Molecular Cell* (Vol. 2).

Bruin, J. E., Petre, M. A., Raha, S., Morrison, K. M., Gerstein, H. C., & Holloway, A. C. (2008). Fetal and neonatal nicotine exposure in wistar rats causes progressive pancreatic mitochondrial damage and beta cell dysfunction. *PLoS ONE*, 3(10). <https://doi.org/10.1371/journal.pone.0003371>

Bunney, P. E., Zink, A. N., Holm, A. A., Billington, C. J., & Kotz, C. M. (2017). Orexin activation counteracts decreases in nonexercise activity thermogenesis (NEAT) caused by high-fat diet. *Physiology and Behavior*, 176, 139–148. <https://doi.org/10.1016/j.physbeh.2017.03.040>

Carlsson, S., Persson, P. ., & Alvarsson, M. (1999). Low birth weight, family history of diabetes and glucose intolerance in Swedish middle-aged men. *Diabetes Care*, 22, 1043–1047.

Carter, P., Gray, L. J., Troughton, J., Khunti, K., & Davies, M. J. (2010). Fruit and vegetable intake and incidence of type 2 diabetes mellitus: Systematic review and meta-analysis. *BMJ (Online)*, 341(7772), 543. <https://doi.org/10.1136/bmj.c4229>

Chang, S. A. (2012). Smoking and type 2 diabetes mellitus. In *Diabetes and Metabolism Journal* (Vol. 36, Issue 6, pp. 399–403). <https://doi.org/10.4093/dmj.2012.36.6.399>

Chen, G. C., Qin, L. Q., & Ye, J. K. (2014). Leptin levels and risk of type 2 diabetes: Gender-specific meta-analysis. *Obesity Reviews*, 15(2), 134–142. <https://doi.org/10.1111/obr.12088>

- Cheong, K. C., Yusoff, A. F., Ghazali, S. M., Lim, K. H., Selvarajah, S., Haniff, J., Khor, G. L., Shahar, S., Rahman, J. A., Zainuddin, A. A., & Mustafa, A. N. (2013). Optimal BMI cut-off values for predicting diabetes, hypertension and hypercholesterolaemia in a multi-ethnic population. In *Public Health Nutrition* (Vol. 16, Issue 3, pp. 453–459). <https://doi.org/10.1017/S1368980012002911>
- Dai, H., Alsalhe, T. A., Chalhaf, N., Riccò, M., Bragazzi, N. L., & Wu, J. (2020). The global burden of disease attributable to high body mass index in 195 countries and territories, 1990–2017: An analysis of the Global Burden of Disease Study. In *PLoS Medicine* (Vol. 17, Issue 7). Public Library of Science. <https://doi.org/10.1371/journal.pmed.1003198>
- Danaei, G., Finucane, M. M., Lu, Y., Singh, G. M., Cowan, M. J., Paciorek, C. J., Lin, J. K., Farzadfar, F., Khang, Y. H., Stevens, G. A., Rao, M., Ali, M. K., Riley, L. M., Robinson, C. A., & Ezzati, M. (2011). National, regional, and global trends in fasting plasma glucose and diabetes prevalence since 1980: Systematic analysis of health examination surveys and epidemiological studies with 370 country-years and 2·7 million participants. *The Lancet*, 378(9785), 31–40. [https://doi.org/10.1016/S0140-6736\(11\)60679-X](https://doi.org/10.1016/S0140-6736(11)60679-X)
- DeFronzo, R. A. (2004). Pathogenesis of type 2 diabetes mellitus. *Medical Clinics of North America*, 88(4), 787–835. <https://doi.org/10.1016/j.mcna.2004.04.013>
- DeFronzo, R. A., Ferrannini, E., & Simonson, D. C. (1989). Fasting hyperglycemia in non-insulin-dependent diabetes mellitus: Contributions of excessive hepatic glucose production and impaired tissue glucose uptake. *Metabolism*, 38(4), 387–395. [https://doi.org/10.1016/0026-0495\(89\)90129-7](https://doi.org/10.1016/0026-0495(89)90129-7)
- Despres, J. P., Lemieux, S., & Lamarche, B. (1995). The insulin resistance-dyslipidemic syndrome: contribution of visceral and therapeutic implications. *Int J Obes Relat Metab Disord*, 19, 76–86.
- Dimas, A. S., Lagou, V., Barker, A., Knowles, J. W., Mägi, R., Hivert, M. F., Benazzo, A., Rybin, D., Jackson, A. U., Stringham, H. M., Song, C., Fischer-Rosinsky, A., Welløv Boesgaard, T., Grarup, N., Abbasi, F. A., Assimes, T. L., Hao, K., Yang, X., Lecoeur, C., ... Prokopenko, I. (2014). Impact of type 2 diabetes susceptibility variants on quantitative glycemic traits reveals mechanistic heterogeneity. *Diabetes*, 63(6), 2158–2171. <https://doi.org/10.2337/db13-0949>
- Donahue, R. P., & Orchard, T. J. (2004). *Diabetes Mellitus and Macrovascular Complications An epidemiological perspective*. <http://diabetesjournals.org/care/article-pdf/15/9/1141/440991/15-9-1141.pdf>
- Drucker, D. J. (2006). The biology of incretin hormones. *Cell Metab*, 3, 153–165.
- Ekpenyong, C. E., Akpan, U. P., Ibu, J. O., & Nyebuk, D. E. (2012a). Gender and age specific prevalence and associated risk factors of type 2 diabetes mellitus

in Uyo metropolis, South Eastern Nigeria. *Diabetologia Croatica*, 41(1), 17–28.

Ekpenyong, C. E., Akpan, U. P., Ibu, J. O., & Nyebuk, D. E. (2012b). Gender and Age Spesific Prevalence and Assosiated Risk Factors of Type 2 Diabetes Mellitus in Uyo Metropolis, South Eastern Nigeria. In *Diabetologia Croatica*.

Elmadfa, I., Meyer, A., Nowak, V., Hasenegger, V., Putz, P., Verstraeten, R., Remaut-DeWinter, A. M., Kolsteren, P., Dostálová, J., Dlouhý, P., Trolle, E., Fagt, S., Biloft-Jensen, A., Mathiessen, J., Velsing Groth, M., Kambek, L., Gluskova, N., Voutilainen, S., Erkkilä, A., ... Cattaneo, A. (2009). European Nutrition and Health Report 2009. *Annals of Nutrition & Metabolism*, 55 Suppl 2, 1–40. <https://doi.org/10.1080/11026480500350294>

Esser, N., Legrand-Poels, S., Piette, J., Scheen, A. J., & Paquot, N. (2014). Inflammation as a link between obesity, metabolic syndrome and type 2 diabetes. In *Diabetes Research and Clinical Practice* (Vol. 105, Issue 2, pp. 141–150). Elsevier Ireland Ltd. <https://doi.org/10.1016/j.diabres.2014.04.006>

Ewers, B., Trolle, E., Jacobsen, S. S., Vististen, D., Almdal, T. P., Vilsbøll, T., & Bruun, J. M. (2019). Dietary habits and adherence to dietary recommendations in patients with type 1 and type 2 diabetes compared with the general population in Denmark. *Nutrition*, 61, 49–55. <https://doi.org/10.1016/j.nut.2018.10.021>

Faria, A. N., Ribeiro Filho, F. F., Ferreira, S. R. G., & Zanella, M. T. (2002). Impact of visceral fat on blood pressure and insulin sensitivity in hypertensive obese women. *Obesity Research*, 10(12), 1203–1206. <https://doi.org/10.1038/oby.2002.164>

Feero, W. G., Gutmacher, A. E., & Mccarthy, M. I. (2010). Genomic Medicine Genomics, Type 2 Diabetes, and Obesity. In *n engl j med* (Vol. 24, Issue 9).

Finucane, M. M., Stevens, G. A., Cowan, M. J., Danaei, G., Lin, J. K., Paciorek, C. J., Singh, G. M., Gutierrez, H. R., Lu, Y., Bahalim, A. N., Farzadfar, F., Riley, L. M., & Ezzati, M. (2011). National, regional, and global trends in body-mass index since 1980: Systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9·1 million participants. *The Lancet*, 377(9765), 557–567. [https://doi.org/10.1016/S0140-6736\(10\)62037-5](https://doi.org/10.1016/S0140-6736(10)62037-5)

Fontana, L., Eagon, J. C., Trujillo, M. E., Scherer, P. E., & Klein, S. (2007). Systemic Inflammation in Obese Humans. *Diabetes*, 56(April), 1010–1013. <https://doi.org/10.2337/db06-1656>

Fuchsberger, C., Flannick, J., Teslovich, T. M., Mahajan, A., Agarwala, V., Gaulton, K. J., Ma, C., Fontanillas, P., Moutsianas, L., McCarthy, D. J., Rivas, M. A., Perry, J. R. B., Sim, X., Blackwell, T. W., Robertson, N. R., Rayner, N. W., Cingolani, P., Locke, A. E., Tajos, J. F., ... McCarthy, M. I.

- (2016). The genetic architecture of type 2 diabetes. *Nature*, 536(7614), 41–47. <https://doi.org/10.1038/nature18642>
- Gharipour, M., Sarrafzadegan, N., Sadeghi, M., Andalib, E., Talaie, M., Shafie, D., & Aghababaie, E. (2013). Predictors of metabolic syndrome in the Iranian population: Waist circumference, body mass index, or waist to hip ratio? *Cholesterol*, 2013. <https://doi.org/10.1155/2013/198384>
- Giacco, F., & Brownlee, M. (2010). Oxidative stress and diabetic complications. In *Circulation Research* (Vol. 107, Issue 9, pp. 1058–1070). <https://doi.org/10.1161/CIRCRESAHA.110.223545>
- Grefhorst, A., Van Den Beukel, J. C., Van Houten, E. L. A., Steenbergen, J., Visser, J. A., & Themmen, A. P. (2015). Estrogens increase expression of bone morphogenetic protein 8b in brown adipose tissue of mice. *Biology of Sex Differences*, 6(1), 1–13. <https://doi.org/10.1186/s13293-015-0025-y>
- Gregg, E. W., Li, Y., Wang, J., Rios Burrows, N., Ali, M. K., Rolka, D., Williams, D. E., & Geiss, L. (2014). Changes in Diabetes-Related Complications in the United States, 1990–2010. *New England Journal of Medicine*, 370(16), 1514–1523. <https://doi.org/10.1056/nejmoa1310799>
- Guilherme, A., Virbasius, J. V., Puri, V., & Czech, M. P. (2008). Adipocyte dysfunctions linking obesity to insulin resistance and type 2 diabetes. *Nature Reviews Molecular Cell Biology*, 9(5), 367–377. <https://doi.org/10.1038/nrm2391>
- Guo, S. S., Zeller, C., Chumlea, W. C., & Siervogel, R. M. (1999). Aging, body composition, and lifestyle: The Fels Longitudinal Study. *American Journal of Clinical Nutrition*, 70(3), 405–411. <https://doi.org/10.1093/ajcn/70.3.405>
- Haghighatdoost, F., Amini, M., Feizi, A., & Iraj, B. (2017). Are body mass index and waist circumference significant predictors of diabetes and prediabetes risk: Results from a population based cohort study. *World Journal of Diabetes*, 8(7), 365. <https://doi.org/10.4239/wjd.v8.i7.365>
- Hajian-Tilaki, K., & Heidari, B. (2015). Is waist circumference a better predictor of diabetes than body mass index or waist-to-height ratio in Iranian adults? *International Journal of Preventive Medicine*, 2015-January. <https://doi.org/10.4103/2008-7802.151434>
- Haldar, S., Chia, S. C., & Henry, C. J. (2015). Body Composition in Asians and Caucasians: Comparative Analyses and Influences on Cardiometabolic Outcomes. In *Advances in Food and Nutrition Research* (1st ed., Vol. 75). Elsevier Inc. <https://doi.org/10.1016/bs.afnr.2015.07.001>
- Hall, J. N., Moore, S., Harper, S. B., & Lynch, J. W. (2009). Global Variability in Fruit and Vegetable Consumption. *American Journal of Preventive Medicine*, 36(5), 402–409.e5. <https://doi.org/10.1016/j.amepre.2009.01.029>
- Hamer, M., & Chida, Y. (2007). Intake of fruit, vegetables, and antioxidants and

risk of type 2 diabetes: Systematic review and meta-analysis. *Journal of Hypertension*, 25(12), 2361–2369.
<https://doi.org/10.1097/HJH.0b013e3282efc214>

Hamman, R. F. (1992). *Genetic and Environmental Determinants of Non-Insulin-Dependent Diabetes Mellitus (NIDDM)*.

Handschin, C., & Spiegelman, B. M. (2008). The role of exercise and PGC1 α in inflammation and chronic disease. In *Nature* (Vol. 454, Issue 7203, pp. 463–469). Nature Publishing Group. <https://doi.org/10.1038/nature07206>

Harbuwono, D. S., Tahapary, D. L., Edi Tarigan, T. J., & Yunir, E. (2020). New proposed cut-off of waist circumference for central obesity as risk factor for diabetes mellitus: Evidence from the Indonesian Basic National Health Survey. *PLoS ONE*, 15(11 November).
<https://doi.org/10.1371/journal.pone.0242417>

Hirani, V., Zaninotto, P., & Primatesta, P. (2008). Generalised and abdominal obesity and risk of diabetes, hypertension and hypertension-diabetes co-morbidity in England. *Public Health Nutrition*, 11(5), 521–527.
<https://doi.org/10.1017/S1368980007000845>

Holst, J. J., Knop, F. K., & Visbol, T. (2011). Loss of incretin effect is a specific, important, and early characteristic of type 2 diabetes. *Diabetes Care*, 34(2), S251-7.

Hou, X., Lu, J., Weng, J., Ji, L., Shan, Z., Liu, J., Tian, H., Ji, Q., Zhu, D., Ge, J., Lin, L., Chen, L., Guo, X., Zhao, Z., Li, Q., Zhou, Z., Shan, G., Yang, Z., Yang, W., & Jia, W. (2013). Impact of Waist Circumference and Body Mass Index on Risk of Cardiometabolic Disorder and Cardiovascular Disease in Chinese Adults: A National Diabetes and Metabolic Disorders Survey. *PLoS ONE*, 8(3). <https://doi.org/10.1371/journal.pone.0057319>

Hsiung, D. Y., Liu, C. W., Cheng, P. C., & Ma, W. F. (2015). Using non-invasive assessment methods to predict the risk of metabolic syndrome. *Applied Nursing Research*, 28(2), 72–77. <https://doi.org/10.1016/j.apnr.2014.12.001>

Hwang, L. C., Bai, C. H., Sun, C. A., & Chen, C. J. (2012). Prevalence of metabolically healthy obesity and its impacts on incidences of hypertension, diabetes and the metabolic syndrome in Taiwan. *Asia Pacific Journal of Clinical Nutrition*, 21(2), 227–233.

Imamura, F., O'Connor, L., Ye, Z., Mursu, J., Hayashino, Y., Bhupathiraju, S. N., & Forouhi, N. G. (2015). Consumption of sugar sweetened beverages, artificially sweetened beverages, and fruit juice and incidence of type 2 diabetes: Systematic review, meta-analysis, and estimation of population attributable fraction. *BMJ (Online)*, 351, 1–12.
<https://doi.org/10.1136/bmj.h3576>

International Diabetes Federation. (2021). *IDF Diabetes Atlas 10th Edition*.

- Jallut, D., Golay, A., Munger, R., Frascarolo, P., Schutz, Y., Jéquier, E., & Felber, J. P. (1990). Impaired glucose tolerance and diabetes in obesity: A 6-year follow-up study of glucose metabolism. *Metabolism*, 39(10), 1068–1075. [https://doi.org/10.1016/0026-0495\(90\)90168-C](https://doi.org/10.1016/0026-0495(90)90168-C)
- Janghorbani, M., & Amini, M. (2010). Comparison of body mass index with abdominal obesity indicators and waist-to-stature ratio for prediction of type 2 diabetes: The Isfahan diabetes prevention study. *Obesity Research and Clinical Practice*, 4(1). <https://doi.org/10.1016/j.orcp.2009.07.002>
- Jazet, I. M., Piji, H., & Meinders, A. E. (2000). Adipose tissue, an endocrine organ: impart on insulin resistance. *Neth J Med*, 61, 194–212.
- Kahn, B. B., & Flier, J. S. (2000). Obesity and insulin resistance. *Clinical Invest*, 104, 473–481.
- Kaku, K. (2010). Pathophysiology of Type 2 Diabetes and Its Treatment Policy. *JMAJ*, 53(1), 41–46.
- Kautzky-Willer, A., Harreiter, J., & Pacini, G. (2016). Sex and gender differences in risk, pathophysiology and complications of type 2 diabetes mellitus. *Endocrine Reviews*, 37(3), 278–316. <https://doi.org/10.1210/er.2015-1137>
- Kementerian Kesehatan RI. (2019). Laporan Nasional Riskesdas 2018. *Jakarta: Badan Penelitian Dan Pengembangan Kesehatan*.
- Kodama, S., Horikawa, C., Fujihara, K., Heianza, Y., Hirasawa, R., Yachi, Y., Sugawara, A., Tanaka, S., Shimano, H., Iida, K. T., Saito, K., & Sone, H. (2012). Comparisons of the strength of associations with future type 2 diabetes risk among anthropometric obesity indicators, including waist-to-height ratio: A meta-analysis. In *American Journal of Epidemiology* (Vol. 176, Issue 11, pp. 959–969). <https://doi.org/10.1093/aje/kws172>
- Kohli, S., Sniderman, A. D., Tchernof, A., & Lear, S. A. (2010). Ethnic-specific differences in abdominal subcutaneous adipose tissue compartments. *Obesity*, 18(11), 2177–2183. <https://doi.org/10.1038/oby.2010.94>
- Kusumo, M. P., & Kusumawati, W. (2022). *Barriers to Understanding Health Education in Patients with Diabetes Mellitus (DM) in Yogyakarta : Qualitative Study*. 11(1), 1–9.
- Lam, D. W., & D, L. (2015). *Metabolic syndrome*.
- Lê, K. A., Faeh, D., Stettler, R., Ith, M., Kreis, R., Vermathen, P., Boesch, C., Ravussin, E., & Tappy, L. (2006). A 4-wk high-fructose diet alters lipid metabolism without affecting insulin sensitivity or ectopic lipids in healthy humans. *American Journal of Clinical Nutrition*, 84(6), 1374–1379. <https://doi.org/10.1073/pnas.1115183109>
- Lear, S. A., Chockalingam, A., Kohli, S., Richardson, C. G., & Humphries, K. H. (2012). Elevation in cardiovascular disease risk in south asians is mediated

- by differences in visceral adipose tissue. *Obesity*, 20(6), 1293–1300. <https://doi.org/10.1038/oby.2011.395>
- Lee, D. H., Keum, N. N., Hu, F. B., Orav, E. J., Rimm, E. B., Willett, W. C., & Giovannucci, E. L. (2018). Comparison of the association of predicted fat mass, body mass index, and other obesity indicators with type 2 diabetes risk: two large prospective studies in US men and women. *European Journal of Epidemiology*, 33(11), 1113–1123. <https://doi.org/10.1007/s10654-018-0433-5>
- Leone, A., Battezzati, A., Di Lello, S., Ravasenghi, S., Mohamed-Iahdih, B., Saleh, S. M. L., & Bertoli, S. (2020). Dietary habits of Saharawi type II diabetic women living in Algerian refugee camps: Relationship with nutritional status and glycemic profile. *Nutrients*, 12(2), 1–16. <https://doi.org/10.3390/nu12020568>
- Lesser, I. A., Gasevic, D., & Lear, S. A. (2013). The effect of body fat distribution on ethnic differences in cardiometabolic risk factors of Chinese and Europeans. *Applied Physiology, Nutrition and Metabolism*, 38(7), 701–706. <https://doi.org/10.1139/apnm-2012-0125>
- Li, A., Li, Y., Xiang, F., Zhang, Y., Yang, Q., Hu, Z., & Peng, Q. (2019). Optimal Cut-off Points of BMI WC and WHR for Screening of Pre-Diabetes and Diabetes Over 35 Years Old People. *World Journal of Public Health*, 4(1), 1. <https://doi.org/10.11648/j.wjph.20190401.11>
- Liu, C. J. (2012). Prevalence and risk factors for non-alcoholic fatty liver disease in Asian people who are not obese. *Journal of Gastroenterology and Hepatology (Australia)*, 27(10), 1555–1560. <https://doi.org/10.1111/j.1440-1746.2012.07222.x>
- Low, S., Chin, M. C., Stefan, M., Heng, D., & Deurenberg-Yap, M. (2009). Rationale for redefining obesity in asians. *Annals of the Academy of Medicine Singapore*, 38(1), 66–74.
- Mary, G., Caroline, H., Naomi, H., Helen, O., Lorraine, O., Anna, C., & Louisa, E. (2014). Adult Obesity and Type 2 Diabetes About Public Health England. *Public Health England*, 1–39.
- Meisinger, C., Döring, A., Thorand, B., Heier, M., & Löwel, H. (2006). *Body fat distribution and risk of type 2 diabetes in the general population: are there differences between men and women? The MONICA/KORA Augsburg Cohort Study* 13. <https://academic.oup.com/ajcn/article-abstract/84/3/483/4648894>
- Milita, F., Handayani, S., & Setiaji, B. (2021). Kejadian Diabetes Mellitus Tipe II pada Lanjut Usia di Indonesia (Analisis Riskesdas 2018). *Jurnal Kedokteran Dan Kesehatan*, 17(1), 9–20. <https://www.bing.com/search?q=prevalensi+diabetes+melitus+di+indonesia+jurnal&cvid=e26a04e174e441dcb70594134d15936a&aqs=edge..69i57.1757>

6j0j9&FORM=ANAB01&PC=ASTS

- Muscelli, E., Mari, A., & Casolaro, A. (2008). Separate impact of obesity and glucose tolerance on the incretin effect in normal subjects and type 2 diabetic patients. *Diabetes*, 57, 1340–1348.
- Nazare, J. A., Smith, J. D., Borel, A. L., Haffner, S. M., Balkau, B., Ross, R., Massien, C., Alm  ras, N., & Despr  s, J. P. (2012). Ethnic influences on the relations between abdominal subcutaneous and visceral adiposity, liver fat, and cardiometabolic risk profile: The international study of prediction of intra-abdominal adiposity and its relationship with cardiometabolic risk/intra-. *American Journal of Clinical Nutrition*, 96(4), 714–726. <https://doi.org/10.3945/ajcn.112.035758>
- Neeland, I. J., Turer, A. T., Ayers, R., Powell-Wiley, T. M., Vega, G. L., Farzaneh-Far, R., Grundy, S. M., Khera, A., McGuire, D. K., & Lemos, J. A. (2013). Dysfunctional Adiposity and the Risk of Prediabetes and Type 2 Diabetes in Obese Adults. *JAMA*, 308(11), 1150–1159. <https://doi.org/10.1001/2012.jama.11132.Dysfunctional>
- Nordstr  m, A., Hadr  vi, J., Olsson, T., Franks, P. W., & Nordstr  m, P. (2016). Higher prevalence of type 2 diabetes in men than in women is associated with differences in visceral fat mass. *Journal of Clinical Endocrinology and Metabolism*, 101(10), 3740–3746. <https://doi.org/10.1210/jc.2016-1915>
- Oestenson, C. .-G. (2001). *The pathophysiology of type 2 diabetes mellitus: an overview*.
- Perkeni. (2021). *Pedoman Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 di Indonesia*.
- Polak, K., Czyzyk, A., Simoncini, T., & Meczekalski, B. (2017). New markers of insulin resistance in polycystic ovary syndrome. In *Journal of Endocrinological Investigation* (Vol. 40, Issue 1). Springer International Publishing. <https://doi.org/10.1007/s40618-016-0523-8>
- Power, M. L., & Schulkin, J. (2008). Sex differences in fat storage, fat metabolism, and the health risks from obesity: Possible evolutionary origins. *British Journal of Nutrition*, 99(5), 931–940. <https://doi.org/10.1017/S0007114507853347>
- Pramono, L. A., Setiati, S., Soewondo, P., Subekti, I., Adisasmita, A., Kodim, N., & Sutrisna, B. (2010). Prevalence and predictors of undiagnosed diabetes mellitus in Indonesia. *Acta Medica Indonesiana*, 42(4), 216–223.
- Qiao, Q., & Nyamdorj, R. (2010). Is the association of type II diabetes with waist circumference or waist-to-hip ratio stronger than that with body mass index. In *European Journal of Clinical Nutrition* (Vol. 64, Issue 1, pp. 30–34). <https://doi.org/10.1038/ejcn.2009.93>
- Rachmi, C. N., Jusril, H., Ariawan, I., Beal, T., & Sutrisna, A. (2020). Eating

behaviour of Indonesian adolescents: A systematic review of the literature.
Public Health Nutrition, 24(Lmic).
<https://doi.org/10.1017/S1368980020002876>

- Ranasinghe, P., Pigera, A. S. A. D., Ishara, M. H., Jayasekara, L. M. D. T., Jayawardena, R., & Katulanda, P. (2015). Knowledge and perceptions about diet and physical activity among Sri Lankan adults with diabetes mellitus: A qualitative study Health behavior, health promotion and society. *BMC Public Health*, 15(1), 1–10. <https://doi.org/10.1186/s12889-015-2518-3>
- Rd, J. P., Mph, J. A. M., Neil, C. E. O., Rd, M. P. H., Hilmers, D. C., Mph, E. E., Ms, Y. L., Nicklas, T. A., & Mph, D. (2008). Original Article A comparison of the prevalence of the metabolic syndrome in the United States (US) and Korea in young adults aged 20 to 39 years. *Asia Pacific Journal of Clinical Nutrition*, 17(August), 471–482.
- Rocco, E. R., Mory, D. B., Bergamin, C. S., Valente, F., Miranda, V. L., Frederico, B., Calegare, A., Silva, R. Q., & Dib, S. A. (2011). Optimal cutoff points for body mass index, waist circumference and HOMA-IR to identify a cluster of cardiometabolic abnormalities in normal glucose-tolerant Brazilian children and adolescents. In *Arq Bras Endocrinol Metab* (Vol. 55, Issue 8).
- Rodelo, C., Guiberna, A., & Reyes, J. A. (2017). *Molecular mechanisms of insulin action*.
- Rodrigues Graciano, M. F., Valle, M. M. R., Kowluru, A., Curi, R., & Carpinelli, A. R. (2011). Regulation of insulin secretion and production of reactive oxygen species by free fatty acids in pancreatic islets. In *Islets* (Vol. 3, Issue 5, pp. 213–223). <https://doi.org/10.4161/isl.3.5.15935>
- Sakot, Y., & Grill, V. E. (1990). *A 48-hour Lipid Infusion in the Rat Time-Dependently Inhibits Glucose-Induced Insulin Secretion and B Cell Oxidation Through a Process Likely Coupled to Fatty Acid Oxidation** (Vol. 127, Issue 4).
- Sattar, N., & Gill, J. M. R. (2014). Type 2 diabetes as a disease of ectopic fat? *BMC Medicine*, 12(1), 1–6. <https://doi.org/10.1186/s12916-014-0123-4>
- Senadheera, S. P. A. S., Ekanayake, S., & Wanigatunge, C. (2016). Dietary Habits of Type 2 Diabetes Patients: Variety and Frequency of Food Intake. *Journal of Nutrition and Metabolism*, 2016. <https://doi.org/10.1155/2016/7987395>
- Shabnam, A. A., Homa, K., Mohammad Reza, M. T., Bagher, L., Mohammad Hossein, F., & Hamidreza, A. (2012). Cut-off points of waist circumference and body mass index for detecting diabetes, hypercholesterolemia and hypertension according to National Non-Communicable Disease Risk Factors Surveillance in Iran. *Archives of Medical Science*, 8(4), 614–621. <https://doi.org/10.5114/aoms.2012.30284>
- Shai, I., Jiang, R., Manson, J. A. E., Stampfer, M. J., Willett, W. C., Colditz, G.

- A., & Hu, F. B. (2006). Ethnicity, obesity, and risk of type 2 diabetes in women: A 20-year follow-up study. *Diabetes Care*, 29(7), 1585–1590. <https://doi.org/10.2337/dc06-0057>
- Sigit, F. S., Tahapary, D. L., Trompet, S., Sartono, E., Willems Van Dijk, K., Rosendaal, F. R., & De Mutsert, R. (2020). The prevalence of metabolic syndrome and its association with body fat distribution in middle-aged individuals from Indonesia and the Netherlands: A cross-sectional analysis of two population-based studies. *Diabetology and Metabolic Syndrome*, 12(1), 1–11. <https://doi.org/10.1186/s13098-019-0503-1>
- Sinaga, M., Worku, M., Yemane, T., Tegene, E., Wakayo, T., Girma, T., Lindstrom, D., & Belachew, T. (2018). Optimal cut-off for obesity and markers of metabolic syndrome for Ethiopian adults. *Nutrition Journal*, 17(1). <https://doi.org/10.1186/s12937-018-0416-0>
- Sulistyoningrum, E. (2010). Tinjauan molekular dan Aspek Klinis Resistensi Insulin. *Mandala of Health*, 4(2), 131–138.
- Sun, J., Liu, Z., Zhang, Z., Zeng, Z., & Kang, W. (2022). The Correlation of Prediabetes and Type 2 Diabetes With Adiposity in Adults. *Frontiers in Nutrition*, 9(April), 5–10. <https://doi.org/10.3389/fnut.2022.818263>
- Takahashi, M., Shimomura, K., Proks, P., Craig, T., Negishi, M., Akuzawa, M., Hayashi, R., Shimomura, Y., & Kobashi, I. (2009). A Proposal of Combined Evaluation of Waist Circumference and BMI for the Diagnosis of Metabolic Syndrome. In *Endocrine Journal* (Vol. 56, Issue 9).
- Tyrovolas, S., Chalkias, C., Morena, M., Tsiligianni, I., Zeimbekis, A., Gotsis, E., Metallinos, G., Bountziouka, V., Polychronopoulos, E., Lionis, C., & Panagiotakos, D. (2011). Health care access and prevalence of the metabolic syndrome among elders living in high-altitude areas of the mediterranean islands: The MEDIS study. *Review of Diabetic Studies*, 8(4), 468–476. <https://doi.org/10.1900/RDS.2011.8.468>
- Tyrovolas, S., Koyanagi, A., Garin, N., Olaya, B., Ayuso-Mateos, J. L., Miret, M., Chatterji, S., Tobiasz-Adamczyk, B., Koskinen, S., Leonardi, M., & Haro, J. M. (2015). Diabetes mellitus and its association with central obesity and disability among older adults: A global perspective. *Experimental Gerontology*, 64, 70–77. <https://doi.org/10.1016/j.exger.2015.02.010>
- Vetrivel Venkatasamy, V., Pericherla, S., Manthuruthil, S., Mishra, S., & Hanno, R. (2013). Effect of physical activity on insulin resistance, inflammation and oxidative stress in diabetes mellitus. In *Journal of Clinical and Diagnostic Research* (Vol. 7, Issue 8, pp. 1764–1766). <https://doi.org/10.7860/JCDR/2013/6518.3306>
- Willett, W., Manson, J., & Liu, S. (2002). *Glycemic index, glycemic load, and risk of type 2 diabetes* 1–3. 76(April 2001), 274–280.

- Xie, X. T., Liu, Q., Wu, J., & Wakui, M. (2009). Impact of cigarette smoking in type 2 diabetes development. In *Acta Pharmacologica Sinica* (Vol. 30, Issue 6, pp. 784–787). <https://doi.org/10.1038/aps.2009.49>
- Yanbaeva, D. G., Dentener, M. A., Creutzberg, E. C., Wesseling, G., & Wouters, E. F. M. (2007). Systemic effects of smoking. *Chest*, 131(5), 1557–1566. <https://doi.org/10.1378/chest.06-2179>
- Zaccardi, F., Webb, D. R., Yates, T., & Davies, M. J. (2016). Pathophysiology of type 1 and type 2 diabetes mellitus: A 90-year perspective. *Postgraduate Medical Journal*, 92(1084), 63–69. <https://doi.org/10.1136/postgradmedj-2015-133281>
- Zhang, F. L., Ren, J. X., Zhang, P., Jin, H., Qu, Y., Yu, Y., Guo, Z. N., & Yang, Y. (2021). Strong Association of Waist Circumference (WC), Body Mass Index (BMI), Waist-to-Height Ratio (WHtR), and Waist-to-Hip Ratio (WHR) with Diabetes: A Population-Based Cross-Sectional Study in Jilin Province, China. *Journal of Diabetes Research*, 2021. <https://doi.org/10.1155/2021/8812431>
- Zhang, L., Curhan, G. C., Hu, F. B., Rimm, E. B., & Forman, J. P. (2011). Association between passive and active smoking and incident type 2 diabetes in women. *Diabetes Care*, 34(4), 892–897. <https://doi.org/10.2337/dc10-2087>
- Zhang, Y., Gu, Y., Wang, N., Zhao, Q., Ng, N., Wang, R., Zhou, X., Jiang, Y., Wang, W., & Zhao, G. (2019). Association between anthropometric indicators of obesity and cardiovascular risk factors among adults in Shanghai, China. *BMC Public Health*, 19(1), 1–9. <https://doi.org/10.1186/s12889-019-7366-0>