

DAFTAR PUSTAKA

- Aldujeli A, Hamadeh A, Briedis K, et al. 2020. 'Delays in presentation in patients with acute myocardial infarction during the COVID-19 pandemic'. *Cardiol Res* 2020;11:386–91, diakses 5 Januari 2022
<https://doi.org/10.14740/cr1175>; PMID: 33224384.
- Anand SS, Islam S, Rosengren, et al, INTERHEART Investigators. Risk factors for myocardial infarction in women and men: insights from the INTERHEART study. *Eur Heart J* 2008;29:932-40.
[doi:10.1093/eurheartj/ehn018](https://doi.org/10.1093/eurheartj/ehn018)
- Antman EM, Anbe DT, Armstrong PW, et al. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction: A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2004;44:e1–211.
- Antoniucci D, Valenti R, Migliorini A, et al. 2002. 'Relation of time to treatment and mortality in patients with acute myocardial infarction undergoing primary coronary angioplasty'. *Am J Cardiol* 2002;89:1248–52, diakses 27 Desember 2021
[https://doi.org/10.1016/s0002-9149\(02\)02320-2](https://doi.org/10.1016/s0002-9149(02)02320-2); PMID: 12031722.
- Berger PB, Ellis SG, Holmes DR, et al. 1999. 'Relationship between delay in performing direct coronary angioplasty and early clinical outcome in patients with acute myocardial infarction.' *Circulation* 1999;100:14–20, diakses 28 Desember 2021
<https://doi.org/10.1161/01.cir.100.1.14>; PMID: 10393675
- Bots SH, Peters SAE, Woodward M. Sex differences in coronary heart disease and stroke mortality: a global assessment of the effect of ageing between 1980 and 2010. *BMJ Global Health* 2017;2:e000298.
[doi:10.1136/bmjgh-2017-000298](https://doi.org/10.1136/bmjgh-2017-000298)
- Brodie BR, Gersh BJ, Stuckey T, et al. 2010. 'When is door-to-balloon time critical? Analysis from the HORIZONS-AMI (Harmonizing Outcomes with Revascularization and Stents in Acute Myocardial Infarction) and CADILLAC (Controlled Abciximab and Device Investigation to Lower Late Angioplasty Complications) trials'. *J Am Coll Cardiol* 2010;56:407–13, diakses 28 Desember 2021.
<https://doi.org/10.1016/j.jacc.2010.04.020>; PMID: 20650362.
- Cannon CP, Gibson CM, Lambrew CT, et al. 2000. 'Relationship of symptom-onset-to-balloon time and door-to-balloon time with mortality in patients undergoing angioplasty for acute myocardial infarction'. *JAMA*

2000;283:2941–7, diakses 27 Desember 2021

<https://doi.org/10.1001/jama.283.22.2941>; PMID: 10865271.

Carstensen S, Nelson GC, Hansen PS, Macken L, Irons S, Flynn M, Kovoov P, Soo Hoo SY, Ward MR, Rasmussen HH. Field triage to primary angioplasty combined with emergency department bypass reduces treatment delays and is associated with improved outcome. *Eur Heart J*. 2007;28:2313–2319

Chew NW, Sia CH, Wee HL, et al. 2021. ‘Impact of the COVID-19 pandemic on door-to-balloon time for primary percutaneous coronary intervention – results from the Singapore Western STEMI network’. *Circ J* 2021;85:139–49, diakses 5 Januari 2022
<https://doi.org/10.1253/circj.CJ-20-0800>; PMID: 33162491.

Çinier G, Hayiroğlu M, Pay L, et al. 2020. ‘Effect of the COVID-19 pandemic on access to primary percutaneous coronary intervention for ST-segment elevation myocardial infarction’. *Turk Kardiyol Dern Ars* 2020;48:640–5, diakses 5 Januari 2022
<https://doi.org/10.5543/tkda.2020.95845>; PMID: 33034585.

Claeys MJ, Argacha JF, Collart P, et al. 2020. ‘Impact of COVID-19- related public containment measures on the ST elevation myocardial infarction epidemic in Belgium: a nationwide, serial, cross-sectional study’. *Acta Cardiol* 2020:1–7, diakses 5 Januari 2022
<https://doi.org/10.1080/00015385.2020.1796035>; PMID: 32727305.

Daoulah A, Hersi AS, Al-Faifi SM, et al. 2020. ‘STEMI and COVID-19 pandemic in Saudi Arabia’. *Curr Probl Cardiol* 2020:100656, diakses 5 Januari 2022
<https://doi.org/10.1016/j.cpcardiol.2020.100656>; PMID: 32839042.

De Luca G, Suryapranata H, Ottervanger JP, Antman EM. 2004. ‘Time delay to treatment and mortality in primary angioplasty for acute myocardial infarction’. *Circulation* 2004;109:1223–5, diakses 28 Desember 2021
<https://doi.org/10.1161/01.cir.0000121424.76486.20>; PMID: 15007008.

Dharma S, Iwan D, Isman F, Siska SD, Dian Z, et al. 2021. ‘Performance of Primary Angioplasty for STEMI during the COVID-19 Outbreak’. *Int J Angiol* 2021;30:148-154, diakses 1 Juni 2022
<https://doi.org/10.1055/s-0041-1727133>. ISSN 1061-1711.

Dhungel S, Rabi M, Chandramani A, Arun M, Rajib R, Ranjit S, Himanshu N, Binaya R, Dharmanath Y. Door-to-balloon time and the determining factors in a tertiary cardiac center in Nepal. *Indian Heart Journal*. 2018;70:S309–S312.
<https://doi.org/10.1016/j.ihj.2018.07.011>

Dhruva VN, Abdelhadi SI, Anis A, Gluckman W, Hom D, Dougan W, Kaluski E, Haider B, Klapholz M. ST-Segment Analysis Using Wireless Technology in Acute Myocardial Infarction (STAT-MI) trial. *J Am Coll Cardiol*.

- Erol MK, Kayıkçioğlu M, Kılıçkap M, et al. 2020. 'Treatment delays and in-hospital outcomes in acute myocardial infarction during the COVID-19 pandemic: a nationwide study'. *Anatol J Cardiol* 2020;24:334–42, diakses 5 Januari 2022
<https://doi.org/10.14744/AnatolJCardiol.2020.98607>; PMID: 33122486.
- Fileti L, Vecchio S, Moretti C, et al. 2020. 'Impact of the COVID-19 pandemic on coronary invasive procedures at two Italian high-volume referral centers'. *J Cardiovasc Med (Hagerstown)* 2020;21:869–73, diakses 5 Januari 2022
<https://doi.org/10.2459/JCM.0000000000001101>; PMID: 33009170.
- Fu XY, Shen XF, Cheng YR, et al. 2021. 'Effect of COVID-19 outbreak on the treatment time of patients with acute ST-segment elevation myocardial infarction'. *Am J Emerg Med* 2021;44:192–7, diakses 5 Januari 2022
<https://doi.org/10.1016/j.ajem.2020.09.038>; PMID: 33039221.
- Garcia S, Stanberry L, Schmidt C, et al. 2020. 'Impact of COVID-19 pandemic on STEMI care: An expanded analysis from the United States'. *Catheter Cardiovasc Interv* 2020, diakses 5 Januari 2022
<https://doi.org/10.1002/ccd.29154>; PMID: 32767652.
- Hammad TA, Parikh M, Tashtish N, et al. 2021. 'Impact of COVID-19 pandemic on ST-elevation myocardial infarction in a non-COVID-19 epicenter'. *Catheter Cardiovasc Interv* 2021;97:208–14, diakses 5 Januari 2022
<https://doi.org/10.1002/ccd.28997>; PMID: 32478961.
- Huxley RR, Woodward M. Cigarette smoking as a risk factor for coronary heart disease in women compared with men: a systematic review and meta-analysis of prospective cohort studies. *Lancet* 2011;378:1297-305.
[doi:10.1016/S0140-6736\(11\)60781-2](https://doi.org/10.1016/S0140-6736(11)60781-2)
- Khrisnan VS, Manoj R, et al. 'Door to balloon time in patients presenting with acute ST elevation myocardial infarction and time factors influencing it; an observational study from a tertiary care teaching hospital in India'. *Indian Heart Journal* 73 (2021) 359—361, diakses 10 Desember 2021
<https://doi.org/10.1016/j.ihj.2021.02.008>
- Kumar A, Cannon CP. Acute Coronary Syndromes: Diagnosis and Management, Part I. *Mayo Clinic proceedings*. 2009;84(10):917-38.
- Kwok CS, Gale CP, Kinnaird T, et al. 'Impact of COVID-19 on percutaneous coronary intervention for ST-elevation myocardial infarction'. *Heart* 2020;106:1805–11, diakses 5 Januari 2022
<https://doi.org/10.1136/heartjnl-2020-317650>; PMID: 32868280.
- Lambert L, Brown K, Segal E, et al. 2010. 'Association between timeliness of

reperfusion therapy and clinical outcomes in ST-elevation myocardial infarction'. *JAMA* 2010;303:2148–55, diakses 28 Desember 2021
<https://doi.org/10.1001/jama.2010.712>; PMID: 20516415.

Lee KD, Lee SB, Lim JK, et al. 2020. 'Providing essential clinical care for non-COVID-19 patients in a Seoul metropolitan acute care hospital amidst ongoing treatment of COVID-19 patients'. *J Hosp Infect* 2020;106:673–7, diakses 6 Januari 2022
<https://doi.org/10.1016/j.jhin.2020.09.031>; PMID: 33011308.

Lu Q, Wang JL, Chen ZN, Fu WB, Liu HJ, Ding SF. The impact of COVID-19 on short-term prognosis of ST-segment elevation myocardial infarction patients receiving primary percutaneous coronary intervention in Wuhan China. *Rev Cardiovasc Med.* 2021;22(1):247-56.

Lubovich A, Hamood H, Behar S, Rosenchein U. Bypassing the emergency room to reduce door-to-balloon time and improve outcomes of patients with ST elevation myocardial infarction: the Acute Coronary Syndrome Israeli Survey experience. *Isr Med Assoc J.* 2011; 13:216 – 219.

Mathews R, Peterson ED, Li S, Roe MT, Glickman SW, Wiviott SD, Saucedo JF, Antman EM, Jacobs AK, Wang TY. Use of emergency medical service transport among patients with ST-segment-elevation myocardial infarction: findings from the National Cardiovascular Data Registry Acute Coronary Treatment Intervention Outcomes Network Registry-Get With The Guidelines. *Circulation.* 2011;124:154–163.

McCabe JM, Ehrin JA, Kurt SH, Prashant DB, et al. Impact of Door-to-Activation Time on Door-to-Balloon Time in Primary Percutaneous Coronary Intervention for ST-Segment Elevation Myocardial Infarctions. *Circulation: Cardiovascular Quality and Outcomes, American Heart Association.* 2012. doi: <https://doi.org/10.1161/CIRCOUTCOMES.112.966382>

Menees DS, Peterson ED, Wang Y, et al. Door-to-Balloon time with primary percutaneous coronary intervention for acute myocardial infarction impacts late cardiac mortality in high-risk patients and patients presenting early after the onset of symptoms. *J Am Coll Cardiol.* 2006;47:289 – 295, diakses 2 Juni 2022.

Mously H, Nischay S, Zachary Z, Ibrahim A, Adham K, Rahul J, Setevn J, Daniel I, Mehdi S, Farshad F. Door-to-balloon Time for ST-elevation MI in the Coronavirus Disease 2019 Era, *USC Volume 15* 2021, diakses 23 Maret 2022. <https://doi.org/10.15420/usc.2021.05>

Murat Bektas, Eylem Kivanc, Selda Murat. Gender differences in clinical characteristics and in-hospital and one-year outcomes of young patients with ST-segment elevation myocardial infarction under the age of 40. *J Cardiovasc Thorac Res.* 2021; 13(2): 116–124. Published online 2021 Feb 8.

- Nallamothu BK, Bates ER. 2003. 'Percutaneous coronary intervention versus fibrinolytic therapy in acute myocardial infarction: is timing (almost) everything?'. *Am J Cardiol* 2003;92:824–6, diakses 28 Desember 2021
[https://doi.org/10.1016/s0002-9149\(03\)00891-9](https://doi.org/10.1016/s0002-9149(03)00891-9); PMID: 14516884
- Nishiga M, Wang DW, Han Y, et al. 2020, 'COVID-19 and Cardiovascular Disease: From Basic Mechanisms to Clinical Perspectives'. *Nature Reviews Cardiology*, diakses 5 Januari 2022
<https://doi.org/10.1038/s41569-020-0413-9>
- Tim Penyusunan Panduan Praktik Klinis Tata Laksana Kasus dan Prosedur Tindakan, Tim Penanganan Kejadian Luar Biasa (KLB) Coronavirus Disease 2019 (COVID-19), Rumah Sakit Jantung dan Pembuluh Darah Harapan Kita. 2020. Panduan Praktik Klinis (PPK) Tata Laksana Kasus dan Prosedur Tindakan COVID-19 Kardiovaskular RSJPDHK tahun 2020. ISBN 978-623-6659-67-0.
- Park J, Choi KH, Lee JM, et al. 2019. 'Prognostic implications of door-to-balloon time and onset-to-door time on mortality in patients with ST-segment-elevation myocardial infarction treated with primary percutaneous coronary intervention'. *J Am Heart Assoc* 2019;8:e012188, diakses 28 Desember 2021
<https://doi.org/10.1161/jaha.119.012188>; PMID: 31041869.
- Perhimpunan Dokter Spesialis Kardiovaskular Indonesia (PERKI). 2020. 'Panduan Diagnosis dan tatalaksana Penyakit Kardiovaskular pada Pandemi COVID-19' Edisi Pertama 9 Mei 2020, diakses 10 Januari 2022 secara *online* di <https://inaheart.org/wp-content/uploads/2021/07/FINAL-Panduan-Diagnosis-dan-Tatalaksana-Penyakit-Kardiovaskular-pada-PANDEMI-COVID-19.pdf>
- Peters SAE, Huxley RR, Woodward M. Diabetes as risk factor for incident coronary heart disease in women compared with men: a systematic review and meta-analysis of 64 cohorts including 858,507 individuals and 28,203 coronary events. *Diabetologia* 2014;57:1542-51.
[doi:10.1007/s00125-014-3260-6](https://doi.org/10.1007/s00125-014-3260-6)
- Pinto DS, Kirtane AJ, Nallamothu BK, et al. 2006. 'Hospital delays in reperfusion for ST-elevation myocardial infarction'. *Circulation* 2006;114:2019–25, diakses 27 Desember 2021
<https://doi.org/10.1161/circulationaha.106.638353>; PMID: 17075010.
- Rathore SS, Curtis JP, Chen J, et al. Association of door-to-balloon time and mortality in patients admitted to hospital with ST elevation myocardial infarction: national cohort study. *BMJ (Clin Res Ed)*. 2009;338:b1807, diakses 4 Juni 2022.
<https://doi.org/10.1136/bmj.b1807>.

- Reinstadler SJ, Reindl M, Lechner I, et al. 2020. 'Effect of the COVID-19 pandemic on treatment delays in patients with ST-segment elevation myocardial infarction'. *J Clin Med* 2020;9:2183, diakses 5 Januari 2022
<https://doi.org/10.3390/jcm9072183>; PMID: 32664309.
- Roffi M, Guagliumi G, Ibanez B. 2020. 'The obstacle course of reperfusion for ST-segment-elevation myocardial infarction in the COVID-19 pandemic'. *Circulation* 2020;141:1951–3, diakses 5 Januari 2022
<https://doi.org/10.1161/CIRCULATIONAHA.120.047523>; PMID: 32315205.
- Roffi M, Patrono C, Collet J-P, Mueller C, Valgimigli M, Andreotti F, et al. 2015 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation Task Force for the Management of Acute Coronary Syndromes in Patients Presenting without Persistent ST-Segment Elevation of the European Society of Cardiology (ESC). *European Heart Journal*. 2016;37(3):267-315.
- Sanati HR, Mahjoob MP, Zahedmehr A, et al. Risk Factors of Reperfusion Failure following Primary Angioplasty for ST-Segment Elevation Myocardial Infarction (STEMI). *J Tehran Heart Cent*. 2013;8(3):146-151.
- Secco GG, Zocchi C, Parisi R, et al. 2020. 'Decrease and delay in hospitalization for acute coronary syndromes during the 2020 SARS-CoV-2 pandemic'. *Can J Cardiol* 2020;36:1152–5.
<https://doi.org/10.1016/j.cjca.2020.05.023>; PMID: 32447060.
- Secco GG, Marinucci L, Uguccioni L, Parisi R, Uguccioni S, Fattori R. Transradial versus transfemoral approach for primary percutaneous coronary interventions in elderly patients. *J Invasive Cardiol*. 2013;25(5):254 – 256, diakses 2 Juni 2022.
- Sim WJ, Ang AS, Tan MC, Xiang WW, Foo D, Kong Loh K, et al. Causes of delay in door-to-ballon time in south-east Asian patients undergoing primary percutaneous coronary intervention subjects and methods. 2009; Available from: <https://doi.org/10.1371/journal.pone.0185186>.
- Smith M, Singh A, McElroy D, Mittal S, Pham R. Challenges in managing ST elevation myocardial infarction during the COVID-19 pandemic. *World J Cardiol*. 2021 Apr 26;13(4):76. Available from: [/pmc/artciles/PMC8069518](http://pmc/artciles/PMC8069518).
- Steg PG, James SK, Atar D, Badano LP, Lundqvist CB, Borger MA, et al. ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. The Task Force on the management of ST-segment elevation acute myocardial infarction of the European Society of Cardiology (ESC). *European Heart Journal*. 2012;33(20):2569-619.
- Sugiyono. 2018. Metode Penelitian Kuantitatif Kualitatif dan R&D. Bandung:

Alfabeta.

- Tan W, Parikh RV, Chester R, et al. 2020. 'Single center trends in acute coronary syndrome volume and outcomes during the COVID-19 pandemic'. *Cardiol Res* 2020;11:256–9, diakses 5 Januari 2022
<https://doi.org/10.14740/cr1096>; PMID: 32595811.
- Vavalle JP, Granger CB. The need for regional integrated care for ST-segment elevation myocardial infarction. *Circulation*. 2011;124: 851–856
- Victor SM, Gnanaraj A, V S, Pattabiram S, Mullasari AS. Door-to-balloon: where do we lose time? Single centre experience in India. *Indian Heart J*. 2012; 64: 582 – 587, diakses 2 Juni 2022.
<https://doi.org/10.1016/j.ihj.2012.09.007>.
- Yerasi C, Case BC, Forrestal BJ, et al. 2020. 'Treatment of ST-Segment Elevation Myocardial Infarction During COVID-19 Pandemic'. *Cardiovasc Revasc Med* 2020;21:1024-9, diakses 5 Januari 2022
<https://doi.org/10.1016/j.carrev.2020.05.027>; PMID: 32471712.