

## DAFTAR PUSTAKA

- Al-Kherb, W.A. (2014). Virulence bio-assay efficiency of *Beauveria bassiana* and *Metarhizium anisopliae* for the biological control of *Spodoptera exigua* Hübner (Lepidoptera: Noctuidae) eggs and the 1<sup>st</sup> instar larvae. *Australian Journal of Basic and Applied Sciences*, 8(3), 313–323.
- Al-Momani, F., & Obeidat, M. (2013). Ecology, toxicity, and hydrolytic activities of *Bacillus thuringiensis* in forests. *Turkish Journal of Agriculture and Forestry*, 37(1), 76–82. <https://doi.org/10.3906/tar-1104-33>
- Awmack, C. S., & Leather, S. R. (2002). Host plant quality and fecundity in herbivorous insects. *Annual Review of Entomology*, 47(1), 817–844. <https://doi.org/10.1146/annurev.ento.47.091201.145300>
- Azidah, A. A., & Sofian-Azirun, M. (2006). Fecundity study of *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae) on various host plants. *Journal of Entomology*, 3(3), 261–266. <https://dx.doi.org/10.3923/je.2006.261.266>
- Bechtel, D.B & Bulla, L.A. (1976). Electron microscope study of sporulation and parasporal crystal formation in *Bacillus thuringiensis*. *J Bacteriol* 127:1472–1481. <https://doi.org/10.1128%2Fjb.127.3.1472-1481.1976>
- Bizzarri, M. F., & Bishop, A. H. (2008). The ecology of *Bacillus thuringiensis* on the phylloplane: colonization from soil, plasmid transfer, and interaction with larvae of *Pieris brassicae*. *Microbial Ecology*, 56(1), 133–139. <https://doi.org/10.1007/s00248-007-9331-1>
- Borel, B. (2017). CRISPR, Microbes and more are joining the war against crop killers. *Nature*. 543:302–304. <https://doi.org/10.1038/543302a>
- Badan Pusat Statistik, Statistik Pertanian Hortikultura SPH-SBS/BPS-Statistics Indonesia (2018). (<https://magelangkab.bps.go.id/statistictable/2019/11/01/750/produksi-tanaman-sayuran-menurut-kecamatan-dan-jenis-tanaman-di-kabupaten-magelang-kuintal-2018.html>). Diakses pada tanggal 25 Agustus 2021, pukul 08.45 WIB.
- Badan Pusat Statistika RI. (2020). Distribusi perdagangan komoditas bawang merah Indonesia. *Subdirektorat statistik perdagangan dalam negeri*. Hal. 20–21
- Badan Pusat Statistika Provinsi Jawa Tengah. (2020). Luas panen dan produksi bawang merah 2018-2020. (<https://jateng.bps.go.id/indicator/55/727/1/luas-panen-dan-produksi-bawang-merah.html>). Diakses pada tanggal 25 Agustus 2021, pukul 08.45 WIB.
- Brahma, D., Swargiary, A., & Dutta, K. (2015). A comparative study on morphology and rearing performance of *Samia ricini* and *Samia canningi* crossbreed with reference to different food plants. *Journal of Entomology and Zoology Studies*. 3(5), 12–19.
- Bravo, A., Gill, S. S., & Soberon, M. (2007). Mode of action of *Bacillus thuringiensis* Cry and Cyt toxins and their potential for insect control. *Toxicon*, 49(4), 423–435. <https://doi.org/10.1016%2Fj.toxicon.2006.11.022>
- Bravo, A., Likitvivatanavong, S., Gill, S. S., & Soberon, M. (2011). *Bacillus*

- thuringiensis*: A story of a successful bioinsecticide. *Insect Biochemistry and Molecular Biology*, 41(7), 423–431. <https://doi.org/10.1016/j.ibmb.2011.02.006>
- Burton, S. L., Ellar, D. J., Li, J., & Derbyshire, D. J. (1999). N-acetylgalactosamine on the putative insect receptor aminopeptidase N is recognised by a site on the domain III lectin-like fold of a *Bacillus thuringiensis* insecticidal toxin. *Journal of Molecular Biology*, 287(5), 1011–1022. <https://doi.org/10.1006/jmbi.1999.2649>
- CABI (2022) *Spodoptera exigua*. Invasive species compendium. Distribution maps of plant pests. CAB International, Nosworthy Way, Wallingford, Oxfordshire, OX10 8DE UK. <https://doi.org/10.1079/DMPP/20056600302>
- CABI (2022) *Samia ricini*. Invasive species compendium. Distribution table. <https://www.cabi.org/isc/datasheet/48272#todistributionDatabaseTable>
- Castro, A. A., Legaspi, J. C., Tavares, W.S., Meagher, R. L., Miller, N., Kanga, L., Haseeb, M., Serrao, J.E., Wilcken, C.F., & Zanuncio, J.C. (2018). Lethal and behavioral effects of synthetic and organic insecticides on *Spodoptera exigua* and its predator *Podisus maculiventris*. *Plos One*. 13(11): e0206789. <https://doi.org/10.1371/journal.pone.0206789>
- Chen, R., Ren, X., & Han, Z. (2014). A cadherin-like protein from the beet armyworm *Spodoptera exigua* (Lepidoptera : Noctuidae ) is a putative CryIaC receptor. 86(1), 58–71. <https://doi.org/10.1002/arch.21163>
- Chen, Y., Deng, Y., Wang, J., Cai, J., & Ren, G. (2004). Characterization of melanin produced by a wild-type strain of *Bacillus thuringiensis*. *J Gen Appl Microbiol*. 50(4), 183-8. <https://doi.org/10.2323/jgam.50.183>
- Cohen, E., Rozen, H., Joseph, T., Braun, S., & Margulies, L. (1991). Photoprotection of *Bacillus thuringiensis kurstaki* from ultraviolet irradiation. *Journal of Invertebrate Pathology*, 57(3), 343–351. [https://doi.org/10.1016/0022-2011\(91\)90138-G](https://doi.org/10.1016/0022-2011(91)90138-G)
- Crickmore, N., Berry, C., Panneerselvam, S., Mishra, R., Connor, T. R., & Bonning, B. C. (2020). A structure-based nomenclature for *Bacillus thuringiensis* and other bacteria-derived pesticidal proteins. *Journal of Invertebrate Pathology*, 186(5), 107438. <https://doi.org/10.1016/j.jip.2020.107438>
- Dai, H., Zhang, G., & Zhang, W. (2017). Temperature dependent development parameters and population life table of beet armyworm, *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae). *Arthropods*, 6(4), 117–125.
- Dash, R., Mandal, M., Ghosh, S. K., & Kundu, S. C. (2008). Silk sericin protein of tropical tasar silkworm inhibits UVB-induced apoptosis in human skin keratinocytes. *Molecular and Cellular Biochemistry*, 311(1–2), 111–119. <https://doi.org/10.1007/s11010-008-9702-z>
- Dean, D. H., Rajamohan, F., Lee, M. K., Wu, S. J., Chen, X. J., Alcantara, E., & Hussain, S. R. (1996). Probing the mechanism of action of *Bacillus thuringiensis* insecticidal proteins by site-directed mutagenesis - A minireview. *Gene*, 179(1), 111–7. [https://doi.org/10.1016/S0378-1119\(96\)00442-8](https://doi.org/10.1016/S0378-1119(96)00442-8)
- Deedat, Y. D. (1994). Mini review problems associated with the use of pesticides : an overview. *Insect Science and Its Application*. 15(3), 247–251.

doi:10.1017/S1742758400017537

- Deka, M., Dutta, S., & Devi, D. (2011). Impact of feeding of *Samia cynthia ricini* Boisduval (red variety) (Lepidoptera: Saturniidae) in respect of larval growth and spinning. *International Journal of Pure and Applied Sciences and Technology*, 5(2), 131–140.
- Dulmage, H. T., & Martinez, E. (1973). The effects of continuous exposure to low concentrations of the  $\delta$ -endotoxin of *Bacillus thuringiensis* on the development of the tobacco budworm, *Heliothis virescens*. *Journal of Invertebrate Pathology*, 22(1), 14–22. [https://doi.org/10.1016/0022-2011\(73\)90004-9](https://doi.org/10.1016/0022-2011(73)90004-9)
- El-Sharkawey, A. Z., Ragaei, M., Sabbour, M., Afaf., Mohamed, H. A. L. A., & Samy, R. (2009). Laboratory evaluation of antioxidants as UV-protectants for *Bacillus thuringiensis* against potato tuber moth larvae. *Australian Journal of Basic and Applied Sciences*, 3(2), 358–370.
- Farahani, S., Talebi, A. A., & Fathipour, Y. (2011). Life cycle and fecundity of *Spodoptera exigua* (Lep.: Noctuidae) on five soybean varieties. *Journal of Entomological Society of Iran*, 30(2), 1–12.
- Frankenhuyzen, K. (2009). Insecticidal activity of *Bacillus thuringiensis* crystal proteins. *Journal of Invertebrate Pathology*, 101(1), 1–16. <https://doi.org/10.1016/j.jip.2009.02.009>
- Greenberg, S. M., Sappington, T. W., Legaspi, B. C., Liu, T. X., & Sétamou, M. (2001). Feeding and life history of *Spodoptera exigua* (Lepidoptera: Noctuidae) on different host plants. *Annals of the Entomological Society of America*, 94(4), 566–575. [https://doi.org/10.1603/0013-8746\(2001\)094\[0566:FALHOS\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2001)094[0566:FALHOS]2.0.CO;2)
- Griego, V. M., & Spence, K. D. (1978). Inactivation of *Bacillus thuringiensis* spores by ultraviolet and visible light. *Applied and Environmental Microbiology*, 35(5), 906–910. <https://doi.org/10.1128/aem.35.5.906-910.1978>
- Hart, S. J., Terray, A., Leski, T. A., Arnold, J., & Stroud, R. (2006). Discovery of a significant optical chromatographic difference between spores of *Bacillus anthracis* and its close relative, *Bacillus thuringiensis*. *Analytical Chemistry*, 78(9), 3221–3225. <https://doi.org/10.1021/ac052221z>
- Heckel, D. G. (2012). Learning the ABCs of *Bt*: ABC transporters and insect resistance to *Bacillus thuringiensis* provide clues to a crucial step in toxin mode of action. *Pesticide Biochemistry and Physiology*, 104(2), 103–110. <https://doi.org/10.1016/j.pestbp.2012.05.007>
- Hernandez, C. S., Andrew, R., Bel, Y., & Ferre, J. (2005). Isolation and toxicity of *Bacillus thuringiensis* from potato-growing areas in Bolivia. *Journal of Invertebrate Pathology*, 88(1), 8–16. <https://doi.org/10.1016/j.jip.2004.10.006>
- ITIS. (2022). *Spodoptera exigua*. (<https://www.itis.gov/servlet/SingleRpt/SingleRpt#null>). Diakses tanggal 01 Juni 2021, pukul 08.00 WIB.
- ITIS. (2022). *Bacillus thuringiensis*. (<https://www.itis.gov/servlet/SingleRpt/SingleRpt#null>). Diakses tanggal 01

Juni 2021, pukul 08.00 WIB.

- ITIS. (2022). *Samia ricini*. ([https://www.itis.gov/servlet/SingleRpt/SingleRpt?search\\_topic=TSN&search\\_value=936212#null](https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=936212#null)). Diakses tanggal 01 Juni 2021, pukul 08.00 WIB.
- Jadhav, Y. Z., Deshpande, S. D., & Sindhikar, A. A. (2018). Manufacturing of sulphated castor oil (Turkey red oil) by sulphonation process. 5(7), 6384–6389. <https://doi.org/10.13140/RG.2.2.10604.00645>
- Jayanthi, P. D. K., & Padmavathamma, K. (2001). Joint action of microbial and chemical insecticides on *Spodoptera Litura* (Fab.) (Lepidoptera: Noctuidae). *Journal of Tropical Agriculture*, 39, 142–144.
- Kaur, J., Rajkhowa, R., Tsuzuki, T., Millington, K., Zhang, J., & Wang, X. (2013). Photoprotection by silk cocoons. *Biomacromolecules*, 14(10), 3660–3667. <https://doi.org/10.1021/bm401023h>
- Kawaguchi, Y., Ichida, M., Kusakabe, T., & Koga, K. (2000). Chorion morphology of the Eri-silkworm, *Samia cynthia ricini* (Donovan) (Lepidoptera: Saturniidae). *Applied Entomology and Zoology*, 35(4), 427–434. <https://doi.org/10.1303/aez.2000.427>
- Khetan, S. K. (2001). Microbial Pest Control. *Journal of Phytopathology*. 149, 491–492. <https://doi.org/10.1201/9781482270631>
- Kumar, J. P., & Mandal, B. B. (2019). The inhibitory effect of silk sericin against ultraviolet-induced melanogenesis and its potential use in cosmeceutics as an anti-hyperpigmentation compound. *Photochemical and Photobiological Sciences*, 18(10), 2497–2508. <https://doi.org/10.1039/c9pp00059c>
- Kunz, R. I., Brancalhão, R. M. C., Ribeiro, L. D. F. C., & Natali, M. R. M. (2016). Silkworm sericin: Properties and biomedical applications. *BioMed Research International*. <https://doi.org/10.1155/2016/8175701>
- Kwon, M., Cho, H. M., & Ahn, Y. J. (2006). Relationship between feeding damage by Beet armyworm, *Spodoptera exigua* (Lepidoptera: Noctuidae) and leaf trichome density of Potato. *Journal of Asia-Pacific Entomology*, 9(4), 361–367. [https://doi.org/10.1016/S1226-8615\(08\)60315-5](https://doi.org/10.1016/S1226-8615(08)60315-5)
- Lukmawati, D. (2020). Efektivitas ekstrak kokon *Samia ricini* (Drury, 1773) terhadap patogenitas *Bacillus thuringiensis* var. *kurstaki* Berl. sebagai pengendali ulat grayak *Spodoptera litura* (Fabricius, 1775) pada skala laboratorium. Universitas Gadjah Mada. Skripsi. Universitas Gadjah Mada, Yogyakarta.
- Lantang, D. (2010). Toksisitas isolat lokal *Bacillus thuringiensis* (H-14) serta lama efektivitasnya di dalam air terhadap larva nyamuk *Anopheles farauti* Laveran. *Jurnal Biologi Papua*, 2(2), 53–56. <https://doi.org/10.31957/jbp.562>
- Lee, J., Kiuchi, T., Kawamoto, M., Shimada, T., & Katsuma, S. (2018). Accumulation of uric acid in the epidermis forms the white integument of *Samia ricini* larvae. *Plos One*. 13(10): e0205758. <https://doi.org/10.1371/journal.pone.0205758>
- Loutfi, H., Fayad, N., Pellen, F., Jeune, B. Le, Chakroun, M., Benfarhat, D., Lteif, R., Kallassy, M., Brun, G. Le, & Abboud, M. (2021). Morphological study of *Bacillus thuringiensis* crystals and spores. *Applied Sciences (Switzerland)*, 11(1), 1–15. <https://doi.org/10.3390/app11010155>



- Marsadi, D., Supartha, I. W., & Sunari, A. A. A. S. (2017). Invasi dan tingkat serangan ulat bawang (*Spodoptera exigua* Hubner) pada dua kultivar Tanaman bawang merah di Desa Songan, Kecamatan Kintamani, Kabupaten Bangli. *Jurnal Agroekoteknologi Tropika*, 6(4), 360–369. <https://ojs.unud.ac.id/index.php/JAT>
- Masahiro, S., Yamada, H. & Kato, N. (2000). Consumption of silk protein, sericin elevates intestinal absorption of Zinc, Iron, Magnesium and Calsium in rats. 20(10), 1505–1511. [https://doi.org/10.1016/S0271-5317\(00\)80031-7](https://doi.org/10.1016/S0271-5317(00)80031-7)
- Mehta, V., Jayaram, C. S., Koranga, R., & Negi, N. (2021). Developmental biology of *Spodoptera exigua* (Hubner) (Lepidoptera : Noctuidae) on tomato under mid hills (sub-humid) conditions of India. *Biological Forum-An International Journal*. 13(3a), 11-15.
- Mondal, M., Trivedy, K., & Kumar, S. N. (2007). The silk proteins, sericin and fibroin in silkworm, *Bombyx mori* Linn: A review. *Caspian Journal of Environmental Sciences*. 5(2), 63-76.
- Nickerson, K.W. & L.A. Bulla. (1974). Physiology of spore-forming bacteria associated with insects, minimal nutrition requirements for growth, sporulation and parasporal crystal formation of *Bacillus thuringiensis*. *Appl. Microbiol.* 28(1):124-8. <https://doi.org/10.1128/am.28.1.124-128.1974>
- Padamwar, M. N., & Pawar, A. P. (2004). Silk sericin and its applications: A review. *Journal of Scientific and Industrial Research*. 63(4), 323–329.
- Palma, L., Munoz, D., Berry, C., Murillo, J. & Caballero, P. (2014). *Bacillus thuringiensis* Toxins: An overview of their biocidal activity. 3296–3325. <https://doi.org/10.3390/toxins6123296>
- Park, Y., & Kim, Y. (2013). RNA interference of cadherin gene expression in *Spodoptera exigua* reveals its significance as a specific *Bt* target. *Journal of Invertebrate Pathology*, 114(3), 285–291. <https://doi.org/10.1016/j.jip.2013.09.006>
- Pigott, C. R., & Ellar, D. J. (2007). Role of receptors in *Bacillus thuringiensis* crystal toxin activity . *Microbiology and Molecular Biology Reviews*, 71(2), 255–281. <https://doi.org/10.1128/mmb.00034-06>
- Pozsgay, M., Fast, P., Kaplan, H., & Carey, P. R. (1987). The effect of Sunlight on the protein crystals from *Bacillus thuringiensis* var. *kurstaki* HD 1 and NRD12: A Raman Spectroscopic Study. *Journal of Invertebrate Pathology*. 50(3), 246–253. [https://doi.org/10.1016/0022-2011\(87\)90089-9](https://doi.org/10.1016/0022-2011(87)90089-9)
- Purnamasari, I. (2021). Karakteristik kokon ngengat sutera *Samia cynthia ricini* (biosduval) (Lepidoptera: Saturniidae) dengan pemberian pakan berbeda. Institut Pertanian Bogor. Skripsi. Institut Pertanian Bogor. Bogor
- Pusztai, M., Fast, P., Gringorten, L., Kaplan, H., Lessard, T., & Carey, P. R. (1991). The mechanism of sunlight-mediated inactivation of *Bacillus thuringiensis* crystals. *Biochemical Journal*, 273(1), 43–47. <https://doi.org/10.1042/bj2730043>
- Putrasamedja, S., Setiawati, W., Lukman, L., & Hasyim, A. (2016). Penampilan beberapa klon bawang merah dan bubungannya dengan intensitas serangan organisme pengganggu tumbuhan. *Jurnal Hortikultura*, 22(4), 349. <https://doi.org/10.21082/jhort.v22n4.2012.p349-359>

- Qiu, L., Cui, S., Liu, L., Zhang, B., Ma, W., Wang, X., Lei, C., & Chen, L. (2017). Aminopeptidase N1 is involved in *Bacillus thuringiensis* Cry1Ac toxicity in the beet armyworm, *Spodoptera exigua*. *Scientific Reports*, 7(2), 1–7. <https://doi.org/10.1038/srep45007>
- Qiu, L., Hou, L., Zhang, B., Liu, L., Li, B., Deng, P., Ma, W., Wang, X., Fabrick, J. A., Chen, L., & Lei, C. (2015). Cadherin is involved in the action of *Bacillus thuringiensis* toxins Cry1Ac and Cry2Aa in the beet armyworm, *Spodoptera exigua*. *Journal of Invertebrate Pathology*, 127, 47–53. <https://doi.org/10.1016/j.jip.2015.02.009>
- Rahman, M. M., Roberts, H. L. S., Sarjan, M., Asgarit, S., & Schmidt, O. (2004). Induction and transmission of *Bacillus thuringiensis* tolerance in the flour moth *Ephestia kuehniella*. *Proceedings of the National Academy of Sciences of the United States of America*, 101(9), 2696–2699. <https://doi.org/10.1073/pnas.0306669101>
- Ren, X. L., Chen, R. R., Zhang, Y., Ma, Y., Cui, J. J., Han, Z. J., Mu, L. L., & Li, G. Q. (2013). A *Spodoptera exigua* cadherin serves as a putative receptor for *Bacillus thuringiensis* Cry1ca toxin and shows differential enhancement of Cry1ca and Cry1ac toxicity. *Applied and Environmental Microbiology*, 79(18), 5576–5583. <https://doi.org/10.1128/AEM.01519-13>
- Rusman., Rahmayani, R. F. I. R., & Mukhlis. (2018). Buku ajar kimia larutan. *Syiah Kuala University Press*. Hal.45-52.
- Saeed, S., Sayyed, A. H., & Ahmad, I. (2010). Effect of host plants on life-history traits of *Spodoptera exigua* (Lepidoptera: Noctuidae). *Journal of Pest Science*, 83(2), 165–172. <https://doi.org/10.1007/s10340-009-0283-8>
- Salama, H. S., Foda, M. S., El-Sharaby, A., Matter, M., & Khalafallah, M. (1981). Development of some lepidopterous cotton pests as affected by exposure to sublethal levels of endotoxins of *Bacillus thuringiensis* for different periods. *Journal of Invertebrate Pathology*, 38(2), 220–229. [https://doi.org/10.1016/0022-2011\(81\)90126-9](https://doi.org/10.1016/0022-2011(81)90126-9)
- Sansinenea, E., Salazar, F., Ramirez, M., & Ortiz, A. (2015). An ultra-violet tolerant wild-type strain of melanin-producing *Bacillus thuringiensis*. *Jundishapur Journal of Microbiology*, 8(6). <https://doi.org/10.5812/jjm.20910v2>
- Sarifah, Z., 2020. Pengaruh pemberian ekstrak kokon ulat sutera *Samia ricini* sebagai UV-Protectant pada *Bacillus thuringiensis* untuk pengendalian hama *Spodoptera exigua* di Laboratorium. Universitas Gadjah Mada. Skripsi. Universitas Gadjah Mada, Yogyakarta.
- Sarovart, S., Sudatis, B., Meesilpa, P., Grady, B. P., & Magaraphan, R. (2003). The use of sericin as an antioxidant and antimicrobial for polluted air treatment. *Reviews on Advanced Materials Science*, 5(3), 193–198.
- Satiman, U., Tulung, M., Pelealu, J., Salaki, C. L., Kolondam, B. J., Tallei, T. E., Emran, T. Bin, & Pinaria, A. (2022). Morphology, diversity and phylogenetic analysis of *Spodoptera exigua* (Lepidoptera: Noctuidae) in North Sulawesi by employing partial mitochondrial Cytochrome oxidase 1 gene sequences. *Journal of Advanced Biotechnology and Experimental Therapeutics*, 5(1), 136–147. <https://doi.org/10.5455/JABET.2022.D103>

- Schunemann, R., Knaak, N., & Fiuza, L. M. (2014). Mode of action and specificity of *Bacillus thuringiensis* toxins in the control of caterpillars and stink bugs in soybean culture. <https://doi.org/10.1155/2014/135675>
- Setiyawan, A. I., & Fitasari, E. (2018). Pengaruh perbedaan tiga jenis daun ketela pohon terhadap konsumsi dan konversi pakan ulat sutera *Samia cynthia*. *Ternak tropika Journal of Tropical Animal Production*, 19(1), 32–37. <https://doi.org/10.21776/ub.jtapro.2018.019.01.5>
- Setlow, B., Tautvydas, K. J., & Setlow, P. (1998). Small, acid-soluble spore proteins of the  $\alpha/\beta$  type do not protect the DNA in *Bacillus subtilis* spores against base alkylation. *Applied and Environmental Microbiology*, 64(5), 1958–1962. <https://doi.org/10.1128/aem.64.5.1958-1962.1998>
- Shuqing, L. (2002). Effect of temperature and humidity on the development of *Spodoptera exigua* (Hubner). *Journal Huazhong (Central China)*. Agricultural University. 21(4):352-355.
- Shorey, H. H., & Hale, R. L. (1965). Mass-rearing of the larvae of nine Noctuid species on a simple artificial medium. *Journal of Economic Entomology*, 58(3), 522–524. <https://doi.org/10.1093/jee/58.3.522>
- Silva, S. M. B., Silva-Werneck, J. O., Falcao, R., Gomes, A. C., Fragoso, R. R., Quezado, M. T., Neto, O. B. O., Aguiar, J. B., De Sa, M. F. G., Bravo, A., & Monnerat, R. G. (2004). Characterization of novel Brazilian *Bacillus thuringiensis* strains active against *Spodoptera frugiperda* and other insect pests. *Journal of Applied Entomology*, 128(2), 102–107. <https://doi.org/10.1046/j.1439-0418.2003.00812.x>
- Skudlik, J. O. S., Oprawa, I. Z. P., & Ost, M. A. M. R. (2005). The egg capsule of *Spodoptera Exigua* Hübner, 1808 (Insecta, Lepidoptera, Noctuidae): morphology and ultrastructure. *Zoologica Poloniae*, 1808(10), 25–31.
- Soberon, M., Gill, S. S., & Bravo, A. (2009). Signaling versus punching hole: How do *Bacillus thuringiensis* toxins kill insect midgut cells? *Cellular and Molecular Life Sciences*, 66(8), 1337–1349. <https://doi.org/10.1007/s00018-008-8330-9>
- Soumia, P.S., Karuppaiah, V., Mahajan, V., & Singh, M. (2020). Beet armyworm *Spodoptera exigua*: Emerging threat to Onion production. *The National Academy of Sciences*. <https://doi.org/10.1007/s40009-020-00892-5>
- Sukirno, S., Lukmawati, D., Hanum, S. S. L., Ameliya, V. F., Sumarmi, S., Purwanto, H., Suparmin, S., Sudaryadi, I., Soesilohadi, R. C. H., & Aldawood, A. S. (2022). The effectiveness of *Samia ricini* Drury (Lepidoptera: Saturniidae) and *Attacus atlas* L. (Lepidoptera: Saturniidae) cocoon extracts as ultraviolet protectants of *Bacillus thuringiensis* for controlling *Spodoptera litura* Fab. (Lepidoptera: Noctuidae). *International Journal of Tropical Insect Science*, 42(1), 255–260. <https://doi.org/10.1007/s42690-021-00540-5>
- Sundararajan, B., Moola, A. K., Vivek, K., & Kumari, B. D. R. (2018). Formulation of nanoemulsion from leaves essential oil of *Ocimum basilicum* L. and its antibacterial, antioxidant and larvicidal activities (*Culex quinquefasciatus*). In *Microbial Pathogenesis* (Vol. 125). Elsevier Ltd. <https://doi.org/10.1016/j.micpath.2018.10.017>
- Suparto, J. (2000). Statistik: Teori dan aplikasi. Erlangga. Hal.152-153

- Sutherland, T. D., Young, J. H., Weisman, S., Hayashi, C. Y., & Merritt, D. J. (2010). Insect silk: One name, many materials. *Annual Review of Entomology*, 55, 171–188. <https://doi.org/10.1146/annurev-ento-112408-085401>
- Tampubolon, D., Pangestiniingsih, Y., Zahara, F., & Manik, F. (2013). Uji patogenisitas *Bacillus thuringiensis* dan *Metarhizium anisopliae* terhadap mortalitas *Spodoptera Litura* Fabr (Lepidoptera: Noctuidae) di laboratorium. *Jurnal Agroekoteknologi Universitas Sumatera Utara*, 1(3), 95413. <https://dx.doi.org/10.32734/jaet.v1i3.3004>
- Taylor, J. E., & Riley, D. G. (2008). Artificial infestations of beet armyworm, *Spodoptera exigua* (Lepidoptera: Noctuidae), used to estimate an economic injury level in tomato. *Crop Protection*, 27(2), 268–274. <https://doi.org/10.1016/j.cropro.2007.05.014>
- Tsubota, T., Yamamoto, K., Mita, K., & Sezutsu, H. (2016). Gene expression analysis in the larval silk gland of the eri silkworm *Samia ricini*. *Insect Science*, 23(6), 791–804. <https://doi.org/10.1111/1744-7917.12251>
- Umbanhowar, J., & Hastings, A. (2002). The impact of resource limitation and the phenology of parasitoid attack on the duration of insect herbivore outbreaks. *Theoretical Population Biology*, 62(3), 259–269. <https://doi.org/10.1006/tpbi.2002.1617>
- Vaithanomsat, P., & Kitpreechavanich, V. (2008). Sericin separation from silk degumming wastewater. 59, 129–133. <https://doi.org/10.1016/j.seppur.2007.05.039>
- Wallner, W. E., Dubois, N. R., & Grinberg, P. S. (1983). Alteration of parasitism by *Rogas lymantriae* (Hymenoptera: Braconidae) in *Bacillus thuringiensis*-stressed gypsy moth (Lepidoptera: Lymantriidae) Hosts. *Journal of Economic Entomology*, 76(2), 275–277. <https://doi.org/10.1093/jee/76.2.275>
- Yunianto. (2022). Webinar ento talk X. Kelompok studi entomologi (KSE). Fakultas Biologi. Universitas Gadjah Mada.
- Zhang, Y. Q. (2002). Applications of natural silk protein sericin in biomaterials. *Biotechnology Advances*, 20(2), 91–100. [https://doi.org/10.1016/S0734-9750\(02\)00003-4](https://doi.org/10.1016/S0734-9750(02)00003-4)
- Zhaorigetu, S., Yanaka, N., Sasaki, M., Watanabe, H., & Kato, N. (2003). Inhibitory effects of silk protein, sericin on UVB-induced acute damage and tumor promotion by reducing oxidative stress in the skin of hairless mouse. *Journal of Photochemistry and Photobiology B: Biology*, 71(1–3), 11–17. [https://doi.org/10.1016/S1011-1344\(03\)00092-7](https://doi.org/10.1016/S1011-1344(03)00092-7)
- Zheng, S., Henken, B., Wietsma, W., Sofiari, E., Jacobsen, E., Krens, F. A., & Kik, C. (2000). Development of bio-assays and screening for resistance to beet armyworm (*Spodoptera exigua* Hubner) in *Allium cepa* L. and its wild relatives. *Euphytica*, 114(1), 77–85. <https://doi.org/10.1023/A:1004089424419>
- Zheng, X. L., Cong, X. P., Wang, X. P., & Lei, C. L. (2011a). A review of geographic distribution, overwintering and migration in *Spodoptera exigua* Hübner (Lepidoptera: Noctuidae). *Journal of the Entomological Research Society*, 13(3), 39–48.
- Zheng, X. L., Cong, X. P., Wang, X. P., & Lei, C. L. (2011b). Pupation behaviour,



depth, and site of *Spodoptera exigua*. *Bulletin of Insectology*, 64(2), 209–214.

Zuge, L. C. B., Silva, V. R., Hamerski, F., Ribani, M., Gimenes, M. L., & Scheer, A. P. (2017). Emulsifying properties of sericin obtained from hot water degumming process. *Journal of Food Process Engineering*, 40(1), 1–10. <https://doi.org/10.1111/jfpe.12267>